B 108

Cardiac autonomic function in hand-arm vibration syndrome patients

○M.S. Laskar, M. Iwamoto, T. Wakui, N. Toibana*, N. Harada

Department of Hygiene, Yamaguchi University School of Medicine

*Tokushima Kensei Hospital

Purpose: The purpose of the present study was to investigate non-invasively the cardiac autonomic function in hand-arm vibration syndrome (HAVS) patients and controls of similar age using heart rate variation (HRV) analysis.

Subjects: Twenty one HAVS patients (Patients) and 10 healthy subjects (Controls) volunteered for the study. All patients stopped to be exposed to vibration and were under medical treatment for HAVS. Eleven of the patients were with vibration-induced white finger (VWF (+)) and 10 without VWF (VWF (-)). The means (SD) of age, height and weight were 59.9 (3.7), 160.7 (5.0) and 59.5 (8.2), respectively, for the patients, and 58.3 (4.2), 160.6 (4.9) and 63.0 (8.5), respectively, for the controls. The numbers of current smokers and drinkers in both patients and controls did not differ significantly. The mean (SD) duration of exposure to vibration, time from retirement and time under treatment for the patients were 18.7 (6.7), 4.3 (3.1) and 3.9 (3.0), respectively.

Methods: After a rest for 40 minutes, ECG was recorded during spontaneous breathing in supine resting. Time-domain indexes of HRV (Mean of R-R intervals (MeanRR), standard deviation of R-R intervals (SDRR) and coefficient of variation of R-R intervals (CV)) and frequency-domain indexes of HRV (the normalized low frequency (0.04-0.15 Hz) component (LF, nu (normalized unit)) and normalized high frequency (0.15-0.40 Hz) component (HF, nu) were calculated from two minutes ECG data. The indexes of HRV mentioned above are used to indicate the parasympathetic nervous activity in the supine position without any stimulation.

Results: Table 1 shows HRV in the HAVS patients and controls. The LFs (nu) of the patients were significantly lower than that of the controls ($p < 0.05$). The HFs (nu) of the patients, except for VWF (-), were

significantly lower than that of the controls ($p < 0.05$). When Pearson's correlation analysis was done for the patients using HRV indexes with age, weight, duration of exposure to vibration, time from retirement (TR) and time under treatment (TT), the LF (nu)-TR and LF (nu)-TT correlations were positively and statistically significant ($p < 0.01$). Then, the HAVS patients were divided into three groups as follows according to TR: group A = upto 1 year, group B = from 2 to 4 years and group C = from 5 to 10 years; or according to TT: group X = upto 1 year, group Y = from 2 to 4 years and group Z = from 5 to 10 years (Table 2). The LFs (nu) of the groups A, B, X and Y were significantly lower than that of the controls ($p < 0.05$, $p < 0.01$). The LFs (nu) of the groups B and C were significantly larger than that of the group A ($p < 0.05$). The HFs (nu) of the groups A and X were significantly lower than that of the controls ($p < 0.05$). The HF (nu) of the group Z was significantly larger than that of the group X ($p < 0.05$).

Conclusions: The results of the study indicate decreased cardiac parasympathetic nervous activity in the HAVS patients in comparison with the healthy controls of similar age. The findings of the study also indicate that cessation of exposure to vibration and treatment might have a beneficial effect on the cardiac parasympathetic nervous activity in the HAVS patients.

Table 1 HRV in the HAVS patients and controls

	Patients (n=21)	VWF (+) (n=11)	VWF (-) (n=10)	Controls (n=10)
MeanRR (msec)	894±115	925±135	860±82	885±140
SDRR (msec)	28.0±13.5	31.8±16.2	23.8±8.6	28.4±9.1
CV (%)	3.1±1.3	3.4±1.6	2.8±1.0	3.2±0.9
LF (nu)	13.0±1.5**	13.3±1.3*	12.8±1.8*	14.7±1.7
HF (nu)	27.2±2.7**	27.2±2.7*	27.3±3.0	30.6±4.3

* $P < 0.05$, ** $P < 0.02$ vs controls (Student's *t*-test).**Table 2** HRV in the HAVS patients (n = 21, divided into three groups as follows according to time from retirement: group A = upto 1 year, group B = from 2 to 4 years and group C = from 5 to 10 years; or according to time under treatment: group X = upto 1 year, group Y = from 2 to 4 years and group Z = from 5 to 10 years) and controls

	Group A (n=5)	Group B (n=8)	Group C (n=8)	Group X (n=6)	Group Y (n=8)	Group Z (n=7)	Controls (n=10)
MeanRR (msec)	861±82	879±153	925±94	882±80	866±151	937±94	885±140
SDRR (msec)	21.4±5.8	29.0±12.0	31.1±17.7	25.4±11.2	26.5±10.0	31.8±18.9	28.4±9.1
CV (%)	2.5±0.8	3.3±1.2	3.3±1.7	2.9±1.2	3.0±1.0	3.4±1.9	3.2±0.9
LF (nu)	11.6±1.5**	13.2±0.8*#	13.8±1.6*	11.9±1.6**	13.1±0.9*	14.0±1.6§	14.7±1.7
HF (nu)	25.7±2.9*	26.9±2.7	28.5±2.4	25.7±2.6*	27.0±2.6	28.8±2.4§	30.6±4.3

* $P < 0.05$, ** $P < 0.01$ vs controls; # $P < 0.05$ vs group A; § $P < 0.05$ vs group X (Student's *t*-test). nu, normalized unit.