E107

手腕振動症候群診断のための冷却負荷手指血圧検査における
冷却用ブランケット使用の影響

Laskar MS、Mahbub MH、井上 正岩、横山健治郎、大村 佳代、
原田 規章

Department of Hygiene, Yamaguchi University School of Medicine, Ube, Japan

【Introduction】 Percentage finger systolic blood pressure (%FSBP) response to finger cooling is one of the tests used for assessing vascular components of the hand-arm vibration syndrome (HAVS). The test was introduced in 1977 and has been considered as an International Standard within the International Organization for Standardization after long discussions for optimal provocation and best characterization of an attack of vasospastic Raynaud's phenomenon (ISO 14835-2, 2005). While measuring FSBP on a single-test finger in response to finger cooling by Medimatic apparatus, some researchers used additional body cooling for enhancement of diagnostic significance of the test by placing a cooling blanket perfused with water of about 10°C for 15 minutes on subject body surface in supine position. Simultaneous multi-test- and reference-finger measurements of FSBP can be done to have a large number of data avoiding time-consuming performance by using HVLab Multi-Channel Plethysmograph, University of Southampton, United Kingdom. In the present study, we investigated the effect of additional body cooling on %FSBP response to finger cooling using this multi-test-finger measurement method in healthy subjects.

【Subjects and Methods】 Six healthy male subjects (age: 24.8 ± 5.6 years, BMI: 22.9 ± 1.5 kg/m²) volunteered for the present study. They were in supine position with the hands in neutral position during FSBP measurements. Before measurements, smoking, alcohol intake and time after food intake were controlled. The subjects wore light indoor clothing. All of them rested for at least 30 min in the environment of the examination room without physiological or psychological stress for thermal adaptation. The examination room was kept at $21 \pm 1^\circ\text{C}$. The FSBP measurements were done at 30, 15 and 10°C cuff-water temperatures on the right hand, simultaneously in four test fingers (cooled) and the thumb as reference (not cooled). Three subjects were assigned at first to examination with additional body cooling for 15 minutes by a cooling blanket (Gaymer, USA) perfused with water of 10°C and mounted on subject body surface at measurements using 15 and 10°C cuff-water temperatures. The other three subjects were assigned at first to examination without additional body cooling. The %FSBPs for index, middle, ring and little fingers at 15 and 10°C cuff-water temperatures were calculated. The ISO 14835-2, 2005 was followed to perform the test procedure. Four-way analysis of variance (ANOVA) was performed to determine the independent influence of subject, additional body cooling, test finger and cuff-water temperature factors on %FSBPs response to finger cooling.

【Results】 The results of four-way ANOVA showed that the additional body cooling factor was statistically significant ($p < 0.001$); the mean %FSBP (SD) with and without blanket was 95.2 (23.4) and 85.3 (8.9), respectively. The subject factor was also significant ($p < 0.001$). The subject and additional body cooling interaction factor was also significant ($p < 0.001$).

【Conclusions】 The results of the present study showed that additional body cooling might have an influence on %FSBP response to finger cooling. The effect of additional body cooling may have an implication when assessing peripheral vascular function for diagnosing HAVS using %FSBP response to finger cooling involving the multi-test-finger measurement method.