The Meteorological
Society of
Japan

Scientific Online Letters on the Atmosphere (SOLA)

EARLY ONLINE RELEASE

This is a PDF of a manuscript that has been peer-reviewed and accepted for publication. As the article has not yet been formatted, copy edited or proofread, the final published version may be different from the early online release.

This pre-publication manuscript may be downloaded, distributed and used under the provisions of the Creative Commons Attribution 4.0 International (CC BY 4.0) license. It may be cited using the DOI below.

The DOI for this manuscript is

DOI: 10.2151/sola. 2020-038.

J-STAGE Advance published date: Oct. 22, 2020

The final manuscript after publication will replace the preliminary version at the above DOI once it is available.

**Investigating Aerosol Effects on Maritime Deep Convective Clouds using Satellite
and Reanalysis Data**

Pradeep Khatri, Hiroaki Ooashi, and Hironobu Iwabuchi

*Center for Atmospheric and Oceanic Studies, Graduate School of Science, Tohoku
University, Sendai, Japan*

Corresponding author: Pradeep Khatri, Center for Atmospheric and Oceanic Studies,
Tohoku University, Sendai, Japan, 980-8578, Japan. E-mail:
pradeep.khatri.a3@tohoku.ac.jp.

Abstract

Aerosol effects on deep convective cloud (DCC) have been recognized as one of the complex subjects in climatic studies because of the difficulty in quantifying the sole effect of aerosols on DCC. The complexity further arises if the atmosphere has very strong temporal and spatial variations such as that of Indo-Pacific Warm Pool (IPWP) region. Considering the strong influence of IPWP region on global climate change and water circulation, we investigated aerosol effects on DCC over this region by using data of 2015–2016 El Niño and the 2017–2018 La Niña events. We developed a spectral analysis based framework to identify and decouple the influences of major external factors on aerosol-DCC relationship. We found that temporal variations of aerosols, clouds, and meteorology longer than 2 days' time scale can have larger influences than their diurnal and spatial variations on aerosol-DCC relationship. By removing the effects of those spatial and temporal variations of different scales, the study suggests that aerosols of IPWP region can affect DCC properties with time lags less than ~5 hours and by increasing cloud-top height, cloud coverage, and DCC number concentration with the increase of aerosols.

1. Introduction

The increase of cloud droplets due to the increase of aerosols (Twomey, 1977) can inhibit early rain formation and droplet growth (Albrecht, 1989). Under convective conditions, such delay in precipitation simultaneously increases droplet mobility (Heiblum et al., 2016), leading more water mass aloft in the atmosphere. This results the formation of ice particles by releasing latent heat once the freezing level is reached, and then invigorates cloud vertical development and expansion. This phenomenon, known as Aerosol Invigoration Effect AIE (William et al., 2002; Andreae et al., 2004; Khain et al., 2005,2008; Wang, 2005; van den Heever et al., 2006,2011) can have important roles on convection (e.g., Fan et al., 2018), cloud properties (e.g., Koren et al., 2010), and precipitation (e.g., Rosenfeld et al., 2008).

The Indo-Pacific Warm Pool (IPWP) region is regarded as a natural laboratory for climate research community because of its impact on global climate and water circulation through atmospheric teleconnection. Deep convective clouds (DCCs) over IPWP can expand up to hundreds of kilometers to have climatic impact on a global scale (Houze, 2014), making aerosol–DCC interaction over this region an important aspect in the global climate system. Quantifying aerosol effects on such DCC is an exceptionally challenging task because of the coupling effects of aerosol, radiation, and atmospheric dynamics (Tao et al., 2012). Additionally, the atmospheric variations due to large-scale atmospheric dynamics and radiative processes, i.e., variations of different time scales, such as diurnal, intraseasonal, seasonal, interannual, decadal, and even longer periods (Chang et al., 2004) can add complexity in the study of aerosol-DCC interaction over this region. It is because they can simply overshadow aerosol-DCC interaction occurring in relatively shorter spatial and temporal scales. The aerosol-DCC interaction can be further affected by

atmospheric variations due to small-scale dynamics, i.e., cold pools (e.g., Grant and van den Heever, 2014; Jeon et al., 2018). Since AIE can depend on environmental conditions (e.g., Khain et al., 2005; Fan et al., 2009; Storer et al., 2010), it is important to identify and decouple the influences of dominant natural phenomena for better understanding aerosol-DCC interaction over this region, which is the main goal of this study.

This study is organized as follows: Sections 2 and 3 describe data and methodology; Section 4 provides results and discussion, and Section 5 the conclusion.

2. Data

We used satellite and reanalysis data over the region of 20°S-20°N and 90°E-180°E for selected months (July, October, December, and January) of 2015–2016 El Niño and the 2017–2018 La Niña events. The cloud properties are derived from Integrated Cloud Analysis System ICAS (Iwabuchi et al., 2016) by using Advanced Himawari Imager (AHI) observed infrared signals combined with the ozone and atmospheric profiles of meteorological data from Modern-Era Retrospective Analysis for Research and Application Version-2 MERRA-2 (Gelaro et al., 2017) and incorporating cloud vertical inhomogeneity characteristics (Khatri et al., 2018). The spatial and temporal resolutions of ICAS cloud products are 2-km (at nadir) and 10 minutes, respectively. We further used MERRA-2 aerosol (time-averaged) and meteorological (instantaneous) data. MERRA-2 has a spatial resolution of 0.5°(latitude)x0.625°(longitude) and the temporal resolutions are 1 hour and 3 hours for aerosol and meteorological products. The instantaneous meteorological parameters were interpolated in temporal domain from two pre- and post-adjacent observations; whereas, aerosol data of 1 hour resolution were used.

Both meteorological and aerosol data were linearly interpolated at spatial domain matching with cloud observation.

From high-level cloud products with cloud optical thickness (COT) > 0.1 and cloud-top height (CTH) > 10 km, which are reasonable values for minimum limits of thin anvil COT (e.g., Garnier et al., 2012) and DCC top height (e.g., Chae et al., 2011), DCC was detected using a Detect and Spread technique (Boer and Ramanathan, 1997). Though DCCs were not actually circular in shape, for convenience, an equivalent radius R_{EQ} of DCC was calculated from the total area (A) of DCC by $\sqrt{(A / \pi)}$. DCC number density (N), i.e., DCC counts per unit area (km^2), was calculated by normalizing total DCC counts by total pixel areas corresponding to study region.

Since aerosol number concentration rather than aerosol optical thickness (AOT) can represent cloud condensation nuclei (CCN) concentration more reasonably, we used aerosol index (AI), a product of AOD and Angstrom exponent (AE), as a proxy of aerosol number concentration (Nakajima et al., 2001). Primarily, sea-salt aerosols are dominant over our study region, but aerosols of biomass burning are also profound, especially during October (Eck et al., 2019). Convective Available Potential Energy (CAPE), an indicator of atmospheric instability with higher positive value indicating more unstable atmosphere, was calculated using MERRA-2 data by integrating the buoyancy of air parcel from the level of free convection to the equilibrium level. Vertical wind shear (VWS) was calculated as the difference between the mean low-level (925 and 850 hPa) and high-level (200 hPa) wind speeds (Chakraborty et al., 2016). We further used relative humidity at 850 hPa ($\text{RH}_{850\text{hPa}}$) and updraft velocity at 500 hPa ($\omega_{500\text{hPa}}$) considering the important roles of water vapor of low-level atmosphere on the DCC development and mid-level atmospheric updraft velocity on cloud-top height and anvil cloud expansion.

3. Study method

3.1 Removal of the effects of external factors

Firstly, we binned data to a relatively coarse grid of 1° (latitude) $\times$ 1.25° (longitude) to reasonably undermine the effect of non-located observations of aerosols and clouds in the study of aerosol indirect effect (e.g., Anderson et al., 2003; Avey et al., 2007). For each grid, we performed arithmetic averaging for aerosol and meteorological data, and weighted averaging for cloud properties with cloud coverage as the weight.

Secondly, for each month, we filled the missing data, which may result due to retrieval failure and/or low quality, in temporal and spatial domains by cubic spline interpolation before applying a 3-dimensional fast Fourier transform (FFT). This allowed us to convert the temporal domain of each spatial box into frequency (cycle/day) domain. The frequencies lower than 0.2 cycle/day for aerosol and 0.4 cycle/day for other data (meteorological fields and clouds) were removed by using high-pass filters. They were then transferred back to time series by using an inverse FFT, and data filled up before FFT were removed. Those frequencies mentioned above were removed assuming that they were the results of some meteorological and natural phenomena occurring over the study region as summarized in Supplement I. Thus, our data can correspond the variations associated with relatively short-period local-scale fluctuations.

Thirdly, for each month, we removed the diurnal variations by subtracting hourly averages from the instantaneous values. It is because DCC modification due to solar insolation time and land-sea temperature difference can be prominent even without the effect of aerosols.

Finally, we removed the spatial contrasts by subtracting the mean values of each spatial box from instantaneous values. The spatial contrast can arise depending on the distance of DCC from land surface and/or due to region specific natural phenomenon. The external factors stated onward are low-frequency, diurnal, and spatial variations.

When performing a sensitivity study, we removed the effect(s) of factor(s) that was(were) concerned. For example, anomalies were calculated without excluding the effects of diurnal and low-frequency variations while studying the sole effect of spatial variation.

3.2 Timescale for AIE and composite analysis

DCCs are affected by aerosols and environment via dynamical, microphysical, and radiative processes. Thus, a time delay can be expected for changes in DCC properties due to change in the atmosphere. To understand such time delay, we analyzed DCC properties (y) of observation time t_y with AI data (x) of observation time $t_x = t_y - \Delta t$, where Δt is time difference between cloud and aerosol observations. AIs within $\pm 2\sigma_x$ of the mean value, where σ_x is the standard deviation, were divided into 2000 bins to average AIs of each bin and DCC properties corresponding to them, and then to calculate regression coefficient $\partial y / \partial x$ using a least square regression method. A change in y for a σ_x change in x was calculated as $\Delta y = \partial y / \partial x \sigma_x$.

We further used a composite analysis to study AIE after eliminating the effects of external factors mentioned above. First, we calculated the anomalies of AI within $\pm 2\sigma_x$ of the mean values and DCC properties corresponding to them. For 10 bins of AI anomalies ranging from -0.1 to 0.1 (bin width = 0.02), the corresponding DCC anomalies were used

to calculate the mean and 95% confidence interval values by using a bootstrap method as summarized in Appendix II.

4. Results and Discussion

4.1 Effects of external factors

We performed a sensitivity study by eliminating the effect of each factor outlined in Section 3.1 individually and collectively. Figure 1 shows changes in ΔR_{EQ} for 1- σ change in AI for different time lags between aerosol and cloud observations. The mean R_{EQ} for data of our study was 38.3 km. For the case without eliminating the effect of any factor, ΔR_{EQ} was nearly constant regardless of time lag. This result might be mistakenly interpreted as no response of AI of different time lags on cloud properties. In this case, the external factors, including low-frequency variations with period longer than 48 hours (see Supplement I), simply overshadow aerosol-cloud interaction process of lifetime shorter than 48 hours. When the effect of diurnal variation was removed, ΔR_{EQ} responded with the change of lag hour, suggesting the influence of diurnal variation in aerosol-DCC relationship. The effect of diurnal variation seems to be larger than effects of other factors, which is plausibly due to the strong influence of solar insolation time and land-sea surface temperature difference on diurnal cycle of DCC than that for aerosols. Next, removal of the spatial variation effect can be seen to change the ΔR_{EQ} -lag hour relationship as well. A small peak appeared at lag hour of ~ 6 , indicating that spatial variation was capable to influence aerosol-DCC relationship as well. However, we noted a negative ΔR_{EQ} for such peak value. This negative ΔR_{EQ} , i.e., negative $\partial R_{EQ} / \partial AI * \sigma_{AI}$ (see Section 3.2), is possible if relatively larger R_{EQ} values are affected significantly than smaller R_{EQ} values while

removing spatial variation effects. The spatial mean values of R_{EQ} used to remove spatial variation effect of native R_{EQ} values are non-uniform with larger mean values for regions that have larger native R_{EQ} values. Thus, while removing spatial variation effect, relatively larger native R_{EQ} values can be affected significantly than smaller native R_{EQ} values to give a negative $\partial R_{EQ}/\partial AI$ and then negative ΔR_{EQ} . In contrast, removal of low-frequency variation produced a peak after ~ 3 hours. The dominant DCCs over our study region were with R_{EQ} of ~ 30 -40 km, whose typical lifetime could be ~ 6 hours according to Inoue et al. (2009). Therefore, the noted peak with time lag less than 6 hours seems reasonable, indicating more effectiveness of the removal of low-frequency variation.

We also performed a sensitivity study by removing the low-frequency at first, followed by diurnal and spatial variations, which showed no significant difference in ΔR_{EQ} -lag hour relationship as compared to the removal of low-frequency variation alone. This indicated low-frequency variation as the most important factor among the three considered in this study. The results onward correspond to the removal of low-frequency, diurnal, and spatial variations.

The importance and necessity of removing the effects of such external factors are further emphasized in Figure 2, which shows the correlations among AI and different meteorological parameters. If strong dependencies (correlations) among meteorological parameters exist, one may expect a severe meteorological condition affecting DCC promptly by overshadowing aerosol effect, and vice versa. After removing the external factors, we noted weaker relationships among the meteorological parameters (Figure 2b) than before (Figure 2a). Such a weak meteorological field can be more favorable and reliable for evaluating aerosol effects on DCC. Note that those meteorological parameters correspond to relatively coarse grid giving overall view of relatively larger area.

Especially, ω_{500hpa} gives an overall view on the distribution of both ascending and descending motions. This plausibly explains relatively poor correlation between CAPE and ω_{500hpa} in Figure 2(b).

4.2 Timescale for AIE

Figure 3 shows the differences in DCC properties (ΔR_{EQ} , ΔCTH , and ΔN) for $1-\sigma$ (0.023) change in AI after eliminating the effects of all factors. The mean values of R_{EQ} , CTH , and N for data of this study were 38.3 km, 13.1 km and $1.1 \times 10^{-4} \text{ km}^{-2}$, respectively. The peak values of ΔR_{EQ} , ΔCTH , and ΔN , determined by generating fine resolution data via cubic spline fitting, were for time lags of 1, -1.5 and 4.5 hours, respectively. We speculate that precipitation effect was important for negative time lag for ΔCTH . Kubar and Hartmann (2008) suggested that heavily precipitating clouds have higher CTH than moderately precipitating or non-precipitating high clouds. ΔCTH was derived as $\partial CTH / \partial AI * \sigma_{AI}$ (see Section 3.2). Thus, if precipitating clouds wash out aerosols, a negative time lag is prominent, which can be weakly compensated by possibly positive time lag for non-precipitating clouds to result overall negative time lag in a composite analysis. Gasparini et al. (2019) using a modelling study suggested that clouds start to spread horizontally after reaching the maximum height, indicating that time lag for ΔCTH is shorter than that for ΔR_{EQ} . Though negative in value, our observation-based results are consistent with modelling study of Gasparini et al. (2019). According to Figure 3, we suggest that the optimum time lag for aerosol effects on DCC is less than ~ 5 hours.

4.3 AIE

Figure 4 shows the relationships between AI and DCC properties (R_{EQ} , CTH , and N) anomalies with 95% confidence bands. Figure 4 suggests that aerosols can invigorate DCCs over our study region. Except for exceptionally dominant absorbing aerosols (e.g., Jiang et al., 2018), aerosols can increase CTH and DCC coverage via AIE as discussed in Section 1. Such results could be achieved for our study region after removing the effects of external factors mentioned in Section 4.1. Though AIE in forms of CTH and R_{EQ} increments has been widely studied, AIE in the form of N modification is still limited in number. Xue et al. (2008) suggested that an increase of aerosols can increase N of shallow cumulus clouds using large eddy simulation. To the best of our knowledge, there are no observation-based studies to shed further light on aerosol effects on N . Figure 4(c) suggests that aerosol increment also leads to increasing N . Below the baseline (mean) value of AI, the change in N was not significant; however, it changed rapidly after the baseline value. Tracking of DCC is difficult because DCC frequently undergoes aggregation and decomposition. Further, tracking of concentrated DCCs can be more difficult than unconcentrated DCCs. Probably, this may lead tracked DCCs broadly into two groups of low and high concentrations. Another speculation is that DCCs might have occurred over the study region by broadly dividing into two groups of relatively low and high densities. We speculate they could be plausible reasons for rapid change of N after the baseline value of AI in Figure 4(c). We suggest two possible reasons for increasing N with increasing AI. First, precipitation suppression via AIE (see Section 1) can suppress atmospheric stability, generating a favorable condition to seed a new convective cloud and/or to delay the dissipation of existing DCC. Second, an increase in updraft velocity via AIE is also suggested to increase downdraft velocity as a compensation (Zhou et al.,

2017), which can form a cold pool. Such cool pool formation can make it favorable for a next-generation convection (Houze Jr, 2014; Torri et al., 2015; van den Heever et al., 2006; Tao et al., 2007).

5. Conclusions

This study investigated aerosol effects on DCC over IPWP region by identifying and decoupling important factors that could distort relationship between aerosol and DCC properties. The study used data of 2015–2016 El Niño and the 2017–2018 La Niña events. The spectral analysis suggested that clouds, aerosols, and meteorological parameters over the IPWP region were strongly correlated with atmospheric variations of different spatio-temporal scales. Our sensitivity study indicated that such spatio-temporal variations could significantly affect the quantification of aerosol effect on DCC. Specifically, we found that the influence of low-frequency variations with periods longer than 2 days was stronger than that of diurnal and spatial variations. After eliminating the effects of those external factors, we noted the maximum effects of increased aerosols on cloud properties (CTH , R_{EQ} , and N) for time lags less than ~ 5 hours. Those cloud properties were noted to increase with increased AI, suggesting aerosol invigoration effect over the IPWP region.

Acknowledgements

This study is supported by the 1st Research Announcement (RA) on the Earth Observations of the Japan Aerospace Exploration Agency (JAXA; PI no. RA1R306), a Grant-in-Aid for Scientific Research (C) 17K05650 from Japan Society for the Promotion of Science (JSPS), and “Virtual Laboratory for Diagnosing the Earth’s Climate System” program of MEXT, Japan. The Himawari-8 and MERRA-2 data used in this study are obtained from the National Institute of Information and Communication Technology (NICT) and National Aeronautics and Space Administration (NASA), respectively. The authors would like to express sincere thanks to anonymous reviewers for constructive comments.

Supplements

Supplement I: Additional information for study region and data.

Supplement II: Additional information for data analysis method.

References

- Albrecht, B. A., 1989: Aerosols, cloud microphysics, and fractional cloudiness. *Science*, **245**, 1227-1230, doi: 10.1126/science.245.4923.1227.
- Anderson, T. L., R. J. Charlson, D. M. Winker, J. A. Ogren, and K. Holmén, 2003: Mesoscale variations of tropospheric aerosols. *J. Atmos. Sci.*, **60**, 119–136, doi:10.1175/1520-0469(2003)060%3C0119:MVOTA%3E2.0.CO;2.
- Andreae, M. O., D. Rosenfeld, P. Artaxo, A. A. Costa, G. P. Frank, K. M. Longo, and M. A. F. Silva-Dias, 2004: Smoking rain clouds over Amazon. *Science*, **303**, 1337-1342, doi:10.1126/science.1092779.
- Avey, L., T. J. Garrett, and A. Stohl, 2007: Evaluation of the aerosol indirect effect using satellite, tracer transport model, and aircraft data from the International Consortium for Atmospheric Research on Transport and Transformation. *J. Geophys. Res.*, **112**, doi:10.1029/2006JD007581.
- Boer, E. and V. Ramanathan, 1997: Lagrangian approach for deriving cloud characteristics from satellite observations and its implications to cloud parameterization. *J. Geophys. Res.*, **102**, 21383–21399, doi: 10.1029/97JD00930.
- Chakraborty, S., R. Fu, S. T. Massie, and G. Stephens, 2016: Relative influence of meteorological conditions and aerosols on the lifetime of mesoscale convective systems. *Proc. Natl. Acad. Sci.*, **113**, 7426–7431, doi: 10.1073/pnas.1601935113.
- Chae, J. H., D. L. Wu, W. G. Read, and S. C. Sherwood, 2011: The role of tropical deep

convective clouds on temperature, water vapor, and dehydration in the tropical tropopause layer (TTL). *Atmos. Chem. Phys.*, **11**, 3811–3821, doi:10.5194/acp-11-3811-2011, 2011.

Chang, C. P., 2004: *East Asian Monsoon*. World Scientific Series on Meteorology of East Asia, 564pp.

Eck, T.F., B. N. Holben, D. M. Giles, I. Slutsker, A. Sinyuk, J. S. Schafer, A. Smirnov, M. Sorokin, J. S. Reid, A. M. Sayer, N. C. Hsu, Y. R. Shi, R. C. Levy, A. Lyapustin, M. A. Rahman, S. - C. Liew, S. V. S. Cortijo, T. Li, D. Kalbermatter, K. L. Keong, M. E. Yuggotomo, F. Aditya, M. Mohamad, M. Mahmud, T. K. Chong, H. - S. Lim, Y. E. Choon, G. Deranadyan, S. D. A. Kusumaningtyas, and E. Aldrian, 2019: AERONET remotely sensed measurements and retrievals of biomass burning aerosol optical properties during the 2015 Indonesian burning season. *J. Geophys. Res.*, **124**, doi: 10.1029/2018JD030182.

Fan, J, T. Yuan, J. M. Comstock, S. Ghan, A. Khain, L. R. Leung, Z. Li, V. J. Martins, and M. Ovchinnikov, 2009: Dominant role by vertical wind shear in regulating aerosol effects on deep convective clouds. *J. Geophys. Res.*, **114**, doi:10.1029/2009JD012352.

Fan, J., L. R. Leung, D. Rosenfeld, Q. Chen, Z. Li, J. Zhang, and H. Yan ,2013: Microphysical effects determine macrophysical response for aerosol impacts on deep convective clouds. *Proc. Natl. Acad. Sci.*, **110**, E4581–E4590, doi:10.1073/pnas.1316830110.

- 1 Fan, J., D. Rosenfeld, Y. Zhang, S. E. Giangrande, Z. Li, L. A. T. Machado, S. T. Martin,
2 Y. Yang, J. Wang, P. Artaxo, H. M. J. Barbosa, R. C. Braga, J. M. Comstock, Z.
3 Feng, W. Gao, H. B. Gomes, F. Mei, C. Pöhlker, M. L. Pöhlker, U. Pöschl, and
4 R. A. F. de Souza 2018: Substantial convection and precipitation enhancements by
5 ultrafine aerosol particles. *Science*, **359**, 411–418,
6 doi:10.1126/science.aan8461pmid:29371462.
- 7 Garnier, A., J. Pelon, P. Dubuisson, M. Faivre, O. Chomette, N. Pascal, and D. P. Kratz,
8 2012: Retrieval of cloud properties using CALIPSO imaging infrared radiometer.
9 Part I: Effective emissivity and optical depth. *J. Appl. Meteorol. Climatol.*, **51**,
10 1407–1425, doi:10.1175/JAMC-D-11-0220.1.
- 11 Gasparini, B., P. N. Blossey, D. L. Hartmann, G. Lin, and J. Fan: 2019: What Drives the
12 Life Cycle of Tropical Anvil Clouds? *J. Adv. Model. Earth Syst.*, **11**, 2586–2605, doi:
13 10.1029/2019MS001736.
- 14 Gelaro, R., W. McCarty, M. J. Suárez, R. Todling, A. Molod, L. Takacs, C. A. Randles,
15 A. Darmenov, M. G. Bosilovich, R. Reichle, et al., 2017: The modern-era
16 retrospective analysis for research and applications, version 2 (MERRA-2). *J. Clim.*,
17 **30**, 5419–5454, doi: 10.1175/JCLI-D-16-0758.1.
- 18 Grant, L. D. and S. C. van den Heever, 2014: Aerosol - cloud - land surface interactions
19 within tropical sea breeze convection. *J. Geophys. Res.*, **119**, 8340–8361,
20 doi:10.1002/2014JD021912.

- Heiblum, R. H., Altaratz, O., Koren, I., Feingold, G., Kostinski, A. B., Khain, A. P., Ovchinnikov, M., Fredj, E., Dagan, G., Pinto, L., Yaish, R., and Chen, Q., 2016: Characterization of cumulus cloud fields using trajectories in the center of gravity versus water mass phase space: 2. Aerosol effects on warm convective clouds. *J. Geophys. Res.*, **121**, 6356–6373, doi: 10.1002/2015JD024193.
- Houze Jr, R. A., 2014: *Cloud Dynamics*, Second Edition. Academic Press, 496 pp.
- Inoue, T., D. Vila, K. Rajendran, A. Hamada, X. Wu, and L. A. Machado, 2009: Life cycle of deep convective systems over the eastern tropical Pacific observed by TRMM and GOES-W. *J. Meteor. Soc. Japan*, **87**, 381–391. doi: 10.2151/jmsj.87A.381.
- Iwabuchi, H., M. Saito, Y. Tokoro, N. S. Putri, and M. Sekiguchi, 2016: Retrieval of radiative and microphysical properties of clouds from multispectral infrared measurements. *PEPS*, **3**, doi:10.1186/s40645-016-0108-3.
- Jeon, Y.-L., S. Moon, H. Lee, J.-J. Baik, and J. Lkhamjav, 2018: Non-monotonic dependencies of cloud microphysics and precipitation on aerosol loading in deep convective clouds: A case study using the WRF model with bin microphysics. *Atmosphere*, **9**, doi:10.3390/atmos9110434.
- Khain, A. P., D. Rosenfeld, and A. Pokrovsky, 2005: Aerosol impact on the dynamics and microphysics of deep convective clouds. *Q. J. R. Meteorol. Soc.*, **131**, 2639–2663, doi:10.1256/qj.04.62.
- Khain, A. P., N. BenMoshe, and A. Pokrovsky, 2008: Factors Determining the Impact of Aerosols on Surface Precipitation from Clouds: An Attempt at Classification. *J. Atmos. Sci.*, **65**, 1721–1748, doi:10.1175/2007JAS2515.1.

- 1 Khatri, P., H. Iwabuchi, and M. Saito, 2018: Vertical profiles of ice cloud microphysical
2 properties and their impacts on cloud retrieval using thermal infrared measurement.
3 *J. Geophys. Res.*, **123**, doi: 10.1029/2017JD028165.
- 4 Koren, I., G. Feingold, and L. Remer, 2010: The invigoration of deep convective
5 clouds over the Atlantic: aerosol effect, meteorology or retrieval artifact? *Atmos.*
6 *Chem. Phys.* **10**, 8855–8872, doi: 10.5194/acp-10-8855-2010.
- 7 Kubar, T. L. and D. L. Hartmann, 2008: Vertical structure of tropical oceanic convective
8 clouds and its relation to precipitation. *Geophys. Res. Lett.*, **35**, L03804. doi:
9 10.1029/2007GL032811.
- 10 Nakajima, T., A. Higurashi, K. Kawamoto, and J. E. Penner, 2001: A possible correlation
11 between satellite-derived cloud and aerosol microphysical parameters. *Geophys. Res.*
12 *Lett.*, **28**, 1171–1174. doi: 10.1029/2000GL012186.
- 13 Rosenfeld, D., U. Lohmann, G. B. Raga, C. D. O’Dowd, M. Kulmala, S. Fuzzi, A.
14 Reissell, and M. O. Andreae ,2008: Flood or drought: how do aerosols affect
15 precipitation? *Science*, **321**, , 1309–1313,doi: 10.1126/science.1160606.
- 16 Storer, R. L., S. C. van Den Heever, and G. L. Stephens, 2010: Modeling aerosol impacts
17 on convective storms in different environments. *J. Atmos. Sci.*, **67**, 3904–3915, doi:
18 10.1175/2010JAS3363.1.
- 19 Tao, W. -K, X. Li, A. Khain, T. Matsui, S. Lang, and J. Simpson, 2007: Role of
20 atmospheric aerosol concentration on deep convective precipitation: cloud-resolving
21 model simulations. *J. Geophys. Res.*, **112**, doi: 10.1029/2007JD008728

- 1 Tao,W, -K, J. P, Chen, Z. Li, C. Wang, and C. Zhang, 2012: Impact of aerosols on
2 convective clouds and precipitation. *Rev. Geophys.*, **50**,
3 doi:10.1029/2011RG000369.
- 4 Torri, G., Z. Kuang, and Y. Tian, 2015: Mechanisms for convection triggering by cold
5 pools. *Geophys. Res. Lett.*, **42**, 1943-1950. doi: 10.1002/2015GL063227.
- 6 Twomey, S. ,1977: The Influence of Pollution on the Shortwave Albedo of Clouds. *J.*
7 *Atmos. Sci.*, **34**, 1149–1152, doi:10.1175/1520-
8 0469(1977)034<1149:TIOPOT>2.0.CO;2.
- 9 van den Heever, S. C., G. G. Carrió, W. R. Cotton, P. J. DeMott, and A. J. Prenni, 2006:
10 Impacts of Nucleating Aerosol on Florida Storms. Part I: Mesoscale Simulations. *J.*
11 *Atmos. Sci.*, **63**, 1752–1775, doi:10.1175/JAS3713.1.
- 12 van den Heever, S. C., G. L. Stephens, and N. B. Wood, 2011: Aerosol Indirect Effects
13 on Tropical Convection Characteristics under Conditions of Radiative–Convective
14 Equilibrium. *J. Atmos. Sci.*, **68**, 699–718, doi:10.1175/2010JAS3603.1.
- 15 Wang, C., 2005: A modeling study of the response of tropical deep convection to the
16 increase of cloud condensation nuclei concentration: 1. Dynamics and microphysics.
17 *J. Geophys. Res.*, **110**, 1–16, doi:10.1029/2004JD005720.
- 18 Williams, E., D. Rosenfeld, N. Madden, J. Gerlach, N. Gears, L. Atkinson,
19 N. Dunnemann, G. Frostrom, M. Antonio, B. Biazon, R. Camargo, H.
20 Franca, A. Gomes, M. Lima, R. Machado, S. Manhaes, L. Nachtigall, H.
21 Piva, W. Quintiliano, L. Machado, P. Artaxo, G. Roberts, N. Renno , R.
22 Blakeslee, J. Bailey, D. Boccippio, A. Betts, D. Wolff, B. Roy, J.
23 Halverson, T. Rickenbach, J. Fuentes, and E. Avelino, 2002: Contrasting

convective regimes over the Amazon: Implications for cloud electrification. *J.*

Geophys. Res., **107**, doi:10.1029/2001JD000380.

Xue, H., G. Feingold, and B. Stevens, 2008: Aerosol effects on clouds, precipitation, and

the organization of shallow cumulus convection. *J. Atmos. Sci.*, **65**, 392–406, doi:

10.1175/2007JAS2428.1.

Zhou, X., N. Bei, H. Liu, J. Cao, L. Xing, W. Lei, L. T. Molina, and G. Li, 2017: Aerosol

effects on the development of cumulus clouds over the Tibetan Plateau. *Atmos. Chem.*

Phys., **17**, 7423–7434, doi: 10.5194/acp-2017-148.

List of Figure Captions

Figure 1. Changes in equivalent radius of DCC (R_{EQ}) for $1-\sigma$ change in AI before and after eliminating temporal and spatial variations of aerosols, clouds, and meteorological parameters for combination of AI and cloud data of different time lags. The time lag refers to the difference between cloud and aerosol observation times.

Figure 2. Correlation coefficient (%) among different atmospheric parameters (AI: aerosol index, CAPE: convective available potential energy, RH_{850} : relative humidity at 850 hpa, VWS: vertical wind shear, ω_{500} : updraft velocity at 500 hpa) for (a) before and (b) after removal of temporal and spatial variations in data set.

Figure 3. Differences in cloud-top height (CTH), equivalent radius (R_{EQ}) and number density of DCC for $1-\sigma$ change in AI after eliminating spatial and temporal variations. The time lag refers to the difference between cloud and aerosol observation times.

Figure 4. Relationships of AI anomaly with (a) equivalent radius (R_{EQ}), (b) cloud-top height (CTH), and (c) number density (N) of DCC after eliminating spatial and temporal variations. The data correspond to the time lag of 3 hours. The shaded area shows 95% confidence interval.

Figure 1. Changes in equivalent radius of DCC (R_{EQ}) for $1-\sigma$ change in AI before and after eliminating temporal and spatial variations of aerosols, clouds, and meteorological parameters for combination of AI and cloud data of different time lags. The time lag refers to the difference between cloud and aerosol observation times.

Figure 2. Correlation coefficient (%) among different atmospheric parameters (AI: aerosol index, CAPE: convective available potential energy, RH₈₅₀: relative humidity at 850 hpa, VWS: vertical wind shear, ω₅₀₀: updraft velocity at 500 hpa) for (a) before and (b) after removal of temporal and spatial variations in data set.

Figure 3. Differences in cloud-top height (CTH), equivalent radius (R_{EQ}) and number density of DCC for 1- σ change in AI after eliminating spatial and temporal variations. The time lag refers to the difference between cloud and aerosol observation times.

Figure 4. Relationships of AI anomaly with (a) equivalent radius (R_{EQ}), (b) cloud-top height (CTH), and (c) number density (N) of DCC after eliminating spatial and temporal variations. The data correspond to the time lag of 3 hours. The shaded area shows 95% confidence interval.

Supplement I: Spatio-temporal variations of aerosols, clouds, and meteorological parameters over the study region

Below we briefly highlight the variations of aerosols, clouds, and meteorological parameters over our study region by analyzing one-month data of July 2015. We binned the data with a spatial resolution of 2° (longitude) $\times$ 2.5° (latitude) and a temporal resolution of 1 hour.

S1. Spatial distributions

Figure S1 shows the spatial distributions of meteorological parameters (CAPE, $RH_{850\text{hpa}}$, VWS, and $\omega_{500\text{hpa}}$), DCC properties (DCC fraction and DCC number density), and AI. CAPE and RH_{850} are high around Intertropical Convergence Zone (ITCZ; 10°N , 135°E) and South Pacific Convergence Zone (SPCZ; 10°S , 160°E), indicating unstable and humid atmosphere. Both regions are also characterized by negative values of VWS and $\omega_{500\text{hpa}}$. Similarly, Asian monsoon region (0°N , 90°E) is characterized by high values of CAPE, $RH_{850\text{hpa}}$, and VWS; and negative values of $\omega_{500\text{hpa}}$, indicating unstable atmosphere with large wind shear. The DCC fraction is dominant over ITCZ, SPCZ, and Asian monsoon region. AI is generally high over the Asian monsoon region, but low over ITCZ and SPCZ regions. As suggested by these patterns, aerosols, cloud, and meteorological conditions of our study region have strong spatial variations, especially over ITCZ, SPCZ, and Asian monsoon region.

S2. Spatio-temporal variation

To understand the spatio-temporal variations of aerosol, cloud, and meteorological parameters, we performed spectral analysis in this study. The three-dimensional data (latitude, longitude, and time axes) are converted into two dimensions by taking latitudinal average along longitudinal direction. Since these data have different units, each is normalized by its maximum value. Then a Fast Fourier Transform (FFT) is applied to the time series data of each longitude grid to obtain the power spectrum shown in Figure S2. The longitudes are expressed in zonal wave numbers with a positive (negative) value indicating eastward (westward) propagation. Figure S2 suggests that the time series of meteorological parameters and cloud properties are dominated by low-frequency signals with period (T) > 2.5 days; whereas, AI variation is dominated by signals with $T > 5$ days. For $T > 5$ days, wave propagations both in eastward and westward directions are dominant, suggesting the mixed influences of eastward propagating Kelvin wave and westward propagating equatorial Rossby wave that can have T of ~ 2.5 -20 days and ~ 7 -30 days, respectively (Kiladis et al., 2009). Besides the two waves, the variations associated with ITCZ, SPCZ, and Asian monsoon region also contribute to the spatio-temporal variations on aerosols, clouds, and meteorological parameters (Jiang and Waliser, 2009; Zhao et al., 2018) as presented in Figure S2. Such natural phenomena (external factors) can affect the atmosphere on the scale of thousands of km^2 ; whereas, aerosol-cloud interaction can occur on the scale of hundreds of km^2 . The spatio-temporal variations caused by those external factors can overshadow the aerosol-cloud interaction process, making it important to identify and remove their effects while studying aerosol effects on DCC.

1 **Figures**

2

3 **Figure S1.** Monthly averages of (a) CAPE, (b) RH_{850hpa} , (c) VWS, (d) ω_{500hpa} , (e) DCC

4 fraction, (f) DCC density, and (g) AI.

5

Figure S2. Spatio-temporal spectrum analyses for (a) CAPE, (b) $\text{RH}_{850\text{hpa}}$, (c) VWS, (d) $\omega_{500\text{hpa}}$, (e) DCC fraction, (f) DCC density, and (g) AI. The positive and negative zonal wave numbers denote eastward and westward propagations, respectively.

Supplement II: Bootstrap method for mean values with confidence interval

A bootstrap method is a technique to estimate statistics of a population by sampling data randomly with replacement. The bootstrap method used in this study is described below:

Let us assume a data sample X of n elements, such as $X = \{x_1, x_2, \dots, x_n\}$. From X , we generated m bootstrap samples each with n elements by randomly taking the elements of X with replacement. m in this study was set to 1000. Each bootstrap samples were then averaged to generate m sets of mean and standard deviation values denoted by μ_m and σ_m , respectively. The overall mean (μ) and standard deviation (σ) values were calculated by further averaging the values of μ_m and σ_m , respectively. Finally, the 95% confidence interval was estimated as $\mu \pm z_{\alpha/2} \sigma / \sqrt{m}$, where $\alpha = (1 - 0.95)/2$ and $z_{\alpha/2}$ is the z-score value for $m-1$ degree of freedom.

References

- Jiang, X. and D. E. Waliser, 2009 : Two dominant subseasonal variability modes of the eastern Pacific ITCZ. *Geophys. Res. Lett.*, **36**, doi: 10.1029/2008GL036820.
- Kiladis, G. N., M. C. Wheeler, P. T. Haertel, K. H. Straub, and P. E. Roundy, 2009 : Convectively coupled equatorial waves. *Rev., Geophys.*, **47**, doi:10.1029/2008RG000266.
- Zhao, B., K.-N. Liou, Y. Gu, J. H. Jiang, Q. Li, R. Fu, L. Huang, X. Liu, X. Shi, H. Su, et al., 2018 : Impact of aerosols on ice crystal size. *Atmos. Chem. Phys.*, **18**, 1065–1078, doi: 10.5194/acp-18-1065-2018.