The Meteorological
Society of
Japan

Scientific Online Letters on the Atmosphere (SOLA)

EARLY ONLINE RELEASE

This is a PDF of a manuscript that has been peer-reviewed and accepted for publication. As the article has not yet been formatted, copy edited or proofread, the final published version may be different from the early online release.

This pre-publication manuscript may be downloaded, distributed and used under the provisions of the Creative Commons Attribution 4.0 International (CC BY 4.0) license. It may be cited using the DOI below.

The DOI for this manuscript is

DOI: 10.2151/sola. 2021-038.

J-STAGE Advance published date: Oct. 13, 2021

The final manuscript after publication will replace the preliminary version at the above DOI once it is available.

Estimation of excess deaths during hot summers in Japan

Fumiaki Fujibe¹, and Jun Matsumoto¹

*1 Research Center for Climatology, Department of Geography, Tokyo Metropolitan University,
Hachioji, Japan.*

Corresponding author: Fumiaki Fujibe, Department of Geography, Tokyo Metropolitan University, 1-1 Minami-Osawa, Hachioji, Tokyo 192-0397, Japan. E-mail: ffujibe.bs@gmail.com.

Abstract

Using vital statistics data from 1995 to 2019, the relationships among interannual variations of total mortality, heat-stroke mortality, and temperature in summer were assessed in an attempt to estimate excess deaths in hot summer years in Japan. The number of deaths in July and August increased by 1.1% for each 1°C increment of summer mean temperature over Japan, with an eight-fold larger range of interannual variation than the more narrowly defined heat-stroke deaths. This fact implies that excess deaths due to heat are about eight times more prevalent than heat-stroke deaths and can be on the order of 10,000 in a hot summer year. Analyses by age group and cause of death indicated that excess deaths are largely associated with cardiovascular and respiratory diseases among elderly people.

Introduction

Awareness of heat stroke has increased in Japan as a result of an increasing number of heat-stroke casualties and frequent occurrences of extremely hot summers. There are two main indices of the number of deaths attributable to heat. One is deaths for which heat was determined to be the main cause of death. In the tenth version of the International Statistical Classification of Diseases and Related Health Problems (ICD-10; <https://icd.who.int/browse10/2019/en>), which is a medical classification list by the World Health Organization (WHO), death directly caused by high temperature corresponds to "exposure to excessive natural heat" with a code of X30, and almost equivalently, "effects of heat and light" with a code of T67. In Japan, the annual death toll due to X30 has exceeded 1,000 per year in recent hot summers, with a maximum of 1,731 people in 2010.

The other index of fatalities from heat is the number of excess deaths during a period of high temperature with respect to the average for some other time period. In the 2003 European heat wave, the number of excess deaths in Europe was estimated to be more than 50,000 (Brücker 2005), including 14,800 in France (Fouillet et al. 2006). These include many deaths from diseases such as cardiovascular and respiratory disorders in addition to deaths directly linked to heat. In France, only 2,852 cases among the 11,891 excess deaths in people aged 75 years and over in 2003 were directly linked to heat (Pirard et al.

2005).

It is generally recognized that mortality has a V-shaped dependence on temperature, with a minimum at an "optimum temperature" and increasing trends toward lower and higher temperatures (Honda 2007; Honda et al. 2007, 2014). There have been a large number of studies on mortality affected by high temperature in the world (Hajat and Kosatky 2010; Benmarhnia et al. 2015; Son et al. 2019), including some studies for Japan (Gasparrini et al. 2015; Chung et al. 2015; Lim et al. 2016; Ng et al. 2016; Armstrong et al. 2017; Ma et al. 2018). They have revealed that heat effects are stronger for older people and larger for cardiorespiratory mortality than for other causes of death, with some variability among regions. Although these studies have been designed in elaborate ways, they are based on micro viewpoints, which makes it difficult to provide an answer to the question, How many excess deaths occur in hot summers in Japan? To date, direct estimations of heat-induced excess mortality in Japan have been made only in studies of limited regions and periods (Bai et al. 1995; Qiu et al. 2002).

The aim of the present study was to estimate excess deaths in Japan in recent hot summers based on a macro approach by evaluating the relationships among interannual variations of total mortality, heat-stroke mortality, and temperature in summer.

2. Data and procedure

Monthly data of the number of deaths by gender and 5-year age group from 1995 to 2019 were obtained from the vital statistics website of the Ministry of Health, Labour and Welfare of Japan (<https://www.e-stat.go.jp/stat-search/database?toukei=00450011>). Data by prefecture and simplified cause of death were available from 2009 to 2019. Data of X30 deaths were also obtained from the vital statistics for each year. Annual population data of each prefecture were obtained from the website of the Statistics Bureau of Japan (<https://www.stat.go.jp/data/jinsui/2.html>). Using these data, the numbers of deaths for each year, each prefecture, and each cause of death were adjusted to the population structure of the whole of Japan in 2015 in the same way as was done in Fujibe et al. (2018a, b). The following description is based on the adjusted data unless otherwise stated.

The analysis for temperature was made in the same way as in Fujibe et al. (2018a, b) using hourly data from the network of the Automated Meteorological Data Acquisition System (AMeDAS) operated by the Japan Meteorological Agency (JMA). First, stations were selected on the condition that days with more than four missing records were at most

72 three in any one of the months during the period from 1995 to 2019. Then, the
73 representative temperature of each prefecture was defined as the average of the daily mean
74 temperature over the stations having an elevation of at most 200 m above mean sea level or
75 below the 60th percentile among the stations in the prefecture. The total number of stations
76 used for this analysis was 607. The temperature of a region consisting of more than one
77 prefecture was obtained by the average over the prefectures with a weighting of their
78 population in 2015. The temperature of the entire country of Japan was also calculated in
79 this way.

80 Figure 1a shows the interannual changes of the total number of deaths in July and
81 August (D_{JA}), X30 deaths (D_{X30}), and the average temperature in July and August (T_{JA}) for
82 the entire country of Japan. The value of D_{JA} decreased through the time period because of
83 advancement of the medical condition, whereas D_{X30} displayed an increasing trend. It
84 should be noted that 82% of X30 deaths in Japan occur in July to August (Fujibe et al.
85 2018a), but more total deaths occur in cold seasons than in other seasons (e.g., Nakaji et al.

Fig.1

2004). It should also be noted that the number of deaths without population adjustment increased through the analysis period, from 138,000 in 1995 to 216,000 in 2019, due to a rapid increase of the elderly population. The ≥ 80 years old population increased from 3.9 million (3.1% of the total population) in 1995 to 11.2 million (8.9%) in 2019.

Figure 1b shows the deviation of D_{JA} , D_{X30} , and T_{JA} from the quadratic regressions (dashed lines in Fig. 1a). The deviations of the three quantities tend to be in phase with one another. The correlation between the deviations of D_{JA} and D_{X30} is 0.66, and that between D_{JA} and T_{JA} is 0.46. These are statistically significant at the 1% and 5% level, with a two-sided p -value (hereafter p) of < 0.001 and 0.025, respectively. Hereafter, statistical significance is defined as significance at the 5% level, i.e., $p < 0.05$, unless otherwise stated.

To evaluate the size of the interannual variations of D_{JA} in comparison to those of D_{X30} , a linear regression was applied in the following form:

$$\{D_{JA}(n) - [a D_{X30}(n) + bn + cn^2 + d]\}^2 \longrightarrow \min., \quad (1)$$

where n is the year and a to d are least-squares coefficients. The coefficient a gives the variation of D_{JA} for a unit change of D_{X30} , and b and c account for long-term trends of the two variables in quadratic form. Analysis was performed for all ages and also for age groups: < 20 , $20-40$, $40-60$, $60-80$, and ≥ 80 years old.

The dependence on T_{JA} was evaluated for relative mortality, defined by $M_{JA}(n) = D_{JA}(n)/\langle D_{JA} \rangle$, where $\langle D_{JA} \rangle$ is the average of D_{JA} over the analysis period, using a linear regression:

$$\{M_{JA}(n) - [q T_{JA}(n) + sn + tn^2 + u]\}^2 \longrightarrow \min., \quad (2)$$

where n is the year and q to u are least-squares coefficients. The coefficient q gives the variation of M_{JA} for a 1°C change in T_{JA} , and s and t account for long-term trends. Additionally, Eq. (2) was used to find the relationship between monthly number of deaths and temperature by replacing M_{JA} and T_{JA} by their monthly values.

A similar procedure was applied to the data for each region and the cause of death for the period 2009 to 2019. In this analysis, the term n^2 in Eqs. (1) and (2) was omitted because the data period was short.

3. Results

Figure 2a shows the least-squares coefficient a in Eq. (1) for all ages, each gender, and each age group. For all ages, a has a value of 8.0, indicating that the range of interannual variation of D_{JA} is eight times larger than that of D_{X30} . The difference by gender is small, but a is larger for older groups and is statistically insignificant for groups younger than 40 years old.

Figure 2b shows the least-squares coefficient q in Eq. (2). The value of q for all ages is 1.1%/°C. Differences of q among age groups are small, except for the < 20 years old group, implying that the rate of mortality variation with summer temperature is common to people of adult age, although statistical significance is limited to the ≥ 80 and 60–80 years old groups. However, the variation of the number of deaths with summer temperature is

Fig.2

130 much larger for the older age groups that have higher mortality. A regression for D_{JA}
 131 instead of M_{JA} in Eq. (2) results in a q of 2,614 people per degree, and people ≥ 80 years
 132 old account for 65% of the value, nearly two-thirds of the mortality variation according to
 133 summer temperature.

134 The last five lines of Figs. 2a and 2b give the results obtained by using different options
 135 in the analysis: 1) using a quartic function for Eqs. (1) and (2) instead of a quadratic
 136 function, 2) using data without any population adjustment instead of adjusted data, 3)
 137 excluding the data for 2010, which had a record-breaking hot summer with exceptionally
 138 high D_{JA} and D_{X30} , 4) limiting the period to 2009–2019, and 5) using daily maximum
 139 temperature instead of daily mean temperature when evaluating q . As shown, these
 140 changes hardly affect the a values in comparison to that obtained from the main analysis
 141 (top of Fig. 2a). Likewise, q is hardly affected by options 1), 2), and 5), but analysis 3)
 142 without the 2010 data results in a smaller value of 0.6%/°C that is statistically insignificant
 143 ($p = 0.162$), although an analysis limited to August yielded a value of 1.4%/°C that is

significant at the 1% level ($p = 0.005$; not shown). Additionally, the q obtained for the period 2009 to 2019 in analysis 4) is $1.8\%/^{\circ}\text{C}$, which is larger than the value for 1995 to 2019. Thus, the ratio of the variation of D_{JA} to the variation of D_{X30} seems to be robust to the data sampling method, whereas the dependence of M_{JA} on temperature may vary according to the data period.

Figure 3 shows the monthly values of q for all ages and for the three age groups of ≥ 80 , 60–80, and 40–60 years old. It can be seen that q is positive in summer and negative from late autumn to spring and that it has nearly the same magnitude in all age groups. Thus, mortality increases in a cold winter and a hot summer. However, q values in mid to late winter (January and February) are statistically insignificant due to large uncertainty ranges, implying a contribution from factors other than low temperature, such as influenza (e.g., Kinney et al. 2015; Armstrong et al. 2017). Positive values of q are mostly limited to July and August, in correspondence with the fact that the optimum temperature is higher than the annual mean temperature (Honda et al. 2007).

 A red rectangular box containing the text "Fig.3".

Fig.4

Figure 4 shows a and q values for the ten main causes of death based on data from 2009 to 2019. These ten causes account for 96.2% of the total deaths in this period. The a values are high for circulatory and respiratory diseases and "external causes of morbidity and mortality" (hereafter "external causes"), indicating that they are the main contributors to excess deaths in a hot summer, although the value of a for respiratory diseases is not statistically significant. In comparison, q values for circulatory and respiratory diseases are not so large but are comparable to those of other causes in the lower part of the figure, although q for circulatory diseases is statistically significant. This is because q is defined for the relative mortality in Eq.(2), and does not reflect the total number of deaths from each cause. For neoplasms and senility, a and q are both very small, implying that deaths from these two causes are unaffected by the summer heat. The lower five causes have positive values of a and q , with statistical significance for a of diseases of the genitourinary system, certain infectious and parasitic diseases, and endocrine, nutritional, and metabolic diseases and for q of endocrine, nutritional, and metabolic diseases. Thus, in

addition to major causes, minor causes in the lower part of Fig. 4 can also have excess mortality in a hot summer, although their contribution to the total number of deaths may be small.

The category of "external causes" has a subcategory "other accidents" that includes X30. The a value for this subcategory is 0.94 (not shown), which is near unity, implying that the variation of deaths in this subcategory is mainly due to the variation of X30. The value of q is 47%/°C (not shown), in agreement with the result of Fujibe (2013), who showed that X30 mortality varied by 40–60% for a 1°C variation of T_{JA} . Analysis for each subcategory other than "other accidents" did not yield a statistically significant result, including for "suicide" (X60–X84), for which a is 0.17 ($p = 0.169$). A detailed analysis by Sim et al. (2020) implied that extremely high temperature does not increase the suicide risk in Japan.

Figure 5 shows the values of a and q for each of the nine regions in Japan. The analysis for a was based on unadjusted data of D_{X30} because of the limited availability of

Fig.5

data by prefecture. With large uncertainty ranges, the results show no obvious signal of regional differences, although a and q appear to be smaller in southern regions than in central to northern regions.

4. Summary and remarks

This study obtained two main results, which are summarized below.

1. A statistical analysis over the 25 years from 1995 to 2019 indicated that the range of interannual variation of the number of deaths in July to August (D_{JA}) in Japan is about eight times larger than that of "exposure to excessive natural heat" (D_{X30}). In terms of age, the dependence of D_{JA} on D_{X30} is stronger for older ages.
2. The mortality in July and August (M_{JA}) was found to increase by 1.1% for each 1°C increment of mean temperature in July and August. This dependence is largely contributed by deaths due to circulatory and respiratory diseases in elderly people.

As the range of interannual variability of D_{X30} is 1,000 or more, the factor of eight difference for the variations of D_{JA} and D_{X30} implies on the order of 10,000 excess deaths

in a recent extremely hot summer in Japan. In fact, the D_{JA} values in 2009 and 2010 were 205,000 and 217,000 (175,000 and 191,000 without adjustment), respectively, indicating a difference of 12,000 deaths between the cool summer of 2009 and the extremely hot summer of 2010.

It is difficult to make a direct comparison between the present results and the results of previous studies because of methodological differences. However, the M_{JA} variation of 1.1%/°C using summer mean temperature is in rough agreement with the worldwide review of Hajat and Kosatky (2010), who stated "in almost half of the locations, the risk of mortality increased by between 1% and 3% per 1°C change in high temperature." In terms of regional difference, a number of researchers have reported a larger heat effect in northern regions than in southern regions (Chung et al. 2015; Lim et al. 2016; Ma et al. 2018). Their findings may agree with the implication of a smaller number of excess deaths in southern Japan in the present study (Fig. 5).

Ng et al. (2016) showed that the excess mortality attributable to summer heat in Japan

decreased from 1972–1975 and from 2006–2010, meaning that the mortality increase for a 1°C temperature increment became smaller. This suggests an effect from adapting to the recent warmer climate. Fujibe et al. (2020) demonstrated a decreasing trend of X30 mortality relative to the number of heat-stroke ambulance transports during the 2010s, implying the possibility that enhanced public awareness of heat stroke has prompted early emergency responses that have helped prevent deaths. On the other hand, the number of heat-stroke deaths is increasing due to the increase of elderly people, and the excess mortality rate is still high, especially in a year of hot summer. Further investigation of the climatology of heat-related deaths will facilitate their prevention.

Acknowledgments

The study was supported financially by JSPS KAKENHI (grant numbers 26220202 and 17H06116).

References

- Armstrong, B., M. L. Bell, M. de Sousa Zanotti Stagliorio Coelho, Y.-L. L. Guo, Y. Guo, P. Goodman, M. Hashizume, Y. Honda, H. Kim, E. Lavigne, P. Michelozzi, P. Hilario N. Saldiva, J. Schwartz, M. Scortichini, F. Sera, A. Tobias, S. Tong, C. Wu, A. Zanobetti, A. Zeka, and A. Gasparrini, 2017: Longer-term impact of high and low temperature on mortality: An international study to clarify length of mortality displacement. *Environ. Health Perspect.*, **125**, 107009. doi:10.1289/EHP1756.
- Bai, H., M. N. Islam, H. Kuroki, K. Honda, and C. Wakasugi, 1995: Deaths due to heat waves during the summer of 1994 in Osaka Prefecture, Japan. *J. Leg. Med.*, **49**, 265–274. (in Japanese with English abstract)
- Benmarhnia, T., S. Deguen, J. S. Kaufman, and A. Smargiassi, 2015: Vulnerability to heat-related mortality: A systematic review, meta-analysis, and meta-regression analysis. *Epidemiology*, **26**, 781–793. doi:10.1097/EDE.0000000000000375.
- Brücker, G., 2005: Vulnerable populations: Lessons learnt from the summer 2003 heat waves in Europe. *Eurosurveillance*, **10**, 147.

- 244 Chung, Y., Y.-H. Lim, Y. Honda, Y.-L. Guo, M. Hashizume, M. L. Bell, B.-Y. Chen, and
245 H. Kim, 2015: Mortality related to extreme temperature for 15 cities in northeast Asia.
246 *Epidemiology*, **26**, 255–262. doi:10.1097/EDE.0000000000000229.
- 247 Fouillet, A., G. Rey, F. Laurent, G. Pavillon, S. Bellec, C. Guihenneuc-Jouyaux, J. Clavel,
248 E. Jougl, and D. Hémon. 2006: Excess mortality related to the August 2003 heat wave
249 in France. *Int. Arch. Occup. Environ. Health*, **80**, 16–24. doi:10.1007/s00420-006-0089-
250 4.
- 251 Fujibe, F., 2013: Long-term variations in heat mortality and summer temperature in Japan.
252 *Tenki*, **60**, 371–381 (in Japanese with English abstract).
- 253 Fujibe, F., J. Matsumoto, and H. Suzuki, 2018a: Spatial and temporal features of heat
254 stroke mortality in Japan and their relation to temperature variations, 1999–2014. *Geogr.*
255 *Rev. Japan*, Ser. B, **91**, 17–27. doi:10.4157/geogrevjapanb.91.17.
- 256 Fujibe, F., J. Matsumoto, and H. Suzuki, 2018b: Regional features of the relationship
257 between daily heat-stroke mortality and temperature in different climate zones in Japan.

- 258 *SOLA*, **14**, 144–147. doi:10.2151/sola.2018-025.
- 259 Fujibe, F., J. Matsumoto, and H. Suzuki, 2020: Regional features and temporal variations
260 of heat-stroke ambulance transport rates in Japan: Comparison with mortality. *J. Heat Isl.*
261 *Inst. Int.*, **15**, 1–13. (in Japanese with English abstract)
- 262 Gasparrini, A., Y. Guo, M. Hashizume, E. Lavigne, A. Zanobetti, J. Schwartz, A. Tobias, S.
263 Tong, J. Rocklöv, B. Forsberg, M. Leone, M. De Sario, M. L. Bell, Y.-L. L. Guo, C. Wu,
264 H. Kan, S.-M. Yi, M. de Sousa Zanotti Stagliorio Coelho, P. H. N. Saldiva, Y. Honda, H.
265 Kim, and B. Armstrong, 2015: Mortality risk attributable to high and low ambient
266 temperature: a multicountry observational study. *Lancet*, **386**, 369–375.
267 doi:10.1016/S0140-6736(14)62114-0.
- 268 Hajat, S., and T. Kosatky, 2010: Heat-related mortality: A review and exploration of
269 heterogeneity. *J. Epidemiol. Community Health*, **64**, 753–760.
270 doi:10.1136/jech.2009.087999.
- 271 Honda, Y., 2007: Impact of climate change on human health in Asia and Japan. *Glob.*

- 272 *Environ. Res.*, **11**, 33–38.
- 273 Honda, Y., M. Kabuto, M. Ono, and I. Uchiyama, 2007: Determination of optimum daily
274 maximum temperature using climate data. *Environ. Health Prev. Med.*, **12**, 209–216.
275 doi:10.1265/ehpm.12.209.
- 276 Honda, Y., M. Kondo, G. McGregor, H. Kim, Y.-L. Guo, Y. Hijioka, M. Yoshikawa, K.
277 Oka, S. Takano, S. Hales, and R. S. Kovats, 2014: Heat-related mortality risk model for
278 climate change impact projection. *Environ. Health Prev. Med.*, **19**, 56–63.
279 doi:10.1007/s12199-013-0354-6.
- 280 Kinney, P. L., J. Schwartz, M. Pascal, E. Petkova, A. Le Tertre, S. Medina, and R. Vautard,
281 2015: Winter season mortality: will climate warming bring benefits? *Environ. Res. Lett.*,
282 **10**, 064016. doi:10.1088/1748-9326/10/6/064016.
- 283 Lim, Y.-H., C. E. Reid, Y. Honda, and H. Kim, 2016: Temperature deviation index and
284 elderly mortality in Japan. *Int. J. Biometeor.*, **60**, 991–998, doi:10.1007/s00484-015-
285 1091-x.

- 286 Ma, C., Y. Honda, and T. N. Dang, 2018: Comparison of wet-bulb globe temperature
287 (WBGT) and mean temperature for assessment of heat-related mortality: Evidence from
288 47 Japanese prefectures. *Japan. J. Health Hum. Ecol.*, **84**, 52–72.
289 doi:10.3861/kenko.84.2_52.
- 290 Nakaji, S., S. Parodi, V. Fontana, T. Umeda, K. Suzuki, J. Sakamoto, S. Fukuda, S. Wada,
291 and K. Sugawara, 2004: Seasonal changes in mortality rates from main causes of death
292 in Japan (1970–1999). *Eur. J. Epidemiol.*, **19**, 905–913. doi:10.1007/s10654-004-4695-8.
- 293 Ng, C. F. S., M. Boeckmann, K. Ueda, H. Zeeb, H. Nitta, C. Watanabe, and Y. Honda,
294 2016: Heat-related mortality: Effect modification and adaptation in Japan from 1972 to
295 2010. *Glob. Environ. Change*, **39**, 234–243. doi:10.1016/j.gloenvcha.2016.05.006.
- 296 Pirard, P., S. Vandentorren, M. Pascal, K. Laaidi, A. Le Tertre, S. Cassadou, and M.
297 Ledrans, 2005: Summary of the mortality impact assessment of the 2003 heat wave in
298 France. *Eurosurveillance*, **10**, 153–156.
- 299 Qiu, D., T. Tanihata, H. Aoyama, T. Fujita, Y. Inaba, and. M. Minowa, 2002: Relationship

between a high mortality rate and extreme heat during the summer of 1999 in Hokkaido

Prefecture, Japan. *J. Epidemiol.*, **12**, 254–261. doi:10.2188/jea.12.254.

Sim, K., Y. Kim, M. Hashizume, A. Gasparrini, B. Armstrong, F. Sera, C. F. S. Ng, Y.

Honda, and Y. Chung, 2020: Nonlinear temperature-suicide association in Japan from

1972 to 2015: Its heterogeneity and the role of climate, demographic, and socioeconomic

factors. *Environ. Int.*, **142**, 105829. doi:10.1016/j.envint.2020.105829.

Son, J.-Y., J. C. Liu, and M. L. Bell, 2019: Temperature-related mortality: A systematic

review and investigation of effect modifiers. *Environ. Res. Lett.*, **14**, 073004.

doi:10.1088/1748-9326/ab1cdb.

Figure captions

Fig. 1 (a) Interannual changes of the total number of deaths in July and August (D_{JA}), the X30 deaths (D_{X30}), and the average temperature in July and August (T_{JA}). Dotted lines indicate the quadratic regression. (b) Deviation of each quantity from the quadratic regression.

Fig. 2 (a) Least-squares coefficient a in Eq. (1) and (b) least-squares coefficient q in Eq. (2) obtained for each specified condition. Bars indicate 95% confidence ranges, and p -values are written in digits.

Fig. 3 Monthly values of q for all ages and for three age groups. Vertical bars indicate 95% confidence ranges.

Fig. 4 Same as Fig. 2, but for each cause of death. The corresponding ICD-10 codes and the contributions to the total deaths are indicated in parentheses.

Fig. 5 Same as Fig. 2, but for each region shown in the map at left. The letter "J" denotes the entire nation of Japan. The regional populations in 2015 (in thousands) are indicated in parentheses.

Figure 1 (a) Interannual changes of the total number of deaths in July and August (D_{JA}), the X30 deaths (D_{X30}), and the average temperature in July and August (T_{JA}). Dotted lines indicate the quadratic regression. (b) Deviation of each quantity from the quadratic regression.

Figure 2 (a) Least-squares coefficient a in Eq. (1) and (b) least-squares coefficient q in Eq. (2) obtained for each specified condition. Bars indicate 95% confidence ranges, and p -values are written in digits.

Figure 3 Monthly values of q for all ages and for three age groups. Vertical bars indicate 95% confidence ranges.

Figure 4 Same as Fig. 2, but for each cause of death. The corresponding ICD-10 codes and the contributions to the total deaths are indicated in parentheses.

Figure 5 Same as Fig. 2, but for each region shown in the map at left. The letter "J" denotes the entire nation of Japan. The regional populations in 2015 (in thousands) are indicated in parentheses.