Revision Surgery due to Proximal Junctional Failure and Rod Fracture in Adult

Deformity Surgery at a Single Institution in Japan

Running Head Title

Mechanical complication after adult spinal deformity surgery

Tatsuya Yasuda¹, Yu Yamato^{2,3}, Tomohiko Hasegawa², Go Yoshida², Tomohiro Banno²,

Hideyuki Arima², Shin Oe^{2,3}, Yuki Mihara², Koichiro Ide², Yukihiro Matsuyama¹

¹Department of Orthopaedic Surgery, Iwata City Hospital

²Department of Orthopaedic Surgery, Hamamatsu University School of Medicine

³Division of Geriatric Musculoskeletal Health, Hamamatsu University School of
Medicine

Corresponding Author:

Tatsuya Yasuda, MD

Department of Orthopaedic Surgery, Iwata City Hospital, 512 Okubo, Iwata city,
Shizuoka, 438-8550, Japan

Phone: 81538-38-5000

Fax: 81538-38-5050

E-mail: t.yasuda0820@gmail.com

Conflicts of Interest

Yu Yamato and Shin Oe belong to the donated fund laboratory called Division of Geriatric Musculoskeletal Health. The laboratory funded by below organizations.

Sources of Funding

Medtronic Sofamor Danek Inc.

Japan Medical Dynamic Marketing Inc.

Meitoku Medical Institution Jyuzen Memorial Hospital

We have not received funding from the NIH, HHMI, or others.

Type of Contribution of the Authors

Tatsuya Yasuda wrote and prepared the manuscript, and all authors participated in the study design. All authors have read, reviewed, and approved the article.

Approval Code

The study protocol was approved by the Institutional Review Board of the Hamamatsu University School of Medicine, Shizuoka, Japan (No.14-306).

Informed consent: Informed consent for publication was obtained from all participants in this study.

Acknowledgments

None

Revision Surgery Owing to Proximal Junctional Failure and Rod Fracture in Adult Spinal Deformity Surgery at a Single Institution in Japan

ABSTRACT

Introduction

Proximal junctional failure (PJF) and rod fracture (RF) are the primary reasons for revision surgery after a long corrective fusion for the adult spinal deformity (ASD). However, many recent studies on ASD are multicenter studies from the US and European racial characteristics may differ from those of Asians. Therefore, the risk factors for revision surgery because of PJF and RF after ASD surgery were evaluated in Japanese patients.

Methods

Patients with ASD who underwent corrective surgery from the thoracic vertebrae to the ilium at the authors' institution were reviewed. Demographic, surgical, and radiographic parameters were included in the analysis. Univariate and multivariate regression models were used to analyze the risk factors for PJF and RF.

Results

Two hundred and fifty-nine patients were included in the study. A total of 73 patients (28.1%) required revision surgery because of mechanical complications and 15 patients (5.7%) required revision surgery because of PJF on average 380 days after surgery. In PJF cases, body mass index (BMI) and pelvic tilt were significantly higher ($p = 0.01$, $p = 0.048$, respectively). BMI was an independent risk factor for revision owing to PJF (odds ratio [OR], 1.16; $p = 0.013$). A total

of 49 patients (18.9%) required revision owing to RF on average 867 days after surgery. Three-column osteotomy ($p < 0.001$), significant blood loss ($p = 0.048$), number of fusion segments ($p = 0.023$), absence of lateral lumbar interbody fusion ($p < 0.001$), and sagittal imbalance ($p = 0.033$) were risk factors for revision surgery owing to RF in the univariate analysis. Three-column osteotomy (OR 4.41; $p < 0.001$) and number of fusion segments (OR 1.21; $p < 0.009$) were independent factors for revision surgery owing to RF.

Conclusion

PJF occurred in a relatively early phase (approximately 1 year) after surgery in patients with ASD with high BMI. Conversely, RF occurred approximately 2.5 years after surgery in three-column osteotomy and spinal fusion cases that involved longer fusion range.

Keywords

adult spinal deformity; rod fracture; proximal junctional failure; revision surgery; mechanical complication

1 Introduction

2 Patients with adult spinal deformity (ASD) experience back pain and disability, and their quality of life is significantly
3 worse than that of age-matched patients. However, when adequately indicated, surgery has been shown to improve the
4 quality of life compared with conservative treatment¹. As such, the number of adults undergoing surgery has increased².
5 Additionally, surgery for ASD frequently involves long fusion constructs to achieve optimal sagittal and coronal
6 alignment.

7 Long spinal fusion with iliac screws or sacral ala iliac screws is a recent trend in spinal surgery for ASD³⁻⁵. The
8 advantage of long spinal fusion is the maintenance of appropriate spinal alignment, which resulted in decreased spinal
9 motion. However, loss of spinal motion leads to increased mechanical failure due to concentrated load stress to adjacent
10 levels or implants. Mechanical complications are the topic of interest in spinal deformity surgery. The European Spine
11 Study Group proposed a global alignment and proportion (GAP) score as a risk factor simulator for mechanical
12 complications in patients with ASD⁶.

13 Proximal junctional failure (PJF) and rod fracture (RF) are the primary reasons for revision surgery due to
14 mechanical complications^{7, 8}. Although PJF and RF are mechanical complications, their risk factor are thought to be
15 different. Some of the risk factors for PJF and RF have been reported in recent multicenter studies; however, it can be
16 inferred that patient background, such as race, is diverse and that surgical techniques are often not standardized because
17 of multicenter studies. It would be useful to investigate risk factors for PJF and RF in a group of cases with standardized
18 patient background and surgical technique.

This study evaluated the difference in risk factors for PJF and RF after long spinal fusion for ASD at a single institution in Japan.

Materials and Methods

ASD databases (>20 years and coronal Cobb > 20°, sagittal vertical axis [SVA] > 5 cm, pelvic tilt [PT] > 25°, or thoracic kyphosis > 60°) of the authors' institution were queried for Japanese patients with ASD who underwent spinal fusion from the thoracic vertebrae to the ilium with more than 2 years of follow-up since 2010. Demographic, surgical, and radiological variables were prospectively recorded and analyzed.

Demographic variables included age, sex, and body mass index (BMI). Moreover, total blood loss during surgery, total operative time, number of fusion segments, three-column osteotomy (3-CO), lateral lumbar body fusion (LLIF), and number of rods were evaluated as surgical variables. SVA, PT, pelvic incidence minus lumbar lordosis, the distance between the central sacral vertical line and C7, and scoliosis Cobb angle were measured as radiographic parameters before surgery and at first standing after surgery. The GAP alignment score was entered categorically (proportionate, moderately disproportionate, severely disproportionate) and as a compound score.

Revision surgery was recommended for patients with severe pain or neurological symptoms due to PJF or RF or with a progressive loss of correction. In this series, only cases with revision surgery were included in the PJF and RF groups, respectively. First, revision surgeries and reoperations were identified. To examine potential risk factors for revision surgery for PJF and RF, demographic data, surgical data, radiological parameters, and GAP categories were

compared between patients who had undergone revision surgery and those who did not. Risk factors for revision of PJF and RF were identified using univariate analysis for each variable. The identified potential risk factors were then further examined using multiple logistic regression analyses to isolate independent risk factors for revision surgery.

All statistical analyses were performed with EZR (Saitama Medical Center, Jichi Medical University, Saitama, Japan), a graphical user interface for R (The R Foundation for Statistical Computing, Vienna, Austria). Descriptive and bivariate comparisons of demographic variables were performed between the two groups using the independent *t*-test for continuous variables and Fisher's exact test for categorical variables. The level of statistical significance was set at $p < 0.05$.

Results

Of the 393 patients with ASD included in the database with a 2-year follow-up, 259 underwent fusion from the thoracic vertebrae to the ilium. The average age was 68.4 years, and 83.8% were female. A total of 73 patients (28.1%) required revision surgery owing to mechanical complications (Fig. 1). In patients who underwent revision surgery, the average age was 67.7 years and 83.3% were female. The most common cause of revision was RF in 49 cases, followed by PJF in 15 cases.

Proximal Junctional Failure

A total of 15 patients (5.7%) required revision surgeries because of PJF at 380 days (41-1123 days) after surgery. The variables for patients with and without PJF are summarized in Table 1. Regarding demographic variables, BMI in PJF

cases was significantly higher (25.5 vs. 23.0, $p = 0.01$). There was no significant difference in the surgical variables between the two groups. The upper instrumented vertebra in PJF cases was one case each in T2, T4, and T8, two cases in T9, eight cases in T10, and one case in T12. There was no significant difference in the number of fusion segments between the two groups. Regarding radiographic parameters, PT in PJF cases was significantly higher (25.4% vs. 20.4%, $p = 0.048$). In the GAP category, there were no significant differences between the two groups. BMI was an independent risk factor for revision due to PJF (odds ratio [OR], 1.16; $p = 0.013$) in the logistic regression analysis (Table 2).

Rod Fracture

A total of 33 patients with RF received nonsurgical care because of mild symptoms and 49 patients (18.9%) required revision surgery owing to RF at an average of 867 days (181-1738 days) after surgery. In this series, RF cases with revision surgery were defined as the RF group. The variables for the patients with RF and without RF are summarized in Table 3. There was no significant difference in the demographic variables between the two groups. Significantly higher total blood loss (1905 mL vs. 1568 mL, $p = 0.048$), longer fusion segments (10.1% vs. 9.3%, $p = 0.023$), higher rate of three-column osteotomy, and lower rate of LLIF (63.3% vs. 30.5%, $p < 0.001$) were observed in RF cases. With regard to radiographic parameters, preoperative SVA was significantly higher in the RF group. There were no significant differences between the two groups in the GAP category. Both 3-CO (OR, 4.16; $p < 0.001$) and number of fusion segments (OR, 1.20; $p < 0.012$) were independent factors for revision surgery because of RF (Table 4).

RF occurred at the site of 3-CO in 18 cases, posterolateral fusion in 13 cases, posterior interbody fusion in 13 cases,

LLIF in 1 case, and sacroiliac joint in 4 cases. RF levels (except in cases of 3-CO) were L1–2 in 1 case, L3–4 in 4 cases, L4–5 in 6 cases, L5–S1 in 16 cases, and sacroiliac joint in 4 cases. RF at the site of 3-CO was observed in 2 cases for the three-rod construct and in 16 cases for the two-rod construct. Although additional rods were used in 11 of the 27 cases (posterolateral fusion, posterior interbody fusion, and LLIF), RF occurred outside the range of the additional rod in 10 cases.

Discussion

The revision rate for mechanical failure after ASD surgery was relatively high (28.1%). PJF and RF were the leading causes for mechanical revision surgery and each had different characteristics. PJF occurred in a relatively early phase (approximately one year) after surgery. High BMI was an independent risk factor for PJF. Alternatively, revision surgery for RF was performed approximately 2.5 years after surgery. Extended spinal fusion and 3-CO were independent risk factors for RF.

Many recent studies on adult spinal deformities have been conducted in the US and Europe. Multicenter studies have the advantage of accumulating a large number of cases and eliminating bias in the target population. However, they also have disadvantages, such as a lack of uniformity in surgical methods. The advantage of this series is that the surgical techniques were standardized because of their single-center nature. Furthermore, racial differences in spinal alignment have been reported⁹, and studies that focus on specific racial groups would be more revealing.

Previous studies have described various risk factors for PJF after corrective fusion for ASD¹⁰⁻¹⁵. Damage to the

posterior soft tissues and use of pedicle screw constructs were reported as risk factors for PJF related to surgical technique^{10, 12, 14-16}. In this report, surgical variables were not identified as risk factors. Surgical averaging due to the single-center study design might have caused surgical variables were not risk factors for PJF. In agreement with this study, BMI has been reported in previous studies to be a risk factor for PJF^{17, 18}. Moreover, several studies reported that low bone density was also a risk factor for PJF¹⁹; however, the lack of data is a significant limitation of this study. Based on these results, PJF might be caused by a high load applied to a fragile adjacent fixed end. In cases where surgical techniques are standardized, patient factors, such as BMI and bone density, may be responsible for PJF. Even in recent revisions of the GAP score, BMI and bone density were independent risk factors for mechanical failure²⁰. The importance of BMI and bone density in the prevention of PJF should be noted.

Previous studies have reported many risk factors for RF after ASD surgery. The use of 3-CO was reported as a risk factor for RF, similar to the results of this study^{21, 22}. Additionally, several studies reported that a small diameter (< 6mm) was also a risk factor for RF^{23, 24}. Rod diameter was not analyzed because rods of only 6.00 or 6.35 mm in diameter were used. Based on the results, RF might be caused by a high load applied to an unstable area of the fixation segment in 3-CO cases.

In this study, the number of fusion segments was an independent risk factor for RF. This could be attributed to decrease trunk mobility from the more extended fusion, which increased the load on the rods. Unlike PJF, RF was caused by the influence of surgical planning such as 3-CO and more extended fusion and not by patient factors. Therefore, surgeries should be planned to reduce the range of fusion segments and areas of instability as much as

possible to prevent RF. In this study, the additional rod could not reduce the incidence of RF, which may be because several RFs occurred outside the range of the additional rod. Therefore, additional rods should be included in the range from the thoracic vertebrae to the iliac level^{25, 26}. Moreover, LLIF is a dependent risk factor for RF prevention. LLIF using large-sized spacers was considered biomechanically stable and advantageous for the prevention of RF.

The GAP score is a method used to predict mechanical complications⁶. However, a few recent studies have reported no correlation between the GAP score and mechanical complications^{27, 28}. The results of this study also indicated that the GAP score did not correlate with RF and PJF. Furthermore, ideal alignment is thought to differ depending on age, sex, and race. Thus, the cut-off value of the GAP score should be modified depending on the characteristics of the patient group.

Only surgical revision cases were evaluated in this study, but there were several cases of PJF or RF without revision surgery. The indication for revision surgery with PJF or RF was considered on a case-by-case basis without clear criteria. Thus, the risk factors for PJF and RF in this study might be inaccurate as risk factors for developing complications included non-revision surgery, which is a limitation of the study.

Conclusion

PJF and RF were the leading causes of mechanical complication, and each had different characteristics. PJF occurred in a relatively early phase (approximately 1 year) after surgery in patients with ASD with high BMI. Conversely, revision surgery for RF occurred approximately 2.5 years after surgery in 3-CO and spinal fusion cases that involved

more levels.

References

1. Smith JS, Lafage V, Shaffrey CI, et al. Outcomes of Operative and Nonoperative Treatment for Adult Spinal Deformity: A Prospective, Multicenter, Propensity-Matched Cohort Assessment With Minimum 2-Year Follow-up. *Neurosurgery*. 2016;78(6): 851-61.
2. Sing DC, Khanna R, Shaw JD, et al. Increasing Rates of Surgical Management of Multilevel Spinal Curvature in Elderly Patients. *Spine deformity*. 2016;4(5): 365-72.
3. Emami A, Deviren V, Berven S, et al. Outcome and complications of long fusions to the sacrum in adult spine deformity: luque-galveston, combined iliac and sacral screws, and sacral fixation. *Spine*. 2002;27(7): 776-86.
4. Shen FH, Mason JR, Shimer AL, et al. Pelvic fixation for adult scoliosis. *European spine journal : official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society*. 2013;22 Suppl 2: S265-75. .
5. El Dafrawy MH, Raad M, Okafor L, et al. Sacropelvic Fixation: A Comprehensive Review. *Spine deformity*. 2019;7(4): 509-16.
6. Yilgor C, Sogunmez N, Boissiere L, et al. Global Alignment and Proportion (GAP) Score: Development and Validation of a New Method of Analyzing Spinopelvic Alignment to Predict Mechanical Complications After Adult Spinal Deformity Surgery. *The Journal of bone and joint surgery American volume*. 2017;99(19): 1661-

- 72.
7. Martini C, Langella F, Mazzucchelli L, et al. Revision strategies for failed adult spinal deformity surgery. European spine journal : official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society. 2020;29(Suppl 1): 116-25.
8. Ha AS, Hong DY, Coury JR, et al. Partial Intraoperative Global Alignment and Proportion Scores Do Not Reliably Predict Postoperative Mechanical Failure in Adult Spinal Deformity Surgery. Global spine journal. 2020: 2192568220935438.
9. Arima H, Dimar JR, 2nd, Glassman SD, et al. Differences in lumbar and pelvic parameters among African American, Caucasian and Asian populations. European spine journal : official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society. 2018;27(12): 2990-8.
10. Kim JS, Phan K, Cheung ZB, et al. Surgical, Radiographic, and Patient-Related Risk Factors for Proximal Junctional Kyphosis: A Meta-Analysis. Global spine journal. 2019;9(1): 32-40. .
11. Iyer S, Lovecchio F, Elysee JC, et al. Posterior Ligamentous Reinforcement of the Upper Instrumented Vertebrae +1 Does Not Decrease Proximal Junctional Kyphosis in Adult Spinal Deformity. Global spine journal. 2020;10(6): 692-9.
12. Line BG, Bess S, Lafage R, et al. Effective Prevention of Proximal Junctional Failure in Adult Spinal Deformity Surgery Requires a Combination of Surgical Implant Prophylaxis and Avoidance of Sagittal

- 1 Alignment Overcorrection. Spine. 2020;45(4): 258-67.
- 2 13. Hyun SJ, Lee BH, Park JH, et al. Proximal Junctional Kyphosis and Proximal Junctional Failure Following
- 3 Adult Spinal Deformity Surgery. Korean J Spine. 2017;14(4): 126-32.
- 4 14. Yagi M, King AB, Boachie-Adjei O. Incidence, risk factors, and natural course of proximal junctional
- 5 kyphosis: surgical outcomes review of adult idiopathic scoliosis. Minimum 5 years of follow-up. Spine.
- 6 2012;37(17): 1479-89.
- 7 15. Cammarata M, Aubin CE, Wang X, et al. Biomechanical risk factors for proximal junctional kyphosis: a
- 8 detailed numerical analysis of surgical instrumentation variables. Spine. 2014;39(8): E500-7.
- 9 16. Hassanzadeh H, Gupta S, Jain A, et al. Type of Anchor at the Proximal Fusion Level Has a Significant Effect
- 10 on the Incidence of Proximal Junctional Kyphosis and Outcome in Adults After Long Posterior Spinal Fusion.
- 11 Spine deformity. 2013;1(4): 299-305.
- 12 17. Bridwell KH, Lenke LG, Cho SK, et al. Proximal junctional kyphosis in primary adult deformity surgery:
- 13 evaluation of 20 degrees as a critical angle. Neurosurgery. 2013;72(6): 899-906.
- 14 18. Kim YJ, Bridwell KH, Lenke LG, et al. Proximal junctional kyphosis in adult spinal deformity after segmental
- 15 posterior spinal instrumentation and fusion: minimum five-year follow-up. Spine. 2008;33(20): 2179-84.
- 16 19. Yagi M, Fujita N, Tsuji O, et al. Low Bone-Mineral Density Is a Significant Risk for Proximal Junctional
- 17 Failure After Surgical Correction of Adult Spinal Deformity: A Propensity Score-Matched Analysis. Spine.
- 18 2018;43(7): 485-91.

20. Noh SH, Ha Y, Obeid I, et al. Modified global alignment and proportion scoring with body mass index and bone mineral density (GAPB) for improving predictions of mechanical complications after adult spinal deformity surgery. *The spine journal : official journal of the North American Spine Society*. 2020;20(5): 776-84.
21. Smith JS, Shaffrey E, Klineberg E, et al. Prospective multicenter assessment of risk factors for rod fracture following surgery for adult spinal deformity. *Journal of neurosurgery Spine*. 2014;21(6): 994-1003.
22. Smith JS, Shaffrey CI, Ames CP, et al. Assessment of symptomatic rod fracture after posterior instrumented fusion for adult spinal deformity. *Neurosurgery*. 2012;71(4): 862-7.
23. Lertudomphonwanit T, Kelly MP, Bridwell KH, et al. Rod fracture in adult spinal deformity surgery fused to the sacrum: prevalence, risk factors, and impact on health-related quality of life in 526 patients. *The spine journal : official journal of the North American Spine Society*. 2018;18(9): 1612-24.
24. Akazawa T, Kotani T, Sakuma T, et al. Rod fracture after long construct fusion for spinal deformity: clinical and radiographic risk factors. *Journal of orthopaedic science : official journal of the Japanese Orthopaedic Association*. 2013;18(6): 926-31.
25. Yamato Y, Hasegawa T, Kobayashi S, et al. Treatment strategy for rod fractures following corrective fusion surgery in adult spinal deformity depends on symptoms and local alignment change. *Journal of neurosurgery Spine*. 2018;29(1): 59-67.
26. Yamato Y, Hasegawa T, Togawa D, et al. Long additional rod constructs can reduce the incidence of rod

fractures following 3-column osteotomy with pelvic fixation in short term. Spine deformity. 2020;8(3): 481-90.

27. Kawabata A, Yoshii T, Sakai K, et al. Identification of Predictive Factors for Mechanical Complications After Adult Spinal Deformity Surgery: A Multi-Institutional Retrospective Study. Spine. 2020;45(17): 1185-92.

28. Kwan KYH, Lenke LG, Shaffrey CI, et al. Are Higher Global Alignment and Proportion Scores Associated With Increased Risks of Mechanical Complications After Adult Spinal Deformity Surgery? An External Validation. Clinical orthopaedics and related research. 2021;479(2): 312-20.

Figure Legends

Figure 1

Flowchart of patient inclusion criteria. A total of 259 patients were included in this study and 73 patients underwent revision surgery owing to mechanical complications.

Figure 1

233x154mm (150 x 150 DPI)

Table 1. Comparison of PJF and non-PJF

		Non-PJF		PJF		
N		244	(%, SD)	15	(%, SD)	p-value
Demographic Variables						
Age		68.45	9.58	67.47	11.26	0.704
Gender	F	206	84.4	11	73.3	0.276
	M	38	15.6	4	26.7	
BMI		22.97	3.7	25.53	3.97	0.01
Surgical Variables						
Total blood loss(ml)		1635.43	1060.2	1605.87	1107.16	0.917
Total surgical time(min)		440.98	83.14	423.67	103.98	0.441
No. of fusion segments		9.48	2.28	9.07	2.58	0.504
Three-column osteotomy		88	36.1	7	46.7	0.419
LLIF		86	35.2	4	26.7	0.587
No. of rods	2	135	55.8	11	73.3	0.278
	3	32	13.2	0	0.0	
	4	75	31	4	26.7	
Preoperative Radiographic Parameters						
PI-LL(°)		34.52	17.10	28.60	22.16	0.202
SVA(mm)		128.59	82.59	125.34	87.11	0.886
PT(°)		35.86	10.88	34.47	14.93	0.639
C7CSVL(mm)		31.35	32.41	33.05	35.12	0.849
Cobb(°)		29.27	21.61	27.67	24.73	0.576
Postoperative Radiographic Parameters						
PI-LL(°)		7.57	12.91	11	17.47	0.331
SVA(mm)		49.11	41.43	44.95	47.27	0.709
PT(°)		20.37	9.43	25.4	10.26	0.047
C7CSVL(mm)		17.73	17.37	20.93	20.5	0.508
Cobb(°)		9.42	8.99	6.79	6.19	0.282
Global Alignment and Proportional (GAP) Parameters						
GAP score		5.1	3.27	6.07	4.92	0.302
Proportioned		52	24.4	4	28.6	0.214
Moderately disproportional		93	43.7	3	21.4	
Severely disproportional		68	31.9	7	50	

PJF: proximal junctional failure; SD: standard deviation; BMI: body mass index;

PI-LL: pelvic incidence minus lumbar lordosis; SVA: surgical vertical axis; PT: pelvic tilt

C7CSVL; distance between C7 plumb line and central sacral vertical line

Table 2. Risk factors for revision due to PJF (multivariate analysis)

	Odd ratio	95% CI	p-value
BMI	1.16	1.030–1.300	0.013
PT(°)	1.04	0.988–1.100	0.127

BMI, body mass index; PT, pelvic tilt; CI, confidence interval

Table 3. Comparison of RF and non-RF

			Non-RF		RF		
N			210	(%, SD)	49	(%, SD)	p-value
Demographic Variables							
Age			68.41	9.81	68.29	9.09	0.933
Gender	F		177	84.3	40	81.6	0.668
	M		33	15.7	9	18.4	
BMI			23.08	3.74	23.28	3.87	0.73
Surgical Variables							
Total blood loss(ml)			1568.71	1041.12	1905.6	112.24	0.048
Total surgical time(min)			441.64	85.08	432.84	81.54	0.512
Number of fusion segments			9.3	2.24	10.12	2.42	0.023
Three column osteotomy			64	30.5	31	63.3	<0.001
lateral lumber interbody fusion			84	40.0	6	12.2	<0.001
Number of rods	2		135	55.0	31	64.6	0.104
	3		24	11.5	8	16.7	
	4		70	33.5	9	18.8	
Preoperative Radiographic Parameters							
PI-LL(°)			34.30	17.62	33.67	16.83	0.822
SVA(ml)			122.94	17.62	150.98	81.86	0.033
PT(°)			35.18	11.08	38.33	11.08	0.075
C7CSVL(ml)			32.37	32.35	27.85	33.23	0.397
Cobb(°)			30.34	21.36	23.85	22.74	0.063
Postoperative Radiographic Parameters							
PI-LL(°)			7.11	13.45	10.47	11.96	0.111
SVA(ml)			48.9	41.35	48.7	43.58	0.977
PT(°)			20.4	9.78	21.78	8.45	0.368
C7CSVL(ml)			38.79	15.24	44.53	15	0.019
Cobb(°)			9.52	8.49	8.23	10.29	0.367
Global Alignment and Proportional (GAP) Parameters							
GAP score			5.07	3.34	5.54	3.59	0.395
Proportioned			44	24.3	12	26.1	0.144
Moderately disproportional			82	45.3	14	30.4	
Severely disproportional			55	30.4	20	43.5	

PJF: proximal junctional failure; SD: standard deviation; BMI: body mass index;

PI-LL: pelvic incidence minus lumbar lordosis; SVA: surgical vertical axis; PT: pelvic tilt

C7CSVL; distance between C7 plumb line and central sacral vertical line

Table 4. Risk factors for revision due to RF (multivariate analysis)

	Odds ratio	95% CI	p-value
Total blood loss	1.00	1.000–1.000	0.789
No. of fusion segments	1.21	1.050–1.390	0.009
Three-column osteotomy	4.41	2.230–8.720	<0.001
LLIF	0.39	0.123–1.100	0.094
Preoperative SVA	1.00	0.006–0.174	0.858

RF, rod fracture; LLIF, lateral lumbar interbody fusion; SVA; sagittal vertical axis; CI, confidence interval