Studies on *Penaeus orientalis* KISHINOUE—VIII Structure of the Newly Found Lymphoid Organ

Masao OKA*

(Received February 7, 1968)

A number of unknown organs have been discovered in prawns, genus *Penaeus*, suborder *Natantia*. The organ described here is one of those. It was first discovered in Korai prawn, *P. orientalis* KISHINOUE. Consequently, its presence was also confirmed in Ginger prawn, *P. japonica* BATE, *P. carinatus* DANA, White prawn, *Penaeus indicus* de MAN, and *Penaeus merguensis* de MAN. From the histological structure and the manner in which the out-going and in-coming vessels are connected, it can be easily imagined that this organ belongs to the vascular system. However, no such organ in the vascular system has been known to be other than one of the leucopoietic organs¹⁾⁻⁵⁾ on which studies have long been made.^{1,3,5)} The location of the leucopoietic organ being close to the dorsal side of the cardiac stomach can be confirmed in genus *Penaeus* as well as in the others. By comparison of the location, the leucopoietic organ and this newly found organ can be said to be quite different. Further, certain cells which can be seen in this organ resemble lymphocytes more than the lymphocyte type cells^{4,6,7,8)} originating from the leucopoietic organ. In addition, the interstitial tissues surrounding it become lymphoid tissues. For these reason, this organ can be regarded as a lymphoid organ and can be distinguished from the hitherto known leucopoietic organ.

Materials and Methods

Materials used were Korai prawn from the Yellow Sea, Ginger prawn from the Nagasaki coast, and in addition to those, three other species of Genus *Penaeus* from the seas off the Philippines, Singapore and Colombo collected during 1966-1967. Anatomical and histological methods were used for the investigation on the structural of the research materials.

Results and Discussion

The position of this organ differs slightly between the male and the female. In the female of Korai prawn, it lies between the liver and the first branch of the ovary being pressed closely onto the upper anterior part of the former. In the male it lies between the liver and stomach, loosely connected to both. This organ is paired on the left and right side of the anterior portion of the liver, with a pair of afferent and many efferent vessels

* Faculty of Fisheries, Nagasaki University, Nagasaki, Japan (岡 正雄: 長崎大学水産学部)

distributed all over it. The pair of thick-walled vessels leading into the organ through its lower posterior surface and connected with the artery at the other end may be referred to as the afferent vessels. In contrast, the thin-walled vessels and cysts emerging from the entire of the surface organ may be referred to as the efferent vessels. The afferent vessels consist of an endothelial cell layer and loose fibrous tissue surrounding it and in this fibrous tissue mesenchymal reticular cells can be seen. The efferent vessels consist of the adventitia of longitudinally running fibers and the intima of endothelial cells. The afferent vessels enter the organ through the hilus on its lower posterior surface, and an efferent vessel also emerges from this hilus. Tracing the afferent vessels backward from the hilus, it can be seen that they run along the anterior edge of the liver and are connected ultimately with the antennal artery. In the female the afferent vessel joins the antennal artery immediately before passing through the ovary. From this the afferent vessel can be said to be an artery but it differs from the artery in structure, as stated previously. Efferent vessels usually expand to form cysts, but those efferent vessels emerging from the anterior external edge and those from the interior edge are tubiform. The former are connected to the upper part of the leucopoietic organ above the cardiac stomach, to the dorsal epidermis, gonads, etc. The latter are linked with the lymphoid vessels collectively located below the cardiac stomach. The sinuses emerging from several places of the organ are complicatedly bent around it and are linked with the valved lymphoid type vessels. The cystiform vessels link directly with liver go through small openings in the liver tunica. Immediately after the afferent vessel runs into the organ, the fibrous tissue making up the external layer of the vessel becomes loose and joins the fibrous component of the tissues surrounding the artery of this organ. The afferent vessel in the organ is eventually bifurcated to becoming finer and distributed evenly. Then each vessel becomes the central artery of the sheath which is a structural unit of the organ. However it is unknown whether the distal ends open into the sinuses of the organ or not. The surface of this lymphoid organ is covered with collagenous fibers which are the components of the cystiform efferent vessels. The fibrous layer of the tunica is especially dense at the hilus. The base of the efferent vessel penetrates deeply into the organ; the fibers of its wall, as in the afferent vessels are linked up with the organ's fibrous tissue. The lumen of these vessels join the sinuses of the organ.

It is believed that the pulp of this organ consists mainly of lymphoid tissues which surround the central artery forming the peri-arterial sheath⁹⁾. The wall of the central artery is a layer of flat endothelial cells; within this layer are interstices where interchange of substances between blood and the surrounding pulp tissues occurs. Furthermore the endothelial cells are easily transformed and frequently enlarged to protrude into the lumen of the central artery. The peri-arterial sheath is composed of free cells and reticular interstitial tissue. The cells of the reticular tissue are angular ectodermal cells, stained orange-red by Mallory's stain and with phloxin by Gomori's chrome-alum haematoxylin phloxin

Fig. 1. Position of the lymph-like organ in the female of Korai prawn.

Fig. 2. Central artery and peri-arterial sheath which compose a structural unit of the organ.

stain. Lattice fibers of argyrophils are normally in close adherence with the reticular tissue, but in this organ, they are distributed in fragments and concentrically within the peri-arterial sheath slightly distant from the endothelial layer. Fibers which stain well with aniline blue or fuchsin acid are relatively densely distributed outside this argyrophil fiber layer and reticular cells sparsely distributed among these fibers. The free cells which are supposed to be mostly lymph-like cells, are distributed mainly from the argyrophil fiber layer to the outside of the endothelial layer. In the fibrous layer where this kind of lymph-like cells can be seen, there are observed medium sized, poor staining ones which are believed to be their young cells. Moreover, these free cells are also seen inside the artery. Some are extruding into the arterial lumen through interstices in the endothelial cell layer. Hence, it is believed that the lymph-like cells are produced by the reticular tissue and released into the central artery to be transported to where they are needed.

Inside the organ, the spaces between the peri-arterial sheaths take the form of winding, interconnected sinuses. The sinus wall is made up of noncompact endothelial cell layer and argyrophil lattice fibers. Many thinly staining, discoid free cells measuring $6.0\text{--}8.0\ \mu$ in diameter can also be recognized in the sinuses. Lymph-like cells which are the free cells of the peri-arterial sheath, have the diameter of $2.5\text{--}4.0\ \mu$. The cytoplasm is very thin and the nuclei, especially the edges, are stained well by haematoxylin. In the lymphoid tissue, there are observed free cells measuring $6.0\text{--}8.0\ \mu$ in diameter surrounded by clear cytoplasm and containing the nuclei similar to those of lymph-like cells. Upon comparing these free cells with the lymphocyte type cells⁷⁾ produced by the leucopoietic organ⁷⁾ and believed to resemble the former most closely, the latter were found to have the cell diameter of $6.0\text{--}10.0\ \mu$ and nuclear diameter of $4.0\text{--}6.0\ \mu$. From this size difference alone, these two cell types can be said to be different. Moreover, the difference is made more clear by the different staining with chrom-alum haematoxylin stain. For this reason, it appears certain that the free cells of the peri-arterial sheath are not identical to the introduced leucocytes. Though the free cells in the sinuses resemble the lymphocyte type cells of the blood most closely in many aspects, they cannot be discriminated by size alone. The free cells in the sinuses stain extremely poorly with chrome-alum haematoxylin, but this fact alone cannot be the basis for distinguishing the two types of cells. The lymph-like cells of the peri-arterial sheath frequently vary in quantity. It is believed that this phenomenon occurs with abnormal changes in the body. The peri-arterial sheath is filled with lymph-like cells and seemingly their young cells, especially after spawning. The same phenomenon can also be witnessed in the central artery. In abnormal condition, it is occasionally observed that the reticular cells of the peri-arterial sheath degeneratively become fibers and that a large amount of granules appear inside the fibers and they are released into the central artery.

Summary

As a lymphoid organ was found in the vascular system, its anatomical and histological structure was studied.

1. The position of this organ differs slightly in the male and female. In the female, it is pressed by the ovary onto the upper anterior part of the liver.

2. An afferent and many efferent vessels led in and out are observed. The former is connected with the antennal artery and runs along the anterior edge of the liver to be led ultimately into the organ. In the female, the afferent vessel passes through the ovary. It is composed of an endothelial cell layer and surrounding loose fibrous tissue. Mesodermal reticular cells are scattered in this fibrous tissue.

The efferent vessels emerging from several places of this organ usually expand to form cysts but those emerging from the anterior external edge and from the interior edge are tubiform. The former are connected to the upper part of the leucopoietic organ, dorsal epidermis, gonads, etc. The latter are linked with the lymphoid vessels below the cardiac stomach. The cystiform vessels complicatedly bending around the organ are linked with the lymphoid vessels, but the one linked with the liver tunica penetrates the small openings of the tunica into the liver. The efferent vessels are composed of adventitia of longitudinally running fibers and the intima of endothelial cells.

3. The afferent vessel is bifurcated progressively immediately after entering the organ, and each becomes the central artery of the sheath which is a structural unit of this organ. The spaces between the peri-arterial sheaths take the form of winding, interconnected sinuses. The wall of sinus is made up of sporadic endothelial cells and argyrophil lattice fibers.

4. The pulp of this organ is the peri-arterial sheath which surrounds the bifurcated artery in the organ. The wall of the central artery is made up of a layer of flat endothelial cells in which interstices are found. The peri-arterial sheath is composed of free cells and interstitial tissues which consist of reticular tissue and fibrous tissues. In fibrous tissues, argyrophil fibers are distributed in fragments and concentrically within the peri-arterial sheath, and collagenous fibers are distributed outside this layer of argyrophil fibers.

5. The greater part of the free cells in the peri-arterial sheath are lymph-like cells and distributed mainly in the argyrophil fiber layer. In this fibrous layer, there are observed certain cells which are believed to be their young cells. Moreover, these free cells are extruding into the arterial lumen through interstices in the endothelial cell layer. Lymph-like cells measure 2.5–4.0 μ in diameter and they have very thin cytoplasm, and nuclei which are stained well by haematoxylin especially at the peripheries. Discoid free cells, measuring 6.0–8.0 μ in diameter and poorly stained by the haematoxylin can also be recognized in the sinuses. Comparing these free cells with the lymphocyte-type cells produced in leuco-

poietic organ, which resemble each other most closely, the lymph-like cells are clearly distinguishable from the lymphocyte-types, but free cells in the sinuses can not be distinguished from them.

6. The lymph-like cells of the peri-arterial sheath frequently vary in quantity with the abnormal changes in the body. The peri-arterial sheath, as well as the central artery, is filled with lymph-like cells and their young cells. Besides this, the reticular cells of the peri-arterial sheath degeneratively fibers and a large amount of granules appear in the fibers. It can be further confirmed that those cells are released into the central artery.

References

- 1) H. BAUMAN: *Z. wiss. Zool.*, **118**, 247–312 (1921).
- 2) E. FISHER-PIETTE: *Arch. Zool. Exptl. et Gén.*, **74**, 33–52 (1931).
- 3) J. H. LOCHHEAD and M. S. LOCHHEAD: *J. Morphol.*, **68**, 593–632 (1941).
- 4) W. DALL: *Aust. J. mar. Freshwat. Res.*, **15**, 145–161 (1964).
- 5) D. M. MAYNARD: *The physiology of Crustacea—I*, 161–226, Academic Press, N.Y. & London (1960).
- 6) R. NOLD: *Zool. Anz.*, **52**, 277–285 (1921).
- 7) W. C. GEORGE: *J. Morphol.*, **83**, 425–443 (1948).
- 8) M. E. TONY: *Growth*, **22**, 35–50 (1958).
- 9) T. TOKARI: *Histology*, 44–47, 109–121, Nanzando, Tokyo (1964), in Japanese.