Short Paper

Circadian Locomotor Activity in Ayu
*Plecoglossus altivelis*Maung Minh-Nyo,*¹ Mitsuo Tabata,*²
and Mikio Oguri*¹

(Received September 21, 1990)

Although we have demonstrated the circadian rhythmicity of locomotor activity in freshwater fish,¹⁻³⁾ it has not yet been studied in the diurnal freshwater fish. To demonstrate the property of circadian rhythmicity and diel locomotor activity in well known diurnal fish, ayu *Plecoglossus altivelis* was used in the present study.

Ayu of 25–40 g in body weight and 12–15 cm in total length were used. The conditions for an experimental tank, light regime, activity detecting and recording systems were the same as those described elsewhere.^{2,3)} Circadian rhythm was detected by the periodogram analysis.

Fig. 1. Actogram (A) of ayu under LD 12:12 (LD), constant darkness (DD) and constant light (LL). Shadowy and white areas indicate the dark and light phase, respectively. The records were double plotted for clarity. Periodograms (B & C) were obtained from indicated period. The Q(P) indicates a predominant rhythmic component. Slanted line indicates 95% confidence limit.

Locomotor activity was undoubtedly higher under the light phase and lower in the dark phase as shown in the actogram (Fig. 1, A). When the animal was held under continuous darkness (DD, complete darkness), the locomotor activity desynchronized to the previous light and dark (LD) cycles (L=100 lx). Rhythmicity could not be detected as shown in both actogram and periodogram (Fig. 1, A and B). Total amount of locomotor activity in DD increased by an unknown cause. When the animal was reexposed to the LD cycles, diel locomotor activity was apparently observed during several cycles but gradually decreased. However, a distinct 24 hr diurnal activity was confirmed by the periodogram analysis. Under LL, the animal showed free-running locomotor activity. The circadian rhythmicity was observed with period length of 24.0 hr (Fig. 1, C). Despite the presence of circadian locomotor activity in continuous light (LL), it did not last for prolonged cycles. The circadian locomotor activity usually faded-out after 10–14 days which was usually observed also in the nocturnal freshwater fish.²⁻³⁾ Therefore, such persistence of free-running rhythmicity may be a common character in both nocturnal and diurnal freshwater fish.

We examined the shifting experiment of LD regime by 6 hr delaying (light onset 12:00, offset 0:00) and advancing (light onset 6:00, offset 18:00). This experiment is useful to know the contribution of circadian oscillator on the diel locomotor activity. In either direction, ayu synchronized to the shifted LD regime within a day. This was also observed in catfish²⁾ but was slightly different from kawamutsu. These findings indicate that the circadian oscillator of ayu was strongly influenced by LD cycles, which was similar to the result of catfish but different from kawamutsu. It is assumed, therefore, that the diel locomotor activity is one of the behaviors responding directly to the environmental light and dark cycles.

References

- 1) M. Tabata, M. Minh-Nyo, and M. Oguri: *Exp. Biol.*, **47**, 219–225 (1988).
- 2) M. Tabata, M. Minh-Nyo, H. Niwa, and M. Oguri: *Zool. Sci.*, **6**, 367–375 (1989).
- 3) M. Tabata, M. Minh-Nyo, and M. Oguri: *Nippon Suisan Gakkaishi*, **57**, 607–612 (1991).
- 4) M. Tabata: *Dôbutsu Seiri*, **3**, 103–112 (1986).
- 5) M. Minh-Nyo: Studies on locomotor activity and the role of retinal and extraretinal photoreceptor organs in circadian organization of teleosts, doctoral thesis, Nagoya University, Nagoya, Japan, 1989, 1–168.

*¹ Laboratory of Fish Biology, School of Agriculture, Nagoya University, Chikusa, Nagoya 464-01, Japan (M. Minh-Nyo, 小栗幹郎: 名古屋大学農学部).

*² Department of Bioscience, The Nishi-Tokyo University, Uenohara, Yamanashi 409-01, Japan (田畑満生: 西東京科学大学理工学部).