Short Communication

**CYCLIC AMP-DEPENDENT PROTEIN
KINASE-INDEPENDENT ANGIOTENSIN II-INDUCED
INHIBITION OF CALCIUM CURRENT IN HAMSTER
SUBMANDIBULAR GANGLION NEURONS**

TAKAYUKI ENDOH, MITSUHIRO ABE and TAKASHI SUZUKI

*Department of Physiology, Tokyo Dental College,
1-2-2 Masago, Mihama-ku, Chiba 261-8502, Japan*

Received 6 March, 2002/Accepted for Publication 25 March, 2002

Abstract

In a previous study, we demonstrated that angiotensin II (Ang II) inhibited voltage-dependent calcium channels (VDCCs) currents (I_{Ca}) in hamster submandibular ganglion (SMG) neurons. In sinoatrial node cells, it has been reported that Ang II inhibits I_{Ca} by suppressing cyclic AMP production. In this study, to investigate the possible involvement of a cyclic AMP-cyclic AMP dependent protein kinase (PKA) pathway in the Ang II-induced inhibition of I_{Ca} , effects of Ang II were examined in SMG neurons after treatment with an activator and inhibitor of PKA. Neither pretreatment of neurons with membrane permeable cyclic AMP nor intracellular dialysis of PKA blocker attenuated the Ang II-induced inhibition of I_{Ca} . These results indicate that Ang II inhibited I_{Ca} via a cyclic AMP-PKA-independent mechanism in SMG neurons.

Key words: Hamster submandibular ganglion neurons—
Voltage-dependent calcium channels—Angiotensin II—
Cyclic AMP—Protein kinase A

INTRODUCTION

Angiotensin II (Ang II), one of the main active peptides of the renin-angiotensin system, exerts its multiple functions, including drinking, vasopressin and oxytocin release, natriuresis, cardiovascular, endocrine and neuronal targets. This peptide is one of the most important vasoconstrictive hormones but is also known to act as a neuromodulator and a neurotransmitter in the central and peripheral nervous systems. These actions of Ang II involve modulation of membrane ion

channel currents and neuronal activity.

It is well known that cyclic AMP-cyclic AMP dependent protein kinase (PKA) pathways are the most important second messenger pathways which mediate hormone-ion channel regulation. Intracellular Ca^{2+} , which acts as a second messenger, can interact with cyclic AMP, and diacylglycerol (DAG)-protein kinase C (PKC), which is another important second messenger, interacts with the cyclic AMP-PKA pathways.

It has been reported that Ang II receptors cause inhibition of PKA through inhibition of

adenylyl cyclase and reduction in cyclic AMP in various tissues^{1,2,10}). Additionally, it has been demonstrated that Ang II inhibits I_{Ca} by suppressing cyclic AMP production in sinoatrial node cells³).

We previously reported that Ang II, mediated via Ang II type 1 (AT_1) receptors, inhibits voltage-dependent calcium channels (VDCCs) currents (I_{Ca}) in submandibular ganglion (SMG) neurons⁷). It remains unclear, however, whether cyclic AMP-PKA pathways are mediated in the Ang II-induced inhibition of I_{Ca} in this neuron. Thus, the purpose of the present study was to examine the contribution of cyclic AMP-PKA pathways in the Ang II-induced inhibition of I_{Ca} in the SMG neuron.

MATERIALS AND METHODS

Experiments were conducted according to the guidelines for the treatment of experimental animals at Tokyo Dental College. SMG neurons from hamsters were acutely dissociated with a modified version of the method described previously⁷). In brief, SMG neurons were isolated from 4–6 week-old hamsters and maintained in Ca^{2+} -free Krebs solution of the following composition (in mM): 136 NaCl, 5 KCl, 3 $MgCl_2 \cdot 6H_2O$, 10.9 glucose, 11.9 $NaHCO_3$ and 1.1 $NaH_2PO_4 \cdot 2H_2O$. The neurons were treated with collagenase type I (3 mg/ml in Ca^{2+} -free Krebs solution; Sigma, St. Louis, MO, U.S.A.) for 50 min at 37°C, followed by incubation in trypsin type I (1 mg/ml in Ca^{2+} -free Krebs solution; Sigma, St. Louis, MO, U.S.A.) for an additional 10 min. The supernatant was replaced with normal Krebs solution of the following composition (in mM): 136 NaCl, 5 KCl, 2.5 $CaCl_2$, 0.5 $MgCl_2 \cdot 6H_2O$, 10.9 glucose, 11.9 $NaHCO_3$ and 1.1 $NaH_2PO_4 \cdot 2H_2O$. Neurons were then plated onto poly-L-lysine (Sigma)-coated glass coverslips.

Voltage-clamp recordings were conducted using the whole-cell configuration of the patch clamp technique⁴). Fabricated recording pipettes (2–3 M Ω) were filled with an

internal solution with the following composition (in mM): 100 CsCl, 1 $MgCl_2$, 10 HEPES, 10 BAPTA, 3.6 $MgATP$, 14 tris₂phosphocreatine, 0.1 GTP, and 50 U/ml creatine phosphokinase. The pH was adjusted to 7.2 with CsOH. After the formation of a giga seal, in order to record I_{Ca} , the external Krebs solution was replaced by solution containing the following (in mM): 67 cholinechloride, 100 tetraethylammonium chloride, 5.3 KCl, 5 $CaCl_2$ and 10 HEPES. The pH was adjusted to 7.4 with Tris base. Command voltage protocols were generated with a computer software pCLAMP version 8 (Axon Instruments, Union City, CA, U.S.A.) and transformed to an analogue signal using a DigiData 1200 interface (Axon Instruments, Union City, CA, U.S.A.). The command pulses were applied to the cell through an L/M-EPC7 amplifier (HEKA Elektronik, Lambrecht, Germany). The currents were recorded with the amplifier and a computer software pCLAMP 8 acquisition system.

All data analyses were performed using the pCLAMP 8 acquisition system. Statistical analyses were made by Student *t*-test. Probability (*p*) values of less than 0.05 were considered significant.

Ang II was purchased from Peptide Institute. Losartan was a gift from Merck & Co., Inc. 8-(4-chlorophenylthio)-adenosine 3': 5'-cyclic monophosphate (CPT-cAMP) was purchased from Sigma, St. Louis, MO, U.S.A. The PKA inhibitor peptide, PKI (5–24), was purchased from BIOMOL Research Laboratories Inc. (Plymouth, PA, U.S.A.).

RESULTS

An example of the effects of Ang II and AT_1 receptor antagonist, losartan, on I_{Ca} is shown in Fig. 1 (A). Application of 1 μM Ang II inhibited I_{Ca} from $-2,408$ pA to $-1,160$ pA (51.8% inhibition) in this neuron. 1 μM losartan completely antagonized the inhibitory effect of Ang II.

To evaluate the possible contribution of PKA to the Ang II-induced inhibition of I_{Ca} ,

Fig. 1 (A) Effects of Ang II and losartan on I_{Ca} . Superimposed I_{Ca} traces were recorded using a voltage protocol at the times indicated in the time course graph (right). Open points in the graph indicate I_{Ca} . Ang II ($1\mu\text{M}$) and losartan ($1\mu\text{M}$) were bath applied during the times indicated by the filled and open bars, respectively.

(B) Effects of Ang II on I_{Ca} in a neuron pretreated with CPT-cAMP ($10\mu\text{M}$ for 10 min).

(C) Effects of Ang II on I_{Ca} in the presence of PKI (5-24) ($20\mu\text{M}$ for 10 min) contained in the recording pipette.

the effects of Ang II on I_{Ca} in neurons treated with CPT-cAMP ($10\mu\text{M}$ for 10 min), membrane permeable cyclic AMP, were investigated. We have previously demonstrated that this compound has a sufficient effect to cause the saturation of intracellular cyclic AMP levels¹¹. As shown in Fig. 1 (B), Ang II-induced inhibition of I_{Ca} was not affected by CPT-cAMP treatment. In addition, intracellular dialysis of PKI (5–24) ($20\mu\text{M}$ for 10 min), PKA inhibitor, also did not attenuated the Ang II-induced inhibition of I_{Ca} as shown in Fig. 1 (C).

The data showing the effects of altering PKA activity and antagonizing AT_1 receptors are summarized in Fig. 2. On average, the Ang II-induced inhibition of I_{Ca} was $32.1 \pm 2.7\%$ ($n=9$) for control, $7.6 \pm 1.5\%$ ($n=7$) for neurons treated with losartan, $32.7 \pm 4.7\%$ ($n=5$) for the neurons treated with CPT-cAMP, and $25.6 \pm 3.9\%$ ($n=5$) when the recording pipette was filled with PKI (5–24).

DISCUSSION

The present results show that Ang II inhibited I_{Ca} by a cyclic AMP-PKA pathway-independent mechanism. In contrast, there is considerable evidence that Ang II receptors cause inhibition of cyclic AMP-PKA through inhibition of adenylyl cyclase and reduction in cyclic AMP^{1–3,10}. This evidence raises several questions: (1) which processes are absent in Ang II-induced inhibition of I_{Ca} in SMG?; (2) are cyclic AMP-PKA pathways between AT_1 receptors and VDCCs possibly absent in SMG?; and (3) is there a possible lack of AT_1 receptors that is selectively regulating cyclic AMP-PKA pathways in SMG?

Various reports have demonstrated the distinct second messenger pathways in the Ang II-induced ion channel modulations: for instance, PKC pathways and calcium/calmodulin-dependent protein kinase II (CaM KII) pathways in the central nervous system (CNS) neurons^{9,12}, Ras/Raf/mitogen-activated protein (MAP) kinase pathways and Janus kinase/signal transducers and

Fig. 2 Summary of Ang II-induced inhibition of I_{Ca} determined when various conditions. Histogram demonstrating the degree of I_{Ca} inhibition by $1\mu\text{M}$ Ang II in control (untreated neurons), after losartan (AT_1 receptor antagonist), after CPT-cAMP (membrane permeable cyclic AMP) and intracellular dialysis of PKI (5–24) (PKA inhibitor). Ordinates represent the Ang II-induced inhibition of I_{Ca} . Number of neurons tested is indicated in parentheses. *Significant difference relative to control ($p < 0.05$, by post hoc Dunnett's test)

activators of transcription (JAK/STAT) pathways^{6,8}, Fos-regulating kinase and c-Jun NH_2 -terminal kinase⁵.

The present data do not allow us to deduce the exact mechanism by which intracellular pathways may alter VDCCs function. It will be important in future experiments to determine what second messenger pathways contribute to the Ang II-induced inhibition of I_{Ca} in SMG neurons.

ACKNOWLEDGEMENTS

This work was supported by a Grant-in-Aid for Scientific Research from the Japanese Ministry of Education, Culture, Sports, Science and Technology (No.12671816).

REFERENCES

- 1) Anand-Srivastava, M.B. (1983). Angiotensin II receptors negatively coupled to adenylyl cyclase in rat aorta. *Biochem Biophys Res*

- Commun* **117**, 420–427.
- 2) Griendling, K.K., Rittenhouse, S.E., Brock, T.A., Ekstein, L.S., Gimbrone, M.A., Jr. and Alexander, R.H. (1986). Sustained diacylglycerol formation from inositol phospholipids in angiotensin II-stimulated vascular smooth muscle cells. *J Biol Chem* **261**, 5901–5906.
  - 3) Habuchi, Y., Lu, L.-L., Morikawa, J. and Yoshimura, M. (1995). Angiotensin II inhibition of L-type Ca²⁺ current in sinoatrial node cells of rabbits. *Am J Physiol* **268**, H1053–H1060.
  - 4) Hamill, O.P., Marty, A., Neher, E., Sakmann, B. and Sigworth, F.J. (1981). Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflügers Archs* **391**, 85–100.
  - 5) Huang, X.-C., Deng, T. and Sumners, C. (1998). Angiotensin stimulates activation of Fos-regulating kinase and c-Jun NH₂-terminal kinase in neuronal cultures from rat brain. *Endocrinology* **139**, 245–251.
  - 6) Huang, X.-C., Richards, E.M. and Sumners, C. (1996). Mitogen-activated protein kinases in rat brain neuronal cultures are activated by angiotensin II type 1 receptors and inhibited by angiotensin II type 2 receptors. *J Biol Chem* **271**, 15635–15641.
  - 7) Ikegami, H., Endoh, T. and Suzuki, T. (2001). Angiotensin II-induced inhibition of calcium currents in hamster submandibular ganglion neurons. *Neurosci Res* **41**, 227–232.
  - 8) Marrero, M.B., Schieffer, B., Paxton, W.G., Heerdt, L., Berk, B.C., Delafontaine, P. and Bernstein, K.E. (1995). Direct stimulation of Jak/STAT pathway by the angiotensin II AT₁ receptor. *Nature* **375**, 247–250.
  - 9) Sumners, C., Zhu, M., Gelband, C.H. and Posner, P. (1996). Angiotensin II type 1 receptor modulation of neuronal K⁺ and Ca²⁺ currents: Intracellular mechanisms. *Am J Physiol* **271**, C154–C163.
  - 10) Unger, T., Chung, O., Csikos, T., Culman, J., Gallinat, S., Gohlke, P., Hohle, S., Meffert, S., Stoll, M., Stroth, U. and Zhu, Y.Z. (1996). Angiotensin receptors. *J Hypertens Suppl* **14**, S95–103.
  - 11) Yamada, T., Endoh, T. and Suzuki, T. (1999). Inhibition of calcium channels by neurokinin receptor and signal transduction in hamster submandibular ganglion cells. *J Auton Nerv Sys* **76**, 1–8.
  - 12) Zhu, M., Gelband, C.H., Posner, P. and Sumners, C. (1999). Angiotensin II decreases neuronal delayed rectifier potassium current: Role of calcium/calmodulin-dependent protein kinase II. *J Neurophysiol* **82**, 1560–1568.

Reprint requests to:

Dr. Takayuki Endoh
Department of Physiology,
Tokyo Dental College,
1-2-2 Masago, Mihama-ku,
Chiba 261-8502, Japan