The Effect of Anoxia on the Intestinal Secretion.

By

Tatuji Suzuki.

(鈴木 達二)

(From the Physiological Laboratory of Prof. Y. Satake,
Tohoku University, Sendai.)

(Received for publication, July 26, 1949)

Although the effect of anoxia on the secretion of the enteric juice was studied by Northup and Van Liere¹⁾ about ten years ago, their investigations were performed under non-physiological conditions of animals such as anesthesia and laparotomy.

In the present studies, the dog was used, in which the Thirty fistula was prepared with the small intestine 10–20 cm. distal from the duodeno-jejunal flexura. At the time of experiment a gummi catheter was inserted in the fistula, so as to avoid the influence of intestinal movements on the ejection of intestinal juice as possible.²⁾

Placing the animal on the table in the large low pressure chamber, the rate of intestinal secretion for every one hour in non-feeding period was estimated. The barometric pressure in the chamber was lowered to the pressure corresponding to the altitude of 8 km. during one hour.

Comparing the output rate of enteric juice at normal, and lowered barometric pressure of the chamber, we could not find any definite difference among them, although some fluctuations of small magnitude were observed. The result suggests that the intestinal secretion in non-feeding period is not influenced by the anoxia of such severe degree as described above.

Literature.

- 1 Northup and Van Liere, Proc. Soc. Exp. Biol. Med., 1939, **42**, 162.
- 2 Takahasi, Tohoku J. Exp. Med., 1949, **50**, 157.