Biochemical Studies on Carbohydrates.

CXXXIII. Paper Partition Chromatograms of Sugar Components in Glycidamins and Glycoproteins. I.

By

Hajime Masamune and **Zensaku Yosizawa.**

(正宗 一)

(吉澤 善作)

(From the Medico-chemical Institute, Tohoku University, Sendai.)

(Received for publication, October 25, 1949)

We have been long engaged in the study of glycidamins and glycoproteins (Cf. Masamune¹⁾). However, in some of them, sugar components of their polysaccharide or polysaccharides have not been firmly identified due to lack of material or due to difficulty in their separation so that the recent expedient means of paper partition chromatography, which requires only a small quantity of substance, was applied to them anew. The materials at first examined were sero-, ovo-, vitello- and salivary-mucoid, and snail foot- and mucus- and salivary mucin. The results obtained will be tabulated:

Glycidamin or glycoprotein	Sugar components
Seromucoid	Glucosamine, galactose, mannose
Ovomucoid	Glucosamine, chondrosamine, galactose, mannose
Vitellomucoid	Chondrosamine, mannose
Sublingual mucoid	Glucosamine, galactose, glucuronic acid
Sublingual mucin	Glucosamine, galactose, mannose, glucuronic acid, methylpentose
Snail foot mucin	Glucosamine, galactose
Snail mucus mucin	Glucosamine, mannose, galacturonic acid

Seromucoid:—The finding that this mucoid contains glucosamine, mannose and galactose accords with the claim of Rimington²⁾ which we have accepted³⁾ and the result of Werner and Odin.⁴⁾

Ovomucoid:—That it involves glucosamine and mannose has been generally assumed. In the chromatogram a spot corresponding to galactose appeared, though faint, supporting Werner and Odin,⁴⁾ and chondrosamine was further detected.

Vitellomucoid:—Onoe considered the hexosamine in this glycidamin to be glucosamine from F.P. of 199° and mixed melting point test of its benzoate. But we became aware thereafter that this melting point is that of chondrosamine pentabenzoate (cf. Hisamura and Kusuno⁵⁾) and that

the mixed melting point test of chondrosamine-pentabenzoate and impure glucosamine-pentabenzoate⁶⁾ shows no depression.

Sublingual mucoid and mucin and snail mucin:—The results coincide mostly with our previous assertions, but that a spot corresponding to the expected position of methylpentose occurred in the chromatogram of the salivary mucin, though not distinct.

Snail mucus mucin:—The chromatogram suggested the presence of mannose instead of galactose, conflicting with the observation of Suzuki.⁷⁾ To inquire if the chromatograph tells the real fact in this respect, an attempt is in progress to separate from the mucin the sugar in question as a crystalline derivative.

Fig. 1. *n*-Butylacetate-pyridine-water (5:3:1) chromatograms developed with AgNO_3 -ammonia (duration of test, 8.5 hrs.; temp., $19 \pm 1^\circ\text{C}$). a: standard (glucosamine+galactose+mannose+rhamnose), b: standard (chondrosamine+glucuronic acid), c: seromucoid, d: ovomucoid, e: vitellomucoid, f: sublingual mucoid, g: sublingual mucin, h: snail foot mucin, i: snail mucus mucin. 1: chondrosamine, 2: glucosamine, 3: galactose, 4: mannose, 5: rhamnose, 6: glucuronic acid, 7: galacturonic acid.

Fig. 2. *n*-Butylacetate-acetic acid-water (3:2:1) chromatograms developed with AgNO_3 -ammonia (duration of test, 16 hrs.; temp., $19 \pm 1^\circ\text{C}$).

Notations a-j and 1-7 signify the same as in Fig. 1.

EXPERIMENTAL.

Materials.

The preparations of the glycidamins and glycoproteins were those prepared formerly by us: seromucoid (Osaki and Yosizawa⁸), ovomucoid (H. Suzuki⁹), vitellomucoid (Osaki and Yosizawa¹⁰), sublingual mucoid (Tanabe¹¹), sublingual mucin (Tanabe¹²), snail foot mucin (Suzuki⁷), snail mucus mucin (Masamune, Yasuoka, Takahasi and Asagi¹³).

Technique.

The technique was what had been used by one of us (Yosizawa¹⁴) to obtain the chromatograms of monosaccharides in the A-group carbohydrate from pig stomach mucus.

Namely, 100 mgm. each of the materials were hydrolyzed¹⁵ by heating

Fig. 3. Chromatograms run in parallel to those in Fig. 2 and developed with Elson-Morgan reagents.

Notations same as in Fig. 1.

with 5 cc. of 1 n. H_2SO_4 (100°C, 5 hrs.), the hydrolysate removed from H_2SO_4 and after decolorization, distilled in vacuo to 2/5 the volume. The quantities below of the condensates were put on the paper for the resolution: seromucoid, 0.02 cc.; ovomucoid, 0.04 cc.; vitellomucoid, 0.012 cc.; sublingual mucoid, 0.025 cc.; sublingual mucin, 0.03 cc.; snail foot mucin, 0.10 cc.; snail mucus mucin, 0.05 cc. The standard sugar solutions were 1% regarding each sugar and their 0.015 cc. was used.

Results.

Figs. 1-3 illustrate the acquired chromatograms diagrammatically.

Through the Grant Committee for Scientific Researches of the National Research Council was given a grant from the Education Department in aid to us, which is gratefully acknowledged. H. M.

References and Notes.

- 1) Masamune, *Tohoku J. Exp. Med.*, 1949, **50**, 107.
- 2) Rimington, *Bioch. J.*, 1940, **34**, 931.
- 3) Suzuki, *J. Bioch.*, 1941, **34**, 167; 1944, **36**, 395. Osaki and Yosizawa, *Tohoku J. Exp. Med.*, 1949, **51**, 57.
- 4) Werner and Odén, *Uppsala Läkarfören. Förhandl.*, 1949, **54**, 69.
- 5) Hisamura and Kusuno, *J. Bioch.*, 1938, **27**, 375.
- 6) Onoe described to have employed tetrabenzoyl-glucosamine for the mixed melting point test, but his preparation procedure should have given the pentabenzate.
- 7) Suzuki, *J. Bioch.*, 1941, **33**, 377.
- 8) Osaki and Yosizawa, *Tohoku J. Exp. Med.*, 1949, **51**, 57.

- 9) H. Suzuki, J. Bioch., 1944, **36**, 395. Regarding the preparation procedure cf. T. Hasegawa (Tohoku J. Exp. Med., 1943, **46**, 95).
 - 10) Osaki and Yosizawa, Tohoku J. Exp. Med., 1949, **51**, 63.
 - 11) Tanabe, J. Bioch., 1939, **30**, 11.
 - 12) Tanabe, *ibid.*, 1938, **28**, 227.
 - 13) Masamune, Yasuoka, Takahasi and Asagi, Tohoku J. Exp. Med., 1947, **49**, 177.
 - 14) Yosizawa, *ibid.*, 1950, **53**, 123.
 - 15) Snail foot- and mucus-mucin contained comparatively large amounts of insoluble substances.
-