Original article

The Trend of *Vibrio parahaemolyticus* Infections in Southern Thailand from 2006 to 2010

Junthip Thongjun¹, Pimonsri Mittraparp-arthorn¹, Mingkwan Yingkajorn^{1,2}, Jetnapang Kongreung¹, Mitsunaki Nishibuchi³ and Varaporn Vuddhakul^{1*}

Received 7 March, 2013 Accepted 15 June, 2013 Published online 28 November, 2013

Abstract: The bacterium, *Vibrio parahaemolyticus* was isolated from 776 patients at Hat Yai Hospital in Southern Thailand from 2006 to 2010. 51.3–73.6% of the isolates were *tdh*⁺ *trh*⁺ and Group-specific PCR positive pandemic strains. A comparison of the number of *V. parahaemolyticus* isolates in this study and that from the same hospital in 2000–2005 indicates that this region of Thailand is endemic for *V. parahaemolyticus*.

Key words: *V. parahaemolyticus*, pandemic strains, Thailand, GS-PCR

INTRODUCTION

Vibrio parahaemolyticus is a Gram-negative halophilic bacterium detected worldwide in the marine environment. It is a leading cause of gastroenteritis and generally associated with consumption of raw or improperly cooked seafood [1]. The infective dose of pathogenic *V. parahaemolyticus* has been estimated to be 10⁵ to 10⁷ pathogenic organisms [2]. The incubation period ranges from 4 to 96 h and the illness lasts for 2–3 days. Clinical symptoms include abdominal pain, watery diarrhea, nausea, vomiting and the disease is occasionally associated with headache, fever and chills [1]. However, not all strains of *V. parahaemolyticus* are pathogenic. Most clinical strains possess a major virulence factor, the thermostable direct hemolysin (TDH), and exhibit β-hemolysis on Wagatsuma agar [3]. Another virulence factor, the TDH-related hemolysin (TRH) has also been associated with some food-poisoning outbreaks [4]. TDH and TRH are encoded by the *tdh* and *trh* genes respectively. The percentage of *tdh* and *trh* genes detected in the environment varies and depends on both the location and the detection techniques. However, around 90% of clinical isolates possess either the *tdh* gene, the *trh* gene, or both [5–7].

In Thailand, *V. parahaemolyticus* was first reported in 1970 [8]. Since then, infections have been reported from various communities [9, 10]. The incidence of infections in

Thailand was unremarkable until 1996 when a pandemic strain with the *tdh* virulence gene and O3:K6 serotype emerged [11, 12]. Group-specific PCR (GS-PCR) has been established to detect nucleotide variations unique to the pandemic clone within the 1,364 bp *toxRS* region [13]. The first report of the O3:K6 pandemic strain was obtained from patients at a Hat Yai Hospital in southern Thailand. One isolate of the O3:K6 pandemic strain was detected in shellfish purchased from the same area, indicating that people in this area were at risk of *V. parahaemolyticus* infection [12]. Thus, a surveillance program has been established at Hat Yai Hospital, the main public hospital in Songkhla province, as a public health warning. A total of 865 patients infected with *V. parahaemolyticus* were reported from 2000 to 2005 [7]. Approximately 64.1–69.7% of the isolates between 2000 and 2003 were pandemic strains, defined as GS-PCR positive *tdh*⁺ *trh*⁺ (Fig. 1). However, in 2004 and 2005, the percentage of infections by these strains decreased to 56.1 and 55.5%, respectively. It has been postulated that people previously exposed to pandemic strains may have acquired immunity to the pandemic strains although they continue to be susceptible to non-pandemic strains. Hence, we report our continued surveillance of infection by *V. parahaemolyticus* in Hat Yai Hospital from 2006 to 2010.

¹ Department of Microbiology, Faculty of Science, Prince of Songkla University, Hat Yai 90110, Thailand

² Department of Biomedical Sciences, Faculty of Medicine, Prince of Songkla University, Hat Yai 90110, Thailand

³ Center for Southeast Asian Studies, Kyoto University, Kyoto 606-8501, Japan

*Corresponding author:

Department of Microbiology, Faculty of Science, Prince of Songkla University, Hat Yai 90110, Thailand

Tel: +66-74-288328

Fax: +66-74-446661

E-mail: varaporn.v@psu.ac.th

Fig. 1. Number of total and pandemic strains of *V. parahaemolyticus* isolated from patients in Hat Yai Hospital from 2000 to 2005.

MATERIALS AND METHODS

Sample collection and bacteriology

Rectal swabs were obtained from patients with diarrhea in Hat Yai Hospital, Songkhla Province, Thailand, between 2006 and 2010. Each sample was plated on MacConkey, Salmonella-Shigella and Thiosulfate Citrate Bile-Salts Sucrose (TCBS) agars (Difco, Md., U.S.A.). After incubation at 37°C overnight, sucrose non-fermenter colonies on TCBS were selected and were identified as *V. parahaemolyticus* by standard biochemical tests. Confirmation of *V. parahaemolyticus* was performed by PCR targeted to the *toxR* gene.

ToxR investigation

The test isolate was grown in Luria-Bertani (LB) broth containing 1% NaCl with shaking at 160 rpm at 37°C overnight. One milliliter of the broth culture was centrifuged, and the bacterial cells were washed with sterile saline (0.85% NaCl) and then suspended in it. The cell suspension was boiled for 10 min and the supernatant was obtained by centrifugation, diluted 10-fold in distilled water, and used as the DNA template for PCR amplification. To investigate the *toxR* gene, PCR using primers T4 and T7 was performed, as described previously [14].

Detection of *tdh* and *trh* genes

To determine the presence of the *tdh* and *trh* genes, the DNA template was prepared as described above. PCR was carried out using the primers D3–D5 and R2–R6, respectively [15].

Serotyping

The serotype of *V. parahaemolyticus* was established

by the slide-agglutination technique using anti-O and anti-K antibodies (Denka Seiken, Tokyo, Japan). The test isolate was grown in tryptic soy broth containing 3% NaCl at 37°C for 18 h, and the bacterial cells were suspended in saline (3% NaCl). For the determination of the K serotype, the bacterial cell suspension was subjected to agglutination with specific anti-K antibodies. For the O serotype, the bacterial cell suspension was autoclaved at 121°C for 1 h after which the autoclaved bacterial cells were agglutinated with specific anti-O antibodies.

GS-PCR

GS-PCR was carried out using the primers GS-VP1 and GS-VP2 and the reaction was carried out as previously described [13].

Statistical analysis

Student's *t* test was used for comparison of the results.

RESULTS

From 2006 to 2010, a total of 776 *V. parahaemolyticus* isolates were obtained from patients in Hat Yai Hospital (Table 1). The highest number was in 2006 and the lowest number was in 2009. The bacteria were classified into four groups: *tdh*⁺*trh*⁺, *tdh*⁺*trh*[−], *tdh*[−]*trh*[−] and *tdh*[−]*trh*⁺ based on presence or absence of the *tdh* or *trh* toxin genes. Most of the clinical *V. parahaemolyticus* isolates were in the *tdh*⁺*trh*[−] group, which was divided into pandemic (GS-PCR positive) and non-pandemic (GS-PCR negative) strains. The predominant isolates of *V. parahaemolyticus* in each year were pandemic strains of which the highest number, 73.6%, and the lowest number, 51.3%, were from 2008 and 2010, respectively (Table 1). The numbers of isolates in the other three

Table 1. Characteristics of *V. parahaemolyticus* isolates from Hat Yai Hospital from 2006 to 2010.

Year	total	No. of isolates (%)				
		<i>tdh</i> ⁺ <i>trh</i> ⁻		<i>tdh</i> ⁺ <i>trh</i> ⁺	<i>tdh</i> ⁻ <i>trh</i> ⁻	<i>tdh</i> ⁻ <i>trh</i> ⁺
		GS-PCR positive	GS-PCR negative			
2006	214	138 (64.5)	52 (24.3)	7 (3.3)	16 (7.5)	1 (0.4)
2007	139	83 (59.7)	26 (18.7)	11 (7.9)	17 (12.2)	2 (1.5)
2008	193	142 (73.6)	32 (16.6)	6 (3.1)	11 (5.7)	2 (1.0)
2009	111	67 (60.4)	28 (25.2)	4 (3.6)	9* (8.1)	3 (2.7)
2010	119	61 (51.3)	37 (31.1)	6 (5.0)	15 (12.6)	0
total	776	491	175	34	68	8

* Two isolates were GS-PCR positive

groups, *tdh*⁺*trh*⁺, *tdh*⁻*trh*⁻ and *tdh*⁻*trh*⁺, were highest in 2007 (7.9%), 2010 (12.6%) and 2009 (2.7%) respectively. Two isolates of *V. parahaemolyticus* in the *tdh*⁻*trh*⁻ group obtained in 2009 were GS-PCR positive.

Every year the most prevalent serotype was O3:K6, which accounted for 46.8% of the total isolates (Table 2). Also, O3:K6 was the predominant serotype among the pandemic strains, followed by O4:K8 and O1:K25 (Table 3). The serotypes of isolates from non-pandemic strains were variable; the predominant serotypes were O1:K56, O4:K8, O1:KUT and O4:K9 (Table 2).

DISCUSSION

Infections caused by *V. parahaemolyticus* were investigated between 2006 and 2010. It was found that most of the isolates were pandemic strains and belonged to the O3:K6 serotype. This result is consistent with the findings since 2000 in Hat Yai Hospital, where O3:K6 pandemic strains were the main isolates [7]. This pandemic clone is a permanent part of the bacterial flora in this region. The reasons remains unknown, because it has been demonstrated that the effects of environmental factors such as temperature (either 4 or 50°C), acid stress (pH 4) and low-salinity stress on O3:K6 pandemic strains and other non-pandemic strains are not different [16]. In this study, the percentage of O3:K6 pandemic strains between 2006 and 2010 was alternatively high and low, but the total percentage median of pandemic strains detected in this study (63.3%) was not significantly different from that (62.1%) detected in 2000–2005 (Fig. 2). This indicates that the incidence of *V. parahaemolyticus* infection by the pandemic strains in this southern part of Thailand did not increase.

The clonal diversity of the pandemic strains from this study was similar to that observed in our previous investigation in 2000–2005 [7]. In 2010, however, serotype O4:K68 and O1:KUT were not detected, and four other serotypes

(14 isolates) including O1:KUT (11), O4:K9 (2) and O3:KUT (1) were identified as pandemic strains (Table 3).

The number of isolates belonging to *tdh*⁻*trh*⁻ group was higher than the number of isolates belonging to *tdh*⁺*trh*⁺ and *tdh*⁻*trh*⁺ groups and did not differ from the number previously detected in 2000–2005 (Fig. 2). Most clinical isolates of *V. parahaemolyticus* harbor either *tdh* or *trh* or both. These genes have been demonstrated to be the virulence genes involved in diarrhea pathogenicity [17, 18]. The mechanism of the disease caused by *tdh*⁻*trh*⁻ *V. parahaemolyticus* is not clearly understood, but it has been demonstrated that a clinical *V. parahaemolyticus* strain lacking *tdh* gene resulted from the IS-mediated *tdh* gene deletion of a *tdh*⁺ strain [19, 20]. In this study, we also detected two isolates of *tdh*⁻*trh*⁻ that belong to the O3:K6 serotype and GS-PCR positive pandemic group (Table 1). Therefore, it is possible that the *tdh* gene was deleted in those strains.

Mollusks are important sources of *V. parahaemolyticus* because they are filter feeders. In Thailand, consumption of mollusks, especially cockles contaminated with *V. parahaemolyticus*, is probably a major cause of acute gastroenteritis because the popular way to prepare this seafood is by semi-cooking. The average amount of *V. parahaemolyticus* reported from cockles was 5.1×10^3 cfu/g [21]. Infection by this bacterium is regularly encountered. An effective way to reduce *V. parahaemolyticus* contamination in cockles is needed. It has been reported that the number of *V. parahaemolyticus* in oysters decreased after they were stored at low temperatures of 0 to 5°C [22]. Therefore, simple refrigeration of cockles in the market may help to minimize the risk of infection.

In conclusion, investigation of *V. parahaemolyticus* isolates from Hat Yai Hospital from 2006 to 2010 revealed that the predominant isolates were pandemic O3:K6 strains. The number of isolates belonging to each toxin gene profile was not significantly different from the number from the same hospital in 2000–2005. This indicates that people in

Table 2. Serotypes of *V. parahaemolyticus* isolated between 2006 and 2010.

Year (total isolate)	No. of isolates*	Serotype
2006 (214)	107	O3:K6
	31	O4:K68
	19	O1:K56
	13	O4:K8
	5	O1:KUT [†] , O4:K55
	4	O11:KUT, O12:KUT
	3	O2:K3, O3:K29, O5:KUT,
	2	O1:K9, O4:K9, OUT:KUT
	1	O1:K25, O3:K5, O3:K29, O3:K53, O4:K34, O4:K49, O4:K11, O5:K19, O5:K61, O8:KUT, O11:K40
2007 (139)	73	O3:K6
	9	O1:KUT
	7	O1:K25
	5	O4:K8, O4:K68
	4	O1:K56
	3	O13:K5, O4:K11, O5:KUT, O8:KUT, O11:KUT
	2	O1:K69, O3:K5, OUT:KUT
	1	O1:K9, O1:K32, O1:K64, O2:K3, O2:K28, O3:K75, O3:KUT, O4:K13, O4:KUT, O9:KUT, O10:K19, O10:KUT, O12:KUT, O13:K75, OUT:K6
2008 (193)	106	O3:K6
	32	O1:K25
	9	O1:K56
	7	O1:KUT
	6	O4:K8
	3	O12:KUT
	2	O2:K3, O4:K13, O4:K53, O4:K55, O5:KUT, O8:K74, O9:K44
	1	O1:K9, O1:K58, O1:K69, O3:K5, O3:K29, O3:K45, O3:K75, O4:K12, O4:K68, O5:K17, O5:K61, O8:K38, O8:K41, O8:K44, O9:K44, O11:K22
2009 (111)	49	O3:K6
	18	O4:K68
	9	O4:K9
	6	O1:K56
	5	O1:KUT
	4	O2:K3
	3	O4:K8
	2	O1:K25, O3:KUT, O8:K22, O11:K36
	1	O1:K9, O3:K29, O4:K13, O4:K42, O4:K55, O5:K15, O8:K32, O9:K44, O10:KUT
2010 (119)	47	O3:K6
	24	O1:KUT
	15	O4:K9
	7	O1:K56, O4:K8
	4	O3:KUT
	3	O10:KUT, O11:KUT
	2	O4:K4, O4:KUT, O5:KUT
	1	O4:K3, O8:KUT, O8:K70

* No. of isolates per serotype.

[†] UT, Untypeable

Fig. 2. Comparison of the number of *V. parahaemolyticus* isolates in each of five groups of toxin gene profile from patients in Hat Yai Hospital between 2000–2005 and 2006–2010.

No significant difference was detected in any of the groups of *V. parahaemolyticus* ($P > 0.05$)

Table 3. Predominant serotypes of GS-PCR positive strains.

Year	Serotypes			
	O3:K6	O4:K68	O1:K25	Others
2006	105	30	1	2
2007	70	5	7	1
2008	106	1	31	4
2009	47	16	2	2
2010	47	0	0	14

this area are constantly exposed to pathogenic *V. parahaemolyticus* at the same level. Therefore, this region of Thailand is probably endemic for *V. parahaemolyticus*. In the future it would be worthwhile to investigate the factors involved in this infection. It is also important to find methods to reduce the amount of *V. parahaemolyticus* in shellfish in the market.

ACKNOWLEDGEMENTS

This work was supported in part by funds from the Commission on Higher Education, Thailand, the Royal Golden Jubilee Ph.D. program (Grant No. PHD/0066/2550), National Science and Technology Development Agency, Thailand, and the Japan Society for the Promotion of Science, JSPS (KAKENHI 19101010).

REFERENCES

1. Yeung PMS, Boor KJ. Epidemiology, pathogenesis, and prevention of foodborne *Vibrio parahaemolyticus* infections. Foodborne Pathog Dis 2004; 1: 74–88.
2. Sanyal SC, Sen PC. Human volunteer study on the pathogenicity of *Vibrio parahaemolyticus*. In Fujino T, Sakaguchi G, Sakazaki R, Takeda Y, eds. International Symposium on *Vibrio parahaemolyticus*. Tokyo: Saikon Publishing Company; 1974.
3. Nishibuchi M, Kaper JB. Thermostable direct hemolysin gene of *Vibrio parahaemolyticus*: a virulence gene acquired by a marine bacterium. Infect Immun 1995; 63: 2093–2099.
4. Honda T, Ni Y, Miwatani T. Purification and characterization of a hemolysin produced by a clinical isolate of Kanagawa phenomenon-negative *V. parahaemolyticus* and related to the thermostable direct hemolysin. Infect Immun 1988; 56: 961–965.
5. Deepanjali A, Sanath Kumar H, Karunasagar I, Karunasagar I. Seasonal variation in abundance of total and pathogenic *Vibrio parahaemolyticus* bacteria in oysters along the Southwest Coast of India. Appl Environ Microbiol 2005; 71: 3575–3580.
6. Gutierrez West CK, Klein SL, Lovell CR. High frequency of virulence factor genes *tdh*, *trh*, and *tlh* in *Vibrio parahaemolyticus* strains isolated from a Pristine Estuary. Appl Environ Microbiol 2013; 79: 2247–2252. DOI: 10.1128/AEM.03792-12.
7. Wootipoom N, Bhoopong P, Pomwised R, Nishibuchi M, Ishibashi M, Vuddhakul V. A decrease in proportion of infections by pandemic *Vibrio parahaemolyticus* in Hat Yai Hospital, southern Thailand. J Med Microbiol 2007; 56: 1630–1638.
8. Suthienkul O. Infectious Diseases. 2 ed. Bangkok: Holistic publishing; 1998.
9. Udompat P, Sukuonsakul M, Pokawattana L, Guharat S, Hanshaoworakul W. Food poisoning outbreak in a hotel, Bangkok. J Health Sci 2001; 13: 682–688.

10. Jatapai A, Mounghthong B, Thunyaharn S, Huttayanant S, Rangsin R. An acute gastroenteritis outbreak of *Vibrio parahaemolyticus* O4:K55 in Nursing College, Thailand Trop Biomed 2010; 27: 265–274.
11. Okuda J, Ishibashi M, Hayakawa E, Nishino T, Takeda Y, Mukhopadhyay AK, Garg S, Bhattacharya SK, Nair GB, Nishibuchi M. Emergence of a unique O3:K6 clone of *Vibrio parahaemolyticus* in Calcutta, India, and isolation of strains from the same clonal group from Southeast Asian travelers arriving in Japan. J Clin Microbiol 1997; 35: 3150–3155.
12. Vuddhakul V, Chowdhury A, Laohaprerththisan V, Pungrasamee P, Patararungrong N, Thianmontri P, Ishibashi M, Matsumoto C, Nishibuchi M. Isolation of a pandemic O3:K6 clone of a *Vibrio parahaemolyticus* strain from environmental and clinical sources in Thailand. Appl Environ Microbiol 2000; 66: 2685–2689.
13. Matsumoto C, Okuda J, Ishibashi M, Iwanaga M, Garg P, Rammamurthy T, Wong HC, Depaola A, Kim YB, Albert MJ, Nishibuchi M. Pandemic spread of an O3:K6 clone of *Vibrio parahaemolyticus* and emergence of related strains evidenced by arbitrarily primed PCR and toxRS sequence analyses. J Clin Microbiol 2000; 38: 578–585.
14. Kim YB, Okuda J, Matsumoto C, Takahashi N, Hashimoto S, Nishibuchi M. Identification of *Vibrio parahaemolyticus* strains at the species level by PCR targeted to the *toxR* gene. J Clin Microbiol 1999; 37: 1173–1177.
15. Tada J, Ohashi T, Nishimura N, Shirasaki Y, Ozaki H, Fukushima S, Takano J, Nishibuchi M, Takeda Y. Detection of thermostable direct hemolysin gene (*tdh*) and the thermostable direct hemolysin-related hemolysin gene (*trh*) of *Vibrio parahaemolyticus* by polymerase chain reaction. Mol Cell Probes 1992; 6: 477–487.
16. Wong HC, Liu SH, Wang TK, Lee CL, Chiou CS, Liu DP, Nishibuchi M, Lee BK. Characteristics of *Vibrio parahaemolyticus* O3:K6 from Asia. Appl Environ Microbiol 2000; 66: 3981–3986.
17. Honda T, Iida T. The pathogenicity of *Vibrio parahaemolyticus* and the role of the thermostable direct hemolysin and related hemolysins. Rev Med Microbiol 1993; 4: 106–113.
18. Takahashi A, Sato Y, Shiomi Y, Cantarelli VV, Iida T, Lee M, Honda T. Mechanism of chloride secretion induced by thermostable direct hemolysin of *Vibrio parahaemolyticus* in human colonic tissue and intestinal epithelial cell line. J Med Microbiol 2000; 49: 801–810.
19. Bhoopong P, Palittapongarnpim P, Pomwised P, Kiatkittipong A, Kamruzzaman M, Nakaguchi Y, Nishibuchi M, Ishibashi M, Vuddhakul V. Variability of properties of *Vibrio parahaemolyticus* strains isolated from individual patients. J Clin Microbiol 2007; 45: 1544–1550.
20. Kamruzzaman M, Bhoopong P, Vuddhakul V, Nishibuchi M. Insertion sequence-mediated deletion of thermostable direct hemolysin gene (*tdh*) in *Vibrio parahaemolyticus*. Gene 2008; 421: 67–73.
21. Thongchan J, Bhoopong P, Yingkajorn M, Nishibuchi M, Vuddhakul V. Total number, virulence genes, and heterogeneity of *Vibrio parahaemolyticus* in a single shellfish. Science Asia 2013; 39: 230–235.
22. Shen X, Cai Y, Liu C, Liu W, Hui Y, Su YC. Effect of temperature on uptake and survival of *Vibrio parahaemolyticus* in oysters (*Crassostrea plicatula*). Int J Food Microbiol 2009; 136: 129–132.