All attempts to grow the acid-fast bacilli from the original bacillary suspension and the cell suspension at the end of the experiment on artificial culture media have failed. Subcutaneous inoculation of these materials to mice produced the typical lesions caused by rat-leprosy bacilli.

2. Animal-transmission of *M. leprae*

Eighty-five CF#1 mice were inoculated with living leprosy bacilli from an untreated patient with lepromatous leprosy into the left hind foot-pads, and twenty with heat-killed ones, using 10^3 to 10^4 bacilli per foot-pad.

After three to eight months the mice were sacrificed and both left and right hind foot-pads, lungs, livers, kidneys, spleens, and left inguinal lymph nodes were examined for presence of acid-fast bacilli.

Multiplication of the leprosy bacilli in the left hind foot-pads of mice was confirmed in fifty-eight of eighty-five mice inoculated with living bacilli. Acid-fast rods were also found in the tissues other than the inoculated foot-pads.

Discovery of acid-fast bacilli was 64.7 per cent in the mice inoculated with living ones and 50.0 per cent with killed ones.

And acid-fast rods were also found most frequently in the lymph nodes draining the inoculated foot-pads, but the number of bacilli were very small in most cases. The evidence presented suggests that the leprosy bacilli are carried from the site of inoculation to other sites of mice.

Cultures on bacteriological media favorable for the growth of most known mycobacterial species have not shown growth of the bacilli. The third passage of the bacilli is now under experiment.

CHEMOTHERAPY OF HUMAN LEPROSY — MAINLY, ON THE DDS-SENSITIVE STRAIN AND DDS-RESISTANT STRAIN OF MOUSE PASSAGES —

KAZUNARI NAKAMURA

National Institute for Leprosy, Tokyo,

Shepard, et al. and Rees, et al. have described the use of the infection of mouse foot-pads with *M. leprae* as a method for anti-microbial therapy of human leprosy, also, we have previously reported the activity of DDS against *M. leprae* in this system using isolates from Japanese patients.

Therefore, a comparison of the activity of DDS and other drugs against DDS-sensitive, and DDS-resistant strain was carried out in mice inoculated into the foot-pad with mouse passages, and then the inhibitory effect of DDS against the multiplication of *M. leprae* was performed in the mouse foot-pad system. The results obtained were as follows:

The multiplication of the so-called resistant strain was suppressed completely by DDS-administration with containing 0.5 mg per one gram of diet, these resistant

strains were mainly isolated from treated or relapsed patients with Promine- or DDS-administration for a long time.

Even the sensitive strain (no growth was observed in 0.1 mg diet/gram) was not completely suppressed in the case of DDS-administration which was given from the time of 3 to 6 months after the inoculation with *M. leprae*, these strains were mainly isolated from the untreated patients.

On the other hand, DDS-administration to the moderately resistant strain in the mouse foot-pad (slightly growth in 0.1 mg diet/gram) was continued in a period of about 14 to 20 months, this strain was reached to the plateau level (approximately 10^5 to 10^7 bacilli/foot-pad).

The inhibitory effect of Kanamycin and Streptomycin against DDS-resistant strain was not completely suppressed than the DDS-sensitive strain, which was received with each dosage of 0.4 mg per mouse three times a week from the time of 2 to 6 months after the inoculation with these strain.

Differences of the activity of anti-cancer drugs (Mitomycin C, Hydroxyurea, Toyomicin, 5-Fluorouracil) against DDS-resistant strains and DDS-sensitive strains were not observed in mouse foot-pad system.

MEDICAL REHABILITATION OF LEPROSY

TATSUJI TAMAI

*Department of Orthopaedic Surgery, Kumamoto University
Medical School, Kumamoto, Japan*

In Japan, a substantial number of leprosy patients has been rehabilitated to society from the leprosaria, owing to their dramatic response to promin and D. D. S., and to the improvement of orthopaedic treatment for disturbance of function and deformity of extremities.

The problem of personality in leprosy is very important not only for social recovery, but also medical rehabilitation as in the case of other crippeldness.

The observation of personality tells us that the patients differ a great deal from healthy persons in terms of somatic inferiority complex.

According to our experiences, medical rehabilitation of leprosy should be begun immediately after the disease is diagnosed. A success in the rehabilitation of leprosy can only result from cooperation of specialists for leprosy in different fields, besides basic medical treatments of the disease.

Physical therapy is a valuable treatment, using oil massage, vibra-bath, paraffin-bath, under-water exercise and electric treatment and so on. In the case of paralysis of the hands and feet, an early employment of physical therapy has been found extremity valuable and has contributed to recovery of paralysis and to prevention of deformities.

The reconstructive surgery is very effective not only to improve the activity of daily life of patients, but also to remove their inferiority complex, through the