Nutritional characteristics in leaves of plants native to tropical peat swamps and heath forests of Central Kalimantan, Indonesia

Sehat Jaya TUAH^{1,2)}, and Yahya Mika JAMAL²⁾ & Suwido Hester LIMIN²⁾

¹⁾ Graduate School of Agriculture, Hokkaido University, Kita 9, Nishi 9, Kitaku, Sapporo, 060-8589, Japan

²⁾ Center for International Cooperation in Management of Tropical Peatland, Palangka Raya University, Palangka Raya 73112, Central Kalimantan, Indonesia

ABSTRACT Thirteen mineral element concentrations in leaf samples of 478 native plants growing in peat swamps and heath forests of Central Kalimantan were determined. Chemical properties of surface soils in the study sites were also determined. In all sampling sites the mean concentrations of essential macro-elements were less variable than that of micro-elements. The chemical properties of surface soils necessitate that the peat and sandy podzolic soils are nutrient-limited. However, some native plants growing on these soils show a preference to accumulate excessive concentrations of particular micro-elements, such as Na, Mn, Al, and Si, in their leaves. The results of this study confirm that the concentrations of elements in plant leaves appear to be indicative of site variation and are the result of environmental adaptation. Al, Si, and Na were the principal elements determining the nutritional characteristics of the native plant leaves. However, only Al has clear inter-element relationships for each plant group. Therefore, the relationships between Al-Na and Al-Si in leaves reveal important strategies of plant adaptation to the nutrient-poor ecosystems of peat swamps and heath forests. Native plants belonging to the Eurosoid I evolutionary group had very high Al concentrations in their leaves, both in Lahei and Kalamangan peat swamp forests. *Combretocarpus rotundatus* (Anisophylleaceae family) was predominant in the study sites and, could be regarded as an Al accumulating species (leaf Al concentration within the range of 2,000 - 10,000 mg kg⁻¹). Al-accumulating species likely restrict the accumulation of other mineral elements in their leaves, whereas non Al-accumulating species tend to accumulate other mineral elements in their leaves. Although the mechanism for this eco-physiological adaptation remains unclear, it is suspected that Al-accumulating species (c.f. *C. rotundatus*) have developed efficient strategies enabling them to utilize low amounts of essential elements in order to be able to grow in a nutrient-limited environment. Information gathered about Al-accumulation by certain wild plants will be useful in selecting appropriate species to rehabilitate damaged peat swamp and heath forests.

Key words: peat swamp forest, heath forest, mineral element concentrations, native plants, *Combretocarpus rotundatus*, Al-accumulating species, phylogeny

INTRODUCTION

Indonesia contains the largest area of tropical peatland, estimated at 20 million hectares (Radjagukguk 1997) of which 2.5 million hectares are located in Central Kalimantan. Peatlands are wetland ecosystems that are characterized by the accumulation of organic matter, which is produced and deposited at a rate greater than its decomposition, leading to the formation of peat. Within tropical peatlands, inland and transitional peat have been recognized as being particularly nutrient-poor due to their very high acidity and organic matter content, especially lignin (Polak 1975). A lignin content of about 50 - 60% was observed in the peat soil from Central Kalimantan (Artiningsih 1999). High

aluminum (Al) concentration in soil solutions is also known to be the main factor restricting growth of plants on acidic soils (Rao *et al.* 1993; Watanabe & Osaki 2002). However, numerous native plants are well adapted to such environments and do not exhibit symptoms of nutrient deficiency or toxicity and, in fact, grow vigorously. This indicates that these plants have adaptation mechanisms to maintain their growth and development in the harsh environment of the lowland peatland ecosystem. Osaki *et al.* (1998) reported that some native plants such as *Melastoma malabatricum* and *Melaleuca cajuputi*, that grow in various adverse soils in Peninsular Thailand, have outstanding characteristics in relation to Al and Na accumulation in leaves, even though the Al and Na concentrations in the soil are very low. On the other hand, these native plants have low concentrations of Na and Al in soils with high Na and Al concentrations. It is assumed that the Al and Na accumulator plants have a tendency to exclude other mineral elements from their leaves, suggesting that these accumulator plants have developed efficient mechanisms to utilize small amounts of essential nutrients for their growth (Osaki *et al.* 2002).

Native plants constitute important components of ecosystems, as they recycle elements and provide transport pathways from the non-living to the living environment (Hamilton 1995). However, elements within plants are not cycled independently. Because they are involved in the basic structure and function of cells, these elements have similar biochemical pathways (Garten 1976). In an analysis of a wide array of plant traits, Grime *et al.* (1997) suggested that mineral nutrients be considered as the fundamental currency of vegetation processes, from the individual level to the ecosystem level. Thus, assuming that native plants can absorb and accumulate nutrients selectively and maintain a balance of intercellular concentrations to optimize their growth (Garten 1976), it is argued that the specific elemental concentrations of these plants reflect the nutrient concentration ratios in their substrate (Markert 1996).

Studies of elemental traits in native plants, particularly trees, are difficult because of plant size and longevity (Attiwill 1980) and differences in the storage of nutrients in particular plant organs, chemical fractions and cellular compartments (Chapin 1988). Most studies have, therefore, been based on changes in the concentrations of nutrients in various tissues, particularly leaves (Masunaga *et al.* 1997; Osaki *et al.* 1998; Pampasit *et al.* 2000). Because of strong interrelationships between foliar N, P, K, Ca, and Mg concentrations in some species (Garten 1976; Grime *et al.* 1997), Aerts & Chapin (2000) concluded that leaf nutrient concentrations are important determinants of plant function and development. Comparative studies of leaf nutrient concentrations in 54 species of non-vascular and vascular plants, as well as aquatic macrophytes (Garten 1976), 83 species of herbaceous flora (Thompson *et al.* 1997) and native plants of humid tropical lowlands (Osaki *et al.* 2000), have pointed out important relationships between mineral elements. These relationships derive mostly from both species and site specific characteristics. This study, therefore, aimed to clarify a general characteristic of mineral element concentration in leaves and also describe the relationship between the concentrations of mineral elements across a variety of native plant species and site conditions in tropical peat swamps and heath forests in Central Kalimantan, Indonesia.

MATERIALS AND METHODS

Study Sites

Study sites were established at three different locations: Kalampangan, Setia Alam, and Lahei (Fig. 1). The study site in the Kalampangan area comprised of a primary or natural peat swamp forest and a fire-damaged area of inland peat. A primary (natural) peat swamp forest site is defined as a relatively undisturbed area of peat swamp forest, whereas fire-damaged areas are defined as those areas affected by the massive wildfires of 1996-1997. The study site in Setia Alam area included a relatively undisturbed peat swamp forest, as well as a logged forest, whereas the Lahei site included two adjacent, contrasting, forest types of peat swamp and *kerangas* (heath) forest. In addition, the native plants in these three different sites grow in deep, intermediate, and shallow peat depths, respectively.

Fig. 1. Location of the study sites and field samplings in the peat swamps and heath forests in Central Kalimantan, Indonesia.

Species composition within the Kalampangan and Setia Alam sites, as determined by field survey, were relatively similar. The dominant tree species included: *Tetrameristra glabra*, *Calophyllum* sp., *Shorea* sp., *Combretocarpus rotundatus*, *Palaquium* sp., *Buchanania sessifolia*, *Syzygium* sp., *Dactylocladus stenostachys*, *Dyera costulata*, *Ilex cymosa*, *Tristaniopsis obovata*, and *Dyospyros* sp. The species composition at the Lahei site according to Suzuki *et al.* (1998) was dominated by *Buchanania sessifolia*, *Baccaurea macrocarpa*, *Gluta rugulosa*, *Shorea balangeran*, *Dyospyros* sp., *Semecarpus* sp., and *Ganua motleyana* in the peat swamp forest and *Calophyllum* sp., *Palaquium* sp., *Canarium* sp., and some important species of the dipterocarp family in the heath forest.

Sample Collection and Preparation

Mature and fully expanded leaves (including shoots) of some dominant tree species ≥ 10 cm in diameter

at breast height (DBH) were sampled. The upper leaves of several fern species were also collected, particularly within the fire-damaged area of the Kalampangan site.

The collected leaves were washed with deionized water, dried in an air-forced oven at 80°C for 72 hours, and then ground in a tungsten carbide vibrating mixer mill C.M.T. TI-100 (C.M.T. Co. Ltd., Japan). The plant species that were sampled and analyzed in this study are listed in Appendix 1.

Soil samples were also collected at each sampling site using a peat soil auger from Eijkelpamp Agrisearch Equipment (about 50 cm in depth). Five points were sampled at each study site, and the samples collected were subsequently mixed and air-dried at room temperature. The dried soils were ground and sieved through a 2 mm mesh.

Analytical Methods

Prior to determining the total elemental concentration of leaves, about 100 mg of finely ground samples were digested in a mixed solution of $\text{H}_2\text{SO}_4\text{-H}_2\text{O}_2$, according to the method used by Mizuno and Minami (1980). Total N concentration was determined by the semi-micro Kjeldahl method, whereas total concentrations of P, K, Ca, Mg, Na, Mn, Fe, Zn, Cu, Mo, Al, and Si were measured by Inductively Coupled Plasma Atomic Spectrometry (ICPS-7000, SHIMADZU, Japan). The total Si concentration reported in this paper does not include acid insoluble Si, because the digestion method used can only measure acid soluble Si. Fresh soil pH (1:2.5 = soil:deionized water; v/v) was measured in the field using a portable pH combination electrode (TOA Electronics Ltd., Japan). Exchangeable Ca, Mg, and K concentrations in the soils were determined by atomic absorption spectrophotometry (AA-6200, SHIMADZU, Japan) for Ca and Mg and flame photometry for K after extraction with 1N KCl (soil:KCl = 1:20). Atomic absorption spectrophotometry was also used to determine available Cu, Zn and Mn after extraction with 0.05 N HCl. The amount of available P in the soils was determined by the Bray II method, whereas N and C content were measured using a C-N Analyzer (SUMIGRAPH NC-1000, Japan).

Data and Statistical Analysis

The general tendency of mineral elements to accumulate in the plants studied was described according to the phylogenetic classification of each plant, as inferred from analysis of *rbcL*, *atpB*, and 18S rDNA sequences (Chase *et al.* 1993; Angiosperm Phylogeny Group 1998; Soltis *et al.* 1999). The INKEY online program of Watson & Dallwitz (<http://biodiversity.uno.edu/delta/>, version 14th December 2000) was also used for classification.

Statistical analysis was conducted using the SPSS program (SPSS Science, Chicago, USA).

RESULTS

Chemical Properties of the Soils

Surface soils in all the study sites were strongly acidic (KCl pH <3) and the KCl pH was consistently lower than the H_2O pH. The pH (H_2O) of peat soil ranged from 3.3 - 4.0, and was about 4.1 in sandy podzolic soil. Both H_2O pH and KCl pH were slightly higher in sandy podzolic soil than in peat soil. Highly acidic soil may have a low availability of essential nutrients. The chemical properties of surface soils in the study sites showed that there was less N and C, as well as available P, in sandy podzolic soil

than in the peat soil. Exchangeable Ca and Mg were found in moderate amounts in sandy podzolic soil, while exchangeable K was very low. Cu and Mn were found in relatively higher amounts in sandy podzolic soil than in the peat soil, while Zn was detected in moderate concentration. Compared to the other peat soil sampling sites, Kalampangan soil, having a deep peat layer (K2), has relatively high amounts of available P, C, as well as exchangeable Ca and Mg. In general, the chemical properties of surface soils in the study sites, as summarized in Table 1, indicate both peat and sandy podzolic soils are nutrient limited substrates, and most nutrients are likely stored in above-ground biomass. Therefore, it is evident that the native plants growing in these soils compete and have particular strategies for the acquisition of mineral nutrients for growth.

Table 1. Chemical properties of soil from surface to 50 cm depth in the study sites of peat swamps and heath forests in Central Kalimantan, Indonesia.

Sampling Site	Code	Soil type	pH(KCl)	pH(H ₂ O)	Bray II P ₂ O ₅ (mg kg ⁻¹)	N (%)	C (%)	Exchangeable cations			Cu	Zn	Mn
								Ca	Mg	K			
								(cmol _e kg ⁻¹)					
Lahei	L1	Peat (intermediate)	2.4	3.4	98	1.4	37.2	0.13	0.88	0.43	0.12	5.03	0.60
Lahei	L2	Sandy podzolic	2.7	4.1	54	0.5	18.9	0.16	0.55	0.34	0.40	3.37	1.85
Kalampangan	K1	Peat (deep)	2.2	3.3	96	1.3	38.1	0.14	0.79	0.60	0.16	6.89	1.70
Kalampangan	K2	Peat (deep)	2.2	3.7	167	1.1	39.7	0.41	1.58	0.64	0.01	4.15	0.40
Kalampangan	K3	Peat (shallow)	2.4	4.0	44	1.4	38.8	0.04	0.44	0.45	0.14	1.91	0.30
Kalampangan	K4	Peat (shallow)	2.2	3.7	77	1.3	38.2	0.15	0.77	1.23	0.05	3.13	0.10
Setia Alam	S1	Peat (shallow)	2.3	3.7	69	1.4	36.8	0.14	0.78	0.41	0.13	5.07	0.40
Setia Alam	S2	Peat (deep)	2.2	3.6	66	0.9	39.7	0.17	1.12	0.51	0.13	5.55	0.50

Foliar Element Concentrations of Native Plants Growing at Various Sites

The total concentrations of various macro- and micro-elements in the leaves of 478 native plant samples from the three study sites are compiled in Appendix 1. Elemental concentrations in the leaves varied remarkably among plant species growing in tropical peat swamps and heath forests in Central Kalimantan, Indonesia (Table 2). The concentration of some elements was high at certain sampling sites. Compared with the micro-elements, less variation occurred among the macro-elements, indicating that the maintenance of a cellular balance of these nutrients is not site specific. Analytical results (Table 2) show that macro-element concentrations (g kg⁻¹) varied within the range of 4.1-35.0 for N, 0.1-2.5 for P, 1.8-25.6 for K, 0.7-28.7 for Ca, and 0.4 -21.3 for Mg. The concentrations of various micro-elements (mg kg⁻¹) varied within the range of 1.4-3753 for Na, 4.5-553 for Fe, 2.3-2111 for Mn, 0.1-85 for Zn, 0.04-14 for Cu, 0.9-18 for Mo, 0.9-10019 for Al, and 0.4-3616 for Si. In spite of the heterogeneity of the study sites, P concentration remained consistently very low compared to the other macro-elements studied (N, K, Ca, and Mg).

The element concentrations in the leaves of native plants of the study sites were compared with nutrient concentrations in leaves of native plants (mostly trees) from the tropical rain forest in Pinang-pinang, West Sumatra, Indonesia (Masunaga *et al.* 1997), and also native plants from the tropical dry evergreen forest in Sakaerat, Thailand (Sirirattanakon 1999) (Table 2). It is shown that, except for Mg, the macro-elements measured (N, P, K, and Ca) in the plant leaves of the study sites were lower than that of the tropical plant leaves in West Sumatra and Sakaerat. However, the concentrations of micro-elements varied greatly by site and specie. Al and Si, in particular, had a clear preference to accumulate in a considerably high concentration in some native plants' leaves as indicated in all three

different locations (Table 2). Therefore, certain micro-elements, especially Al and Si, are likely to be the key elements of native plant growth in nutrient-limited environments. In this study, element concentrations, especially that of micro-elements, in leaves of native plants varied greatly. Several native plants showed a preference to accumulate particular elements such as Na, Mn, Al, and Si in great concentrations, above the average concentration required for adequate plant growth (Appendix 1). In addition, concentrations of these micro-elements in plant leaves had a high coefficient of variation (CV), indicating that excessive absorption and/or accumulation is a specific characteristic of certain plants. In general, the average concentration of macro-elements in the leaves of the plants studied was greatest for N followed by K, Ca, Mg, P; for micro-elements the greatest concentration was that of Na, followed by Si, Al, Mn, Fe, Mo, Zn, Cu (Table 2).

Table 2. Magnitude of elemental concentrations in leaves of native plants from several sites in Central Kalimantan in comparison to the concentration of mineral elements in leaves of tropical native plants from two different locations: Pinang-pinang, West Sumatra (Indonesia) and Sakaerat, Changwat Nakhon Ratchasima (Thailand). In parenthesis, n = the total number of plant samples from all sampling sites.

Parameters	N	P	K	Ca	Mg	Na	Fe	Mn	Zn	Cu	Mo	Al	Si
	(g kg ⁻¹)					(mg kg ⁻¹)							
Native plants in Central Kalimantan, Indonesia ¹⁾													
Mean±SE (<i>n</i> = 478)	12.6±0.19	0.5±0.01	8.0±0.18	6.5±0.18	2.8±0.09	506±21.5	43±1.7	190±12.4	8.4±0.4	3.3±0.1	8.6±0.2	269±48.1	381±28.7
Min.	4.1	0.1	1.8	0.7	0.4	1.4	4.5	2.3	0.1	0.04	0.9	0.9	0.4
Max.	35.0	2.5	25.6	28.7	21.3	3753	553	2111	85	14	18	10019	3616
CV (%)	33	65	49	60	69	93	85	143	115	75	42	390	165
Native plants in West Sumatra, Indonesia ²⁾													
Mean±SE (<i>n</i> = 608)	18.1±0.19	1.0±0.02	9.4±0.21	16.9±0.42	2.6±0.06	147±2.3	156±3.3	532±21.7	19±1.1	10±0.2	nd	2043±198	6938±361
Min.	8.6	0.4	1.0	0.8	0.2	131	38	4.8	0.4	0.8	nd	1.0	248
Max.	46.6	4.3	34.4	81.7	10.5	861	851	3574	309	64	nd	36924	89984
CV (%)	26	42	54	61	55	39	52	100	137	54	nd	239	128
Native plants in Sakaerat, Thailand ³⁾													
Mean±SE (<i>n</i> = 198)	15.9±0.41	1.0±0.05	11.0±0.54	9.8±0.29	2.6±0.12	43±3.7	121±10.9	524±36.6	26±2.4	9±0.3	nd	2881±313	nd
Min.	6.9	0.1	0.6	0.7	0.3	trace	trace	12.8	2.9	0.4	nd	trace	nd
Max.	31.3	3.7	34.4	25.3	7.2	266	1149	2376	286	27	nd	16568	nd
CV (%)	36	68	70	42	63	122	127	98	126	53	nd	153	nd

1) Average of all native plant samples collected in the tropical peat swamps and heath forests in Central Kalimantan, Indonesia (this study).

2) Average of all native plant samples collected in the tropical rain forest in Pinang-pinang, West Sumatra, Indonesia (Masunaga *et al.* 1997).

3) Average of all native plant samples collected in the tropical dry evergreen forest in Sakaerat, Thailand (Sirirattanakorn 1999).

nd = no data.

Notwithstanding the variability between native plant species, mean concentrations of mineral elements were statistically compared between sampling sites (Table 3). It is shown that the concentrations of various elements measured in the native plants' leaves differed significantly between the study sites. However, Mg and Al concentrations in the native plants' leaves did not differ between the sites, while leaf P concentration was significantly lower in the native plants of both Kalampangan (K1,2,3&4) and Setia Alam (S1&2) than that of the native plants in the Lahei peat swamp and heath forests (L1 and L2). Relatively high concentrations of Na, Fe, Zn, Al, and Si in leaves of the native plants in the peat swamp forest in Lahei (L1) indicate an accumulative behavior of some native plant species at this site. In addition, concentrations of some elements were significantly low in the plant

leaves, because in most cases the tropical acidic soils are poor in nutrients (Table 1). Despite this fact, undisturbed natural forests seldom reveal symptoms of mineral deficiencies which are typical indicators of degradation. It seems that nutrient balances in undisturbed systems are in a “steady state”, and losses, if they occur, are more or less “neutralized” by input rate (Proctor 1987).

Table 3. Concentration of mineral elements (mean $\pm$ SE) in mature leaves of native plants growing on various locations in Central Kalimantan. In parentheses, n = number of plant samples. Different letters in the same row indicate a statistically significant difference (*LSD*-test) among sampling sites at $p<0.05$.

Element*	Sampling site **			
	L1 ($n = 47$)	L2 ($n = 52$)	K1,2,3&4 ($n = 323$)	S1&2 ($n = 56$)
N	14.7 $\pm$ 0.58 ^a	12.8 $\pm$ 0.53 ^b	12.6 $\pm$ 0.24 ^b	10.8 $\pm$ 0.35 ^c
P	0.8 $\pm$ 0.05 ^a	0.7 $\pm$ 0.06 ^a	0.4 $\pm$ 0.01 ^b	0.5 $\pm$ 0.04 ^b
K	7.0 $\pm$ 0.63 ^b	5.5 $\pm$ 0.35 ^d	8.7 $\pm$ 0.22 ^a	6.7 $\pm$ 0.40 ^c
Ca	7.9 $\pm$ 0.53 ^a	7.0 $\pm$ 0.61 ^b	6.0 $\pm$ 0.19 ^c	8.0 $\pm$ 0.72 ^a
Mg	3.4 $\pm$ 0.31 ^a	2.6 $\pm$ 0.24 ^a	2.8 $\pm$ 0.11 ^a	2.6 $\pm$ 0.24 ^a
Na	1311 $\pm$ 89.6 ^a	557 $\pm$ 23.5 ^c	336 $\pm$ 11.3 ^d	737 $\pm$ 99.0 ^b
Fe	66 $\pm$ 3.8 ^a	60 $\pm$ 3.6 ^b	38 $\pm$ 2.18 ^c	30 $\pm$ 1.54 ^d
Mn	264 $\pm$ 51.1 ^b	352 $\pm$ 54.2 ^a	142 $\pm$ 11.3 ^c	271 $\pm$ 46.8 ^b
Zn	19.3 $\pm$ 2.3 ^a	9.6 $\pm$ 1.1 ^b	7.6 $\pm$ 0.41 ^c	3.0 $\pm$ 1.29 ^d
Cu	4.1 $\pm$ 0.43 ^b	4.5 $\pm$ 0.28 ^a	3.1 $\pm$ 0.13 ^c	2.5 $\pm$ 0.26 ^d
Mo	5.4 $\pm$ 0.09 ^d	6.0 $\pm$ 0.13 ^c	8.7 $\pm$ 0.20 ^b	12.4 $\pm$ 0.25 ^a
Al	380 $\pm$ 106.9 ^a	69 $\pm$ 20.5 ^a	312 $\pm$ 68.5 ^a	114 $\pm$ 51.8 ^a
Si	1785 $\pm$ 132.1 ^a	127 $\pm$ 40.6 ^c	279 $\pm$ 19.3 ^b	28 $\pm$ 5.20 ^d

[*] Concentrations of macro-elements (N, P, K, Ca and Mg) and micro-elements (Na, Fe, Mn, Zn, Cu, Mo, Al and Si) expressed in g kg⁻¹ and mg kg⁻¹, respectively.

[**] Peat swamp forest in Lahei (L1); heath forest in Lahei (L2); peat swamp forest in Kalampangan (K1,2,3&4) and peat swamp forest in Setia Alam (S1&2).

In order to clarify the underlying factors associated with elemental characteristics of the native plants, a Principle Component Analysis was performed to derive the grouping pattern of the native plants studied. The three principal components: Al, Si, and Na (Fig. 2), accounted for 62% (Component-1), 25% (Component-2) and 8% (Component-3) of the total variance, respectively (Component Matrix is not shown). Based on these three principal components, all the native plants studied were categorized into three major groups with the Cluster Analysis extraction method. These grouping patterns were predominantly determined by the Al concentration in the native plants' leaves. Therefore, the three major groups mentioned above were separated into groups of native plant species with relatively low, moderate, and notably high Al-concentrations. Figs. 3, 4 and 5 demonstrate a typical relationship between Al, Si and Na, as well as between these three elements and the other elements. However, only Al clearly shows the inter-element relationships for each plant group (Fig. 3). Thus, Al-accumulating species, the third group, are likely restricting the accumulation of other mineral elements in their leaves, whereas non Al-accumulating species, the first and second groups, tend to accumulate other mineral elements in their leaves. Although the mechanism for their eco-physiological adaptation remains unclear, it is suspected that Al accumulating species (c.f. *C. rotundatus*) have developed efficient strategies to utilize lower amounts of essential elements, enabling them to thrive in

the strongly acidic soil of the peat swamps and heath forests.

Fig. 2. Plot of mineral element covariance extracted according to a Principal Component Analysis. Component 1, 2 and 3 associated with the concentrations of Al, Si and Na which accounted for 62%, 25% and 8% of the total variance, respectively.

Fig. 3. Relationships between the concentrations of Al (mg kg^{-1}) and other mineral elements in leaves of native plant species in Central Kalimantan, Indonesia. Concentrations of macro elements (N, P, K, Ca and Mg) and micro elements (Na, Fe, Mn, Zn, Cu, Mo and Si) were expressed in g kg^{-1} and mg kg^{-1} , respectively. (○: group of native plant species with relatively low Al-concentration; □: group of native plant species with moderate Al-concentration; and ◆: group of native plant species with notably high Al-concentration).

Fig. 4. Relationships between the concentrations of Si (mg kg^{-1}) and other mineral elements in leaves of native plant species in Central Kalimantan. Concentrations of macro elements (N, P, K, Ca and Mg) and micro elements (Na, Fe, Mn, Zn, Cu, Mo and Al) were expressed in g kg^{-1} and mg kg^{-1} , respectively. (○: group of native plant species with relatively low Al-concentration; □: group of native plant species with moderate Al-concentration; and ◆: group of native plant species with notably high Al-concentration).

Fig. 5. Relationships between the concentrations of Na (mg kg^{-1}) and other mineral elements in leaves of native plant species in Central Kalimantan. Concentrations of macro elements (N, P, K, Ca and Mg) and micro elements (Fe, Mn, Zn, Cu, Mo, Al and Si) were expressed in g kg^{-1} and mg kg^{-1} , respectively. (○: group of native plant species with relatively low Al-concentration; □: group of native plant species with moderate Al-concentration; and ◆: group of native plant species with notably high Al-concentration).

Accumulation of Elements in Plant Leaves with Different Phylogeny

The accumulation of an element in plant tissue may be an environmental adaptation. The accumulative behavior of native plants may therefore be assumed to be ecologically and genetically related. In Figs. 6 and 7 the patterns of element accumulation in the leaves of the native plants were shown for every element studied within the sampling sites. A comparison was made among native plants of different phylogenetic groups, representing plants that evolved a long time ago and more recently evolved plants. Most of the plants studied in all the sampling sites are members of the Ericales, Eurosid II, and Eurosid I phylogenetic groups of recently evolved plants (Appendix 1).

Fig. 6. Mean concentrations of N, P, K, Ca and Mg (g kg⁻¹) in leaves of native plants from four sampling locations with respect to phylogenetic group. Bars indicate SE value.

Fig. 7. Mean concentrations of Na, Fe, Mn, Zn, Cu, Mo, Al and Si (mg kg^{-1}) in leaves of native plants from four sampling locations with respect to phylogenetic group. Bars indicate SE value.

There was great specie specific variation in leaf elemental concentrations of different phylogenetic groups and sampling sites. Mean concentrations of the essential macro-elements (N, P, K, Ca, and Mg) varied widely among sampling sites, but were relatively less divergent between phylogenetic groups. However, mean concentrations of the micro-elements (Na, Fe, Mn, Zn, Cu, Mo, Al, and Si) demonstrated a considerably high variation between all taxonomic groups and sampling sites. Native plants belonging to the Pteridophyte group showed a high accumulation of K in their leaves, whereas the Ericales plants of the peat swamp forest in Lahei showed a high accumulation of Mg. Some recently evolved native plants, mainly of the Euasterid I, Ericales, Eurosoid II, Eurosoid I, and the Magnoliids group, seemed to accumulate particular micro-elements such as Na, Fe, Mn, Cu, Mo, Al, and Si. However, no particular group had especially excessive accumulation. Several taxa within the Eurosoid I evolutionary group had a considerably high Al concentration in their leaves in plants of both the Lahei and Kalampangan peat swamp forests (Fig. 7).

Effect of Site Variation on Leaf Elemental Concentrations

Because the array of element concentrations in leaves may reflect the concentration ratios of elements in substrates, it is argued that the properties of individual elements in plants may influence plant traits. The possibility of site-specificity of elemental concentrations in the leaves of the native plants was investigated with respect to plant evolutionary groups (Figs. 8, 9, and 10). Native peat swamp forest at the Kalampangan site (K1&3) was investigated for this comparison, because of its relatively greater abundance and diversity of plant species compared with the other sites.

Fig. 8. Relationships of N, P, K, Ca and Mg concentrations (g kg^{-1}) in leaves of native plants with respect to phylogenetic group and sampling location. Bars indicate SE value. L1: peat swamp forest in Lahei; L2: heath forest in Lahei; K1&3: undisturbed peat swamp forest in Kalampangan; K2&4: fire-damaged peat swamp forest in Kalampangan; S1&2: peat swamp forest in Setia Alam. (●: Euasterid II; ■: Euasterid I; ▲: Ericales; ○: Eurosoid II; □: Eurosoid I; △: Magnoliales+Laurales).

Fig. 9. Relationships of Fe, Zn, Cu and Mo concentrations (mg kg^{-1}) in leaves of native plants with respect to phylogenetic group and sampling location. Bars indicate SE value. L1: peat swamp forest in Lahei; L2: heath forest in Lahei; K1&3: undisturbed peat swamp forest in Kalampangan; K2&4: fire-damaged peat swamp forest in Kalampangan; S1&2: peat swamp forest in Setia Alam. (●: Euasterid II; ■: Euasterid I; ▲: Ericales; ○: Eurosoid II; □: Eurosoid I; △: Magnoliales+Laurales).

Fig. 10. Relationships of Mn, Na, Al and Si concentrations (mg kg^{-1}) in leaves of native plants with respect to phylogenetic group and sampling location. Bars indicate SE value. L1: peat swamp forest in Lahei; L2: heath forest in Lahei; K1&3: undisturbed peat swamp forest in Kalampangan; K2&4: fire-damaged peat swamp forest in Kalampangan; S1&2: peat swamp forest in Setia Alam. (●: Euasterid II; ■: Euasterid I; ▲: Ericales; ○: Eurosoid II; □: Eurosoid I; △: Magnoliales+Laurales).

Mean concentrations of essential macro-elements in particular phylogenetic groups were less site-specific than micro-elements (Fig. 8). Most of the sites under peat swamp forest sites (L1, K2&4, and S1&2) showed no remarkable differences, with the exception of the heath forest (L2). However, the mean concentrations of Ca and Mg in the leaves from all study sites were highly site-specific (Fig. 8). In terms of micro-elements, the mineral concentrations were immensely affected by site heterogeneity (Figs. 9 and 10). In both the Kalampangan and Lahei peat swamp forests, plants of the Eurosoid I group showed similar Al accumulation. It is therefore necessary to be aware that the species of the Eurosoid I phylogeny have a preference to accumulate excessive concentrations of Al in their leaves.

DISCUSSION

Concentrations of various elements in the leaves of native plants growing on peat and sandy podzolic soils in tropical peat swamps and heath forests were examined in this study. Growth of native plants is generally limited by the availability of inorganic nutrients in soils and our results confirm that the concentrations of elements in plant leaves appear, to some extent, to be reflective of the site conditions. From a biochemical point of view, all plant species should require a similar quantity of nutrients to construct a given amount of tissue (Chapin 1980; Alonso & Herrera 2001). However, differences do exist in tissue concentrations because of heterogeneous distribution of nutrients in soils and varying uptake efficiency in individual plants (Pugnaire 2001). Adaptation to varying nutrient availability in the substrates, therefore, has presumably led to disparities of nutrient concentrations between different native plants' organs, species and ecosystems. As shown in Appendix 1, the concentrations of individual elements differed widely between species and even between individuals of the same species, as well as between sampling sites. Because mineral nutrients have different functions within plants, their leaf concentrations may differ by orders of magnitude from essential macro-elements to trace elements (Appendix 1, Table 2 and 3).

Table 2 shows that the plants in natural systems demonstrate high inter- and intra-specific variability in nutrient concentrations as indicated by the coefficient of variation (CV) values of each element measured. Some native plant species show a preference for excessive accumulation of particular micro-elements, such as Na, Mn, Al, and Si; while certain species had very low concentrations of other micro-elements (especially Zn, Cu, and Mo) in their leaves. However, notwithstanding the natural variability of nutrient concentrations, native plants grown in adverse environments (Table 2) show a relatively consistent concentration of nutrients in their leaves.

The chemical properties of the soils within the study sites (Table 1) are characterized by a very low availability of most nutrients, particularly Cu, Zn, and Mn. This is probably related to the conditions of peat and sandy podzolic soils, in which nutrients are profoundly limited. Under ideal conditions, the amount of nutrients available in a medium correspond proportionally to the amount of nutrients taken up by the plants (Markert 1996). However, this is not the case in adverse conditions, as observed in this study. Thus, though the physiological mechanism is still unclear, it is assumed that the native plants have developed an efficient strategy enabling them to grow with only limited resources on the highly acidic soils of the peat swamps and heath forests. Mineral element concentrations in plant leaves are variable, not genetically predetermined, and the ability to grow with both low and high accumulation of

particular nutrients may be an important aspect of an individual plant's adjustment to its environment (Alonso & Herrera 2001).

To examine the possible effects of environmental variability on mineral element concentrations in the leaves of the plants studied, a comparison was made between sampling sites (Table 3). It is shown that Mg and Al concentrations were not significantly affected by environmental variability. Yet, the mean concentrations of Na, Fe, Zn, Al, and Si in the leaves of plants of the peat swamp forest in Lahei (L1) were notably higher compared to the other sites. The accumulation of elements in plant tissues is a dynamic process, and therefore the element concentrations of individual plants of inter- or intra-species may vary considerably from one site to another (Verhoeven *et al.* 1996). It has widely been acknowledged that mineral nutrients have various functions within plants, leading to differences in their concentrations over time (Pugnaire 2001).

Considering that the concentrations of nutrients in leaves are important indicators of the functioning of plant species in their ecosystem (Aerts & Chapin 2000), it is necessary to understand the principal factors regarding elemental concentrations in the leaves of plants that may lead to further understanding the plant traits and/or natural groupings. Factor analysis (Fig. 2) specified that three elements, Al, Si, and Na, were the main determinants of the characteristics of mineral elements in the leaves in this study. Among these three principal components, Al was the main component, followed by Si and Na. Figs. 3, 4 and 5 show the relationships between Al, Si and Na, as well as the relationships between these three elements and the other elements. However only Al has clear element interrelationships for each plant group. It is apparent that all the native plants are clearly separated into three groups with relatively low (first group), moderate (second group) and notably high (third group), leaf Al concentrations (Fig. 3). The third group was dominated by the species *C. rotundatus* (data not shown). It is evident that Al-accumulating species (the third group) are likely restricting the accumulation of other mineral elements in their leaves, whereas non Al-accumulating species (the first and second group) tend to accumulate other mineral elements in their leaves. Seemingly, Al accumulating species have developed efficient mechanisms to utilize small amounts of essential elements in order to grow in the nutrient-limited soils of peat swamps and heath forests, though a specific explanation of this adaptation mechanism remains vague (Wanatabe & Osaki 2002).

From these results, we conclude that Al, Si, and Na are likely the most important micro elements for the many native plants well-adapted to peat and podzolic sandy soils. In a study of several plant species well-adapted to temperate and tropical acid soils, Osaki *et al.* (1997) found Al to be beneficial to plant growth. In this study, we also found an antagonistic relationship between Al and Na, and a synergistic interaction between Al and Si, particularly in native plant species that have high concentrations of Al and Si in their leaf tissue. An antagonistic relationship between the concentrations of Al and Na was reported by Osaki *et al.* (1998) as an adaptation strategy of plants in nutrient poor, acidic tropical soils. Meanwhile, there is a possibility that there is a synergistic relationship between Al and Si concentrations in certain plants in which Si alleviates the toxic effects of Al on plants (Hodson and Evans 1995; Cocker *et al.* 1998; Masunaga *et al.* 1998). In their study on the relationship between Al and Si accumulation in *Faramea marginata* (family Rubiaceae), Britez *et al.* (2002) found a positive correlation between Al and Si concentrations and also that these two elements showed a similar distribution in leaf and stem tissues. It was also reported that the Al and Si elution patterns were similar to those of aluminum silicate. These results indicate a formation of an Al-Si complex occurred in

the shoot tissues of *F. marginata*, which may be related to the internal Al detoxification mechanism. Silicon has also been demonstrated to inhibit Al penetration into the root cortex of *Sorghum bicolor*, suggesting that Si forms a complex with Al in either the media or roots and contributes to the detoxification of Al (Hodson & Sangster 1993). In a forest ecosystem, such as in the peat swamps and heath forests, Al is present in plant tissue and thus obviously taken up by the plant. Through litterfall and root dieback, plants redistribute Al into the soil by way of decomposition (Rustad & Cronan 1989). As soil organic matter originates from non-living plant materials, such as litterfall and root dieback, it is presumable that Al in peat soil forms a complex with soluble organic matter. Al in this form is not toxic to plants (Kinraide 1991) and complexation of Al by organic matter appears to be a very important mechanism of Al detoxification of soil solution (Kochian 1995). In addition, during decomposition of plant debris, a whole range of organic compounds are released from the debris and/or synthesized by decomposer microorganisms (Haynes & Mokolobate 2001). Al can bind strongly to many of these compounds (Stevenson & Vance 1989). Therefore, it is evident that various native plants in humid tropical regions with strongly acidic soil can vigorously grow with a high accumulation of Al in their leaves (cf. Osaki *et al.* 1995 & 1997).

Although the accumulation of an element in plant tissues does not necessarily mean that it plays a prominent physiological role (Bargagli 1998), to some extent it may indicate that plants respond differently to environmental variability (Alonso & Herrera 2001), and therefore the accumulative behavior of a plant may also have genetically predetermined origins (Markert 1996). The patterns of mineral element accumulation in the leaves of native plants in this study were elucidated according to plant evolutionary classification. There was considerable specie-specific and site-specific variation in elemental acquisition (Figs. 6 and 7). The mean concentrations of essential macro-elements were less divergent between phylogenetic groups. In contrast, the mean concentrations of micro-elements varied markedly among taxonomic groups and sites. A strong relationship between leaf nutrient concentration and habitat and phylogeny was reported in several taxa of herbaceous flora (Thompson *et al.* 1997). Taxa that are restricted to fertile, lowland habitats (e.g. *Chenopodium*, Urticaceae) tend to have nutrient-rich foliage, whereas those that are mostly confined to upland, infertile habitats (e.g. Cyperaceae, Ericaceae) have nutrient-poor foliage. Thompson *et al.* (1997) therefore argued that leaf nutrient concentrations are essentially a specie-specific trait. In this study, native plants belonging to the Eurosid I phylogeny group had very high Al accumulation in their leaves in both the Kalampangan and Lahei peat swamp forest (Fig. 7). In the Lahei peat swamp forest, Al accumulator species were recorded for only one taxon (Euphorbiaceae), whereas in Kalampangan, it was recorded for two taxa (Anisophylleaceae and Lycopodiaceae). Of these taxa, *C. rotundatus* (Anisophylleaceae) was the species most widely distributed, particularly in the fire-damaged area. In a review of Al accumulation in angiosperms, Jansen *et al.* (2002) reported that the species of the woody plant genus *Combretocarpus* are Al accumulators. As Al in peat soil is presumably bound into organic matter as chelate compounds, it is posited that *C. rotundatus* and other Al-accumulating species have developed their particular ability to absorb and accumulate Al through adaptation mechanisms in adverse environments. Such mechanisms are thought to involve the formation of a complex with an Al ion, as well as intracellular and tissue compartmentalization (Britez *et al.* 2002; Watanabe & Osaki 2002).

Both from an ecological and a nutrient physiological point of view, chemical characterization of ecosystems provides important information on the fate of elements in different systems. Thus, the

pattern of nutrient accumulation could be shaped by environmental and evolutionary factors. Heterogeneity of the sites was examined in this study in relation to elemental concentrations, and the results show that environmental elemental concentrations have a significant impact on leaf element concentrations within each taxonomic group (Figs. 8, 9, and 10). Although the concentrations of Mg and Ca in plant leaves tended to be site-specific, as also reported by Alonso & Herrera (2001), this may not occur consistently in the case of habitats with a larger variety of native plant species. It is argued that the main factors influencing mineral element concentrations are the inherent characteristics of each individual specie. This is mainly because native plants can absorb and accumulate nutrients selectively and maintain a balance of intercellular element concentrations to optimize their growth in spite of low nutrient availability (Garten 1976). Apparently, during their subsequent growth, native plants adjust their elemental composition with respect to the nutritional status of their environment (soil). Also the native plant species that are adapted to nutrient limited forests have developed adaptation strategies which include acquiring high concentrations of Al, Si, and Na in their leaves. This information is considered necessary for the selection of appropriate plant species to rehabilitate damaged peat swamps and heath forests. However, further research is needed to clarify physiological aspects of plant adaptation mechanisms, especially plant species that are considered Al accumulators.

ACKNOWLEDGMENTS We express our gratitude to Professor Mitsuru OSAKI for his invaluable support and encouragement. The assistance of our colleagues at Palangka Raya University during field sampling is greatly acknowledged. We also thank two anonymous reviewers for their critical comments. This research was supported by a research grant from the Japan Society for the Promotion of Science (JSPS) under a Core University Program between Hokkaido University, Japan and R & D Center of Biology, LIPI, Indonesia.

REFERENCES

- Aerts, R. & Chapin III, F. S. 2000. The mineral nutrition of wild plants revisited: a re-evaluation of processes and patterns. *Advances in Ecological Research* **30**:1-67.
- Alonso, C. & Herrera, C.M. 2001. Patterns made patterns: variation and covariation of leaf nutrient concentrations within and between populations of *Prunus mahaleb*. *New Phytologist* **150**: 629-640.
- Artiningsih, T. 1999. Decomposition of organic matter in peat soil by fungi. Environmental Conservation and Land Use Management of Wetland Ecosystem in Southeast Asia (Annual Report for April 1998 - March 1999 of the Core University Program between Hokkaido University, Japan and R&D Center of Biology, LIPI, Indonesia), pp. 229.
- Attiwill, P. M. 1980. Nutrient cycling in a *Eucalyptus obliqua* (L'Hérit) forest (IV): nutrient uptake and nutrient return. *Australian Journal of Botany* **28**: 199-222.
- Bargagli, R. 1998. Trace elements in terrestrial plants: an ecophysiological approach to biomonitoring and biorecovery, pp. 1-47 Springer-Verlag Berlin Heidelberg and R.G. Landes Company, Georgetown, TX, USA.
- Britez, R.M., Watanabe, T., Jansen, S., Reissmann, C.B. & Osaki, M. 2002. The relationship between aluminum and silicon accumulation in leaves of *Faramea marginata* (Rubiaceae). *New Phytologist* **156**: 437-444.
-

- Chapin, F. S. III. 1980. The mineral nutrition of wild plants. *Annual Review of Ecology and Systematics* **11**: 233-260.
- Chapin, F. S. III. 1988. Ecological aspects of plant mineral nutrition. *Advances in Mineral Nutrition* **3**: 61-191.
- Chase, M. W., Soltis, D. E., Olmstead, R. G., Morgan, D., Les, D. H., Mishler, B. D., Duvall, M. R., Price, R. A., Hills, H. G., Qiu, Y. L., Kron, K. A., Rettig, J. H., Conti, E., Palmer, J. D., Manhart, J. R., Sytsma, K. J., Michaels, H. J., Kress, W. J., Karol, K. G., Clark, W. D., Hedren, M., Gaut, B. S., Jansen, R. K., Kim, K. J., Wimpee, C. F., Smith, J. F., Furnier, G. R., Strauss, S. H., Xiang, Q. Y., Plunkett, G. M., Soltis, P. S., Swensen, S. M., Williams, S. E., Gadek, P. A., Quinn, C. J., Eguiarte, L. E., Golenberg, E., Learn Jr., G. H., Graham, S. W., Barrett, S. C. H., Dayanandan, S. & Aldert, V. A. 1993. Phylogenetics of seed plants: an analysis of nucleotide sequences from the plasmid gene *rbcL*. *Annals of the Missouri Botanical Garden* **80**: 528-580.
- Cocker, K. M., Evans, D. E. & Hodson, M. J. 1998. The amelioration of aluminum toxicity by silicon in higher plants: solution chemistry or an in planta mechanism? *Physiologia Plantarum* **104**: 608-614.
- Garten, C. T. 1976. Correlation between concentrations of elements in plants. *Nature* **261**: 686-688.
- Grime, J. P., Thompson, K., Hunt, R., Hodgson, J. G., Cornelissen, J. H. C. & Rorison, I. H. 1997. Integrated screening validates primary axes of specialization in plants. *Oikos* **79**: 259-281.
- Hamilton, E. I. 1995. State of the art of trace element determinations in plants matrices: determination of the chemical elements in plant matrices, an overview. *The Science of the Total Environment* **176**: 3-14.
- Haynes, R.J. & Mokolobate, M.S. 2001. Amelioration of Al toxicity and P deficiency in acid soils by additions of organic residues: a critical review of the phenomenon and the mechanisms involved. *Nutrient Cycling in Agroecosystems* **59**: 47-63.
- Hodson, M. J. & Evans, D. E. 1995. Aluminum/silicon interactions in higher plants. *Journal of Experimental Botany* **46**: 161-171.
- Jansen, S., Broadley, M., Robbrecht, E. & Smets, E. 2002. Aluminum hyperaccumulation in angiosperms: a review of its phylogenetic significance. *The Botanical Review* **68**: 235-269.
- Kinraide, T.B. 1991. Identify of the rhizotoxic aluminum species. *Plant and Soil* **134**: 167-178.
- Kochian, L. V. 1995. Cellular mechanisms of aluminum toxicity and resistance in plants. *Annual Review of Plant Physiology and Plant Molecular Biology* **46**: 237-260.
- Markert, B. 1996. Instrumental element and multi-element analysis of plant samples: methods and applications (Translated by B. Haderlie), pp 9-21. John Wiley & Sons, Chichester.
- Masunaga, T., Kubota, D., Hotta, M. & Wakatsuki, T. 1997. Nutritional characteristics of mineral elements in tree species of tropical rain forest, West Sumatra, Indonesia. *Soil Science and Plant Nutrition* **43**: 405-418.
- Masunaga, T., Kubota, D., Hotta, M. & Wakatsuki, T. 1998. Mineral composition of leaves and bark in aluminum accumulators in a tropical rain forest in Indonesia. *Soil Science and Plant Nutrition* **44**: 347-358.
- Mizuno, N. & Minami, M. 1980. The use of $\text{H}_2\text{SO}_4\text{-H}_2\text{O}_2$ for destruction of plants matter as a preliminary to determination of N, K, Mg, Ca, Fe, Mn. *Jpn. Journal of Soil Science and Plant Nutrition* **51**: 418-420 (in Japanese with English Summary).
- Osaki, M., Sittibush, C. & Nuyim, T. 1995. Nutritional characteristics of wild plants grown in peat and

- acid sulfate soils distributed in Thailand and Malaysia. *In*: Vijarsorn, P., Suzuki, K., Kyuma, K., Wada, E., Nagano, T. & Takai, Y. (eds.). A tropical swamp forest ecosystem and its greenhouse gas emission, pp 63-67. Nodai Research Institute Tokyo University of Agriculture, Tokyo.
- Osaki, M., Watanabe, T. & Tadano, T. 1997. Beneficial effect of aluminum on growth of plants adapted to low pH soil. *Soil Science and Plant Nutrition* **43**: 551-563.
- Osaki, M., Watanabe, T., Ishizawa, T., Nilnond, C., Nuyim, T., Shinano, T., Urayama, M. & Tuah, S. J. 2003. Nutritional characteristics of the leaves of native plants growing in adverse soils of humid tropical lowlands. *Plant Foods for Human Nutrition*: (in press).
- Osaki, M., Watanabe, T., Ishizawa, T., Nilnond, C., Nuyim, T., Sittibush, C. & Tadano, T. 1998. Nutritional characteristics in leaves of native plants grown in acid sulfate, peat, sandy podzolic, and saline soils distributed in Peninsular Thailand. *Plant and Soil* **201**: 175-182.
- Pampasit, S., Khamyong, S., Breulmann, G., Ninomiya, I. & Ogino, K. 2000. Elements concentrations in tree species of tropical hill evergreen forest, Northern Thailand. *Tropics* **9**: 287-308.
- Polak, B. 1975. Character and occurrence of peat deposits in the Malaysian tropics. Pp. 71-81 in G. J. Bartstra & Caspare, W. A. (eds). Modern quaternary research in Southeast Asia. A.A. Belkema, Rotterdam.
- Proctor, J. 1987. Nutrient cycling in old secondary rainforests. *Applied Geography* **7**: 135-152.
- Pugnaire, F.I. 2001. Variability of inorganic nutrient concentrations in leaves. *New Phytologist* **150**: 499-507.
- Radjagukguk, B. 1997. Peat soils of Indonesia: location, classification and problems for sustainability. In Rieley, J. O. & Page, S. E. (eds.). Biodiversity and Sustainability of Tropical Peatlands. Proceedings of the International Symposium on Biodiversity, Environmental Importance and Sustainability of Tropical Peat and Peatlands. Samara Publishing Limited, UK.
- Rao, I. M., Zeigler, R. S., Vera, R. & Sarkarung, S. 1993. Selection and breeding for acid-soil tolerance in crops. *BioScience* **43**: 455-465.
- Rustad, L.E. & Cronan, C.S. 1989. Cycling of aluminum and nutrients in litterfall of a red spruce (*Picea rubens* Sarg.) stand in Maine. *Canadian Journal of Forest Research* **19**: 18-23.
- Sirirattanakorn, S. 1999. Leaf functional and leaf demographic studies on productivity of important species in tropical dry evergreen forest and deciduous dipterocarp forest at Sakaerat, Changwat Nakhon Ratchasima. Dissertation submitted to the Graduate School, Kasetsart University, Thailand, pp. 242.
- Soltis, P. S., Soltis, D. E. & Chase, M. W. 1999. Angiosperm phylogeny inferred from multiple genes as a tool for comparative biology. *Nature* **402**: 402-404.
- Stevenson, F.J. & Vance, G.F. 1989. Naturally occurring aluminum-organic complexes. *In*: Sposito, G. (ed.) The environmental chemistry of aluminum, pp 117-146. CRC Press, Boca Raton.
- Suzuki, E., Kohyama, T., Simbolon, H., Haraguchi, A., Tsuyuzaki, S. & Nishimura, T. 1998. Vegetation of kerangas and peat swamp forests in Lahei, Central Kalimantan. Environmental Conservation and Land Use Management of Wetland Ecosystem in Southeast Asia (Annual Report and Proceedings for International Workshop in Palangka Raya, Central Kalimantan, Indonesia, 6-9 August 1997). Core University Program between Hokkaido University, Japan and R&D Center of Biology, LIPI, Indonesia), pp. 1-10
- The Angiosperm Phylogeny Group. 1998. An ordinal classification for the families of flowering plants.
-

- Annals of the Missouri Botanical Garden* **85**: 531-553.
- Thompson, K., Parkinson, J. A., Band, S. R. & Spencer, R. E. 1997. A comparative study of leaf nutrient concentrations in a regional herbaceous flora. *New Phytologist* **136**: 676-689.
- Watanabe, T. & Osaki, M. 2002. Mechanisms of adaptation to high aluminum condition in native plant species growing in acid soils: a review. *Communications in Soil Science and Plant Analysis* **33**: 1247-1260.
- Watson, L. & Dallwitz, M. J. 1992 onwards. The families of flowering plants: descriptions, illustrations, identification, and information retrieval. Version: 14th December 2000. <http://biodiversity.uno.edu/delta/>.

Received Sep. 6, 2002

Accepted Mar. 18, 2003

Appendix 1. Concentrations of mineral elements in the leaves of native plants from tropical peat swamps and heath forests of Central Kalimantan, Indonesia.

Phylogenetic group*	Family	Species	Form	N	P	K	Ca	Mg	Na	Fe	Mn	Zn	Cu	Mo	Al	Si
				(g kg ⁻¹)					(mg kg ⁻¹)							
Primary heath forest (Lahai)																
Euastrid II	Aquifoliaceae	<i>Ilex cynosa</i> Bl.	tree	10.4	0.2	5.5	8.0	4.2	869	24	497	67.5	3.1	5.6	433	954
Euastrid I	Oleaceae	<i>Chionanthus ramiflorus</i> Roxb.	tree	17.5	0.7	8.4	5.8	3.6	2362	107	273	84.6	5.8	5.2	148	770
Euastrid I	Oleaceae	<i>Chionanthus ramiflorus</i> Roxb.	tree	15.5	0.5	3.3	5.5	2.9	706	75	342	18.3	4.6	5.5	761	2959
Euastrid I	Rubiaceae	<i>Timonius flavescens</i> (Jack) Baker	tree	17.6	0.6	6.8	9.2	2.1	2521	88	52	25.1	3.1	5.3	415	2401
Euastrid I	Rubiaceae	<i>Timonius flavescens</i> (Jack) Baker	tree	14.3	0.5	4.2	13.9	3.0	875	82	67	18.9	3.2	5.2	108	842
Ericales	Ebenaceae	<i>Diospyros hermaphrodita</i> (Zoll.) Bakh.	tree	9.8	0.6	13.6	16.3	9.8	1004	86	2111	10.9	1.1	5.9	164	325
Ericales	Ebenaceae	<i>Diospyros hermaphrodita</i> (Zoll.) Bakh.	tree	16.9	0.8	18.5	5.6	5.7	1023	64	473	12.0	3.0	5.9	108	1079
Ericales	Ebenaceae	<i>Diospyros hermaphrodita</i> (Zoll.) Bakh.	tree	15.5	0.7	11.3	9.3	10.2	857	73	517	10.5	2.3	4.8	357	2370
Ericales	Ebenaceae	<i>Diospyros confertiflora</i> (Hiern.) Bakh.	tree	19.7	0.7	6.9	5.7	2.8	1239	72	81	11.4	3.5	5.9	202	911
Ericales	Ebenaceae	<i>Diospyros confertiflora</i> (Hiern.) Bakh.	tree	16.3	0.6	5.6	8.4	4.3	724	82	364	17.5	3.6	5.4	440	2724
Ericales	Tetrameristaceae	<i>Tetramerista glabra</i> Miq.	tree	6.6	0.5	9.2	6.7	3.7	905	29	26	8.1	1.5	5.5	343	1122
Eurosid II	Anacardiaceae	<i>Buchanania sessifolia</i> Bl.	tree	20.8	0.9	5.4	4.4	3.2	1655	117	71	14.3	3.3	5.5	263	1565
Eurosid II	Anacardiaceae	<i>Buchanania sessifolia</i> Bl.	tree	20.7	2.3	25.6	8.3	2.7	2163	106	59	25.2	6.3	5.4	142	1340
Eurosid II	Anacardiaceae	<i>Buchanania sessifolia</i> Bl.	tree	16.4	1.3	11.0	6.0	2.1	1112	87	60	3.5	5.6	4.7	562	3467
Eurosid II	Bursaceae	<i>Senecarpus longifolius</i> Bl.	tree	17.8	1.0	5.5	3.2	1.5	1386	70	22	25.6	4.8	5.6	148	2152
Eurosid II	Bursaceae	<i>Canarium hirsutum</i> Willd.	tree	8.5	0.8	3.3	12.5	3.4	1513	84	24	37.7	2.9	5.6	135	1509
Eurosid II	Dipterocarpaceae	<i>Corylobium malacoxylon</i> Pierre	tree	10.0	0.7	4.0	7.0	1.1	1303	51	465	30.0	2.2	4.9	85	2038
Eurosid II	Dipterocarpaceae	<i>Shorea balangeran</i> (Korth.) Burk.	tree	17.8	0.9	3.2	9.6	2.2	820	99	497	23.2	10.1	5.2	106	854
Eurosid II	Dipterocarpaceae	<i>Shorea balangeran</i> (Korth.) Burk.	tree	16.1	0.9	4.7	8.7	1.7	1219	111	583	20.1	4.2	5.9	451	2914
Eurosid II	Dipterocarpaceae	<i>Shorea balangeran</i> (Korth.) Burk.	tree	15.3	0.6	3.0	4.2	1.4	1203	82	134	10.6	1.0	7.2	627	3616
Eurosid II	Dipterocarpaceae	<i>Vatica rassak</i> (Korth.) Bl.	tree	12.5	0.8	4.7	5.0	1.5	1444	70	151	24.4	8.6	5.7	100	1340
Eurosid II	Myrtaceae	<i>Myrtaceae</i> (Korth.) Bl.	tree	17.8	0.6	4.7	13.5	6.5	1122	32	38	6.9	9.7	4.1	440	2862
Eurosid II	Myrtaceae	<i>Myrtaceae</i> (Korth.) Bl.	tree	13.2	0.7	4.6	6.1	3.4	858	50	58	10.1	2.3	5.4	343	817
Eurosid II	Myrtaceae	<i>Myrtaceae</i> (Korth.) Bl.	tree	9.4	0.5	2.2	3.3	2.6	1083	32	20	5.7	1.8	5.6	136	1126
Eurosid II	Myrtaceae	<i>Myrtaceae</i> (Korth.) Bl.	tree	11.3	0.6	13.6	4.2	4.6	1057	25	27	7.1	9.9	7.8	68	1026
Eurosid II	Myrtaceae	<i>Tristania sp.</i>	tree	13.2	0.7	4.7	7.7	2.5	3178	69	469	15.2	3.6	5.0	935	3311
Eurosid II	Rutaceae	<i>Tetrastima cf. obovata</i> Merr.	tree	14.0	0.4	5.1	10.0	1.0	815	21	23	5.3	4.0	5.2	88	744
Eurosid II	Rutaceae	<i>Tetrastima cf. obovata</i> Merr.	tree	17.7	0.1	8.9	5.7	3.0	2052	63	330	34.6	12.1	4.9	83	1504
Eurosid II	Sapindaceae	<i>Nephelium cf. mangrove</i> Hiern.	tree	10.5	0.8	6.0	10.3	3.9	1451	61	940	29.1	3.5	4.8	117	1675
Eurosid II	Tiliaceae	<i>Pentace borneensis</i> Pierre	tree	11.7	0.6	4.5	3.1	1.1	1186	69	61	9.9	1.9	4.8	65	1229
Eurosid I	Caesalpiniaceae	<i>Dialium patens</i> Bak.	tree	14.5	0.9	7.1	9.3	1.7	883	112	179	10.5	7.3	4.2	108	1783
Eurosid I	Caesalpiniaceae	<i>Koompassia malaccensis</i> Maing. Ex. Benth.	tree	18.1	1.0	6.1	12.3	2.1	2782	65	619	15.8	5.0	5.0	358	3122
Eurosid I	Chrysobalanaceae	<i>Parastemon urphyllus</i> (A.D.C.)	tree	13.1	1.0	11.8	11.6	2.0	892	66	25	12.5	5.0	5.1	117	1305
Eurosid I	Clusiaceae	<i>Calophyllum hosi</i> Ridley	tree	8.8	0.5	6.4	2.4	2.2	1597	38	32	8.6	2.2	5.5	409	2639
Eurosid I	Clusiaceae	<i>Garcinia dioica</i> Bl.	tree	15.4	0.7	6.0	15.4	5.3	2118	45	341	36.4	3.2	5.2	561	3185
Eurosid I	Euphorbiaceae	<i>Baccharis mucronata</i> (Miq.) Muell. Arg.	tree	14.0	0.6	7.6	9.6	5.8	1079	57	63	17.9	3.1	5.4	5104	2245
Eurosid I	Euphorbiaceae	<i>Christinus mangrove</i> Hook.f.	tree	16.8	0.7	9.6	6.9	5.0	758	54	100	6.7	13.6	5.0	42	963
Eurosid I	Euphorbiaceae	<i>Antidesma montanum</i> Bl.	tree	24.4	1.4	5.4	4.6	1.5	617	58	30	2.7	2.2	4.3	237	1277
Eurosid I	Ixonanthaceae	<i>Ixonanthus reticulata</i> Jack.	tree	8.8	0.6	3.1	6.8	5.0	693	31	350	10.8	0.9	5.4	442	834
Core Eudicot	Oleaceae	<i>Strombosia ceylanica</i> Gardner	tree	15.1	0.8	7.7	6.2	2.6	1134	83	518	9.1	3.5	5.3	479	2930
Magnoliales	Annonaceae	<i>Goniolobium malayense</i> Hook.f. & Thoms.	tree	12.3	0.7	7.0	6.5	2.1	979	47	241	9.0	2.5	5.2	393	2715
Magnoliales	Lauraceae	<i>Actinodaphne borneensis</i> Meisn.	tree	10.4	0.5	3.6	4.3	1.1	628	46	254	26.8	2.3	5.3	78	799
Magnoliales	Lauraceae	<i>Litsea obovata</i> (Bl.) Boerl.	tree	24.0	0.8	6.5	2.7	1.9	650	34	114	9.7	3.0	5.1	196	927
Magnoliales	Myristicaceae	<i>Gymnacanthus eugenifolia</i> (A.D.C.) Sinclair.	tree	14.1	0.9	10.3	16.9	7.6	1435	69	433	39.4	1.0	6.0	89	1086
Magnoliales	Myristicaceae	<i>Koema glauca</i> (Bl.) Warb.	tree	12.1	0.6	4.5	8.2	6.1	1206	39	88	36.5	2.5	6.1	460	2794
Primary heath forest (Lahai)																
Euastrid I	Rubiaceae	<i>Gardenia pterocalyx</i> Val.	tree	9.7	0.3	9.4	3.4	2.7	717	61	38	13.9	6.7	6.0	38	57
Euastrid I	Rubiaceae	<i>Isora hirsutifolia</i> Merr.	tree	9.7	0.4	8.7	17.7	4.0	382	71	57	18.7	4.9	6.1	45	38
Euastrid I	Rubiaceae	<i>Timonius flavescens</i> (Jack) Baker	tree	10.5	0.3	2.7	13.2	2.6	610	48	102	9.9	1.5	7.3	44	148
Ericales	Ebenaceae	<i>Diospyros hermaphrodita</i> (Zoll.) Bakh.	tree	10.2	0.4	3.8	14.8	2.4	368	34	1474	7.8	5.0	4.8	31	27
Ericales	Myrsinaceae	<i>Ardisia laevigata</i> Blume	tree	10.2	0.4	2.8	10.2	1.5	501	35	178	3.3	3.0	5.4	33	50
Ericales	Sapotaceae	<i>Isomandra lanceolata</i> Wight.	tree	19.7	1.5	9.5	10.3	8.1	992	95	193	50.9	7.6	4.7	16	135
Ericales	Sapotaceae	<i>Palauquium leiocarpum</i> Boerl.	tree	7.7	0.5	4.2	12.2	1.6	296	18	45	2.1	2.1	5.7	114	16
Ericales	Sapotaceae	<i>Poyena enderbyi</i> H.J. Lam	tree	8.4	1.1	6.8	4.5	3.5	445	47	34	3.6	2.6	6.8	64	52
Ericales	Theaceae	<i>Ternstroemia coriacea</i> R. Scheffer.	tree	13.9	1.1	7.6	7.4	2.7	652	77	73	5.3	6.4	5.2	29	42
Eurosid II	Anacardiaceae	<i>Canarium hirsutum</i> Willd.	tree	15.6	1.0	5.7	6.3	1.7	632	52	60	10.4	6.8	6.5	20	57
Eurosid II	Bursaceae	<i>Canarium hirsutum</i> Willd.	tree	16.2	0.7	5.4	3.9	1.3	307	38	217	4.2	3.1	7.1	24	196
Eurosid II	Bursaceae	<i>Canarium sp.</i>	tree	10.0	0.5	3.0	19.7	1.2	392	36	1513	6.6	2.1	6.6	8	74
Eurosid II	Dipterocarpaceae	<i>Shorea platycarpa</i> Heim.	tree	13.4	0.7	5.7	4.3	1.4	355	49	823	3.4	5.8	5.8	6	22
Eurosid II	Dipterocarpaceae	<i>Shorea ovalis</i> (Korth.) Bl.	tree	17.2	0.8	5.1	4.0	1.1	670	107	242	8.1	5.7	4.9	63	41
Eurosid II	Dipterocarpaceae	<i>Shorea ovalis</i> (Korth.) Bl.	tree	14.5	0.7	4.9	13.2	1.2	937	75	1186	12.4	4.7	5.8	36	46
Eurosid II	Dipterocarpaceae	<i>Shorea toymanniana</i> Dyer ex Brandis	tree	10.5	0.9	5.5	4.7	1.0	649	74	238	7.0	2.4	4.4	28	35
Eurosid II	Dipterocarpaceae	<i>Shorea toymanniana</i> Dyer ex Brandis	tree	16.4	0.9	2.9	2.8	1.5	565	160	198	10.5	4.5	6.1	53	1174
Eurosid II	Dipterocarpaceae	<i>Vatica umbonata</i> (Hook.f.) Burck	tree	22.7	1.1	7.7	2.9	4.5	381	67	35	21.1	4.2	5.0	59	59
Eurosid II	Meliaceae	<i>Aglaia rubiginosa</i> (Hiern.) Pannell	tree	19.9	2.1	7.4	6.2	9.0	546	61	88	15.5	4.8	5.6	15	23
Eurosid II	Meliaceae	<i>Aglaia odoratissima</i> Bl.	tree	14.9	0.8	13.3	6.0	3.6	446	53	455	10.3	5.9	5.8	40	44
Eurosid II	Myrtaceae	<i>Myrtistina luvina</i> King	tree	17.6	0.9	8.1	4.8	4.6	516	73	89	6.3	4.0	5.9	1007	39
Eurosid II	Myrtaceae	<i>Myrtaceae</i> (Korth.) Bl.	tree	11.3	0.4	4.1	4.6	2.9	433	50	40	2.4	8.6	5.8	78	26
Eurosid II	Myrtaceae	<i>Myrtaceae</i> (Korth.) Bl.	tree	9.4	0.3	5.4	5.8	3.0	423	46	51	3.3	12.8	6.8	35	35
Eurosid II	Rutaceae	<i>Tetrastima cf. obovata</i> Merr.	tree	17.2	1.1	5.5	5.4	5.3	734	55	598	18.2	2.9	6.1	82	105
Eurosid II	Sterculiaceae	<i>Heritiera altiliflora</i> (Ridl.) Kosterm.	tree	14.6	0.5	4.2	6.9	4.6	689	91	421	8.1	4.0	7.1	56	101
Eurosid II	Sterculiaceae	<i>Scaphium longifolium</i> Ridl.	tree	18.2	1.1	9.8	2.5	3.4	832	70	143	7.7	5.0	5.9	11	614
Eurosid I	Caesalpiniaceae	<i>Koompassia malaccensis</i> Maing. Ex. Benth.	tree	18.3	1.5	2.5	6.3	1.4	677	65	640	8.3	5.0	6.5	16	759
Eurosid I	Clusiaceae	<i>Calophyllum inophyllum</i>														

Appendix 1. (continued)

Phylogenetic group*		Family	Species	Form	N	P	K	Ca	Mg	Na	Fe	Mn	Zn	Cu	Mo	Al	Si				
					(g kg ⁻¹)				(mg kg ⁻¹)												
Peridophytes	Nephrolepidaceae		<i>Nephrolepis pectinata</i> (Willd.) Schott	fern	15.9	0.8	14.4	6.1	3.5	639	41	282	0.4	0.5	12.4	15	208				
Peridophytes	Nephrolepidaceae		<i>Nephrolepis pectinata</i> (Willd.) Schott	fern	17.6	0.4	13.4	14.2	9.6	579	19	195	7.0	6.2	12.2	19	6				
Peridophytes	Polypodiaceae		<i>Polypodium hirsutum</i> G. Forst	fern	4.1	0.2	12.9	7.0	5.3	413	46	55	5.8	4.7	15.7	170	19				
Peridophytes	Polypodiaceae		<i>Polypodium hirsutum</i> G. Forst	fern	8.1	0.3	18.4	4.9	3.9	164	17	48	26.8	8.6	12.2	142	74				
Peridophytes	Polypodiaceae		<i>Polypodium hirsutum</i> G. Forst	fern	8.1	0.3	16.8	3.4	3.9	368	42	48	4.0	13.4	13.2	65	60				
Peridophytes	Polypodiaceae		<i>Polypodium hirsutum</i> G. Forst	fern	8.5	0.5	18.6	5.5	3.5	740	21	224	5.3	3.3	10.3	9	32				
Peridophytes	Polypodiaceae		<i>Polypodium hirsutum</i> G. Forst	fern	8.6	0.4	19.5	7.1	5.0	501	31	148	11.1	2.0	16.5	80	55				
Peridophytes	Polypodiaceae		<i>Polypodium hirsutum</i> G. Forst	fern	8.7	0.4	19.4	5.1	4.0	543	23	79	11.2	0.9	16.3	60	44				
Peridophytes	Polypodiaceae		<i>Polypodium hirsutum</i> G. Forst	fern	8.7	0.7	16.6	5.1	4.6	567	69	302	15.9	0.7	12.3	68	45				
Peridophytes	Polypodiaceae		<i>Polypodium hirsutum</i> G. Forst	fern	8.8	0.3	16.7	6.9	4.5	203	50	181	4.4	0.7	17.8	87	32				
Peridophytes	Polypodiaceae		<i>Polypodium hirsutum</i> G. Forst	fern	9.5	0.4	15.8	6.3	5.1	223	52	175	16.0	6.2	11.7	57	61				
Peridophytes	Polypodiaceae		<i>Polypodium hirsutum</i> G. Forst	fern	9.7	0.5	18.7	7.3	6.3	634	33	283	12.0	2.4	11.6	3	41				
Peridophytes	Polypodiaceae		<i>Polypodium hirsutum</i> G. Forst	fern	10.0	0.3	17.3	5.7	6.4	379	30	106	19.7	2.8	14.8	66	43				
Peridophytes	Dennstaedtiaceae		<i>Pteridium esculentum</i> L.	fern	6.2	0.3	13.3	0.7	0.9	502	21	22	3.3	7.5	16.4	1	63				
Peridophytes	Dennstaedtiaceae		<i>Pteridium esculentum</i> L.	fern	7.5	0.8	14.2	2.0	1.9	721	20	16	7.5	5.1	18.3	28	31				
Peridophytes	Dennstaedtiaceae		<i>Pteridium esculentum</i> L.	fern	8.3	0.8	11.8	2.9	1.9	385	6	98	9.0	1.8	11.1	5	20				
Peridophytes	Dennstaedtiaceae		<i>Pteridium esculentum</i> L.	fern	8.7	0.5	14.8	7.4	7.5	141	46	198	8.8	8.8	13.9	42	33				
Peridophytes	Dennstaedtiaceae		<i>Pteridium esculentum</i> L.	fern	9.3	0.9	11.9	2.9	1.4	845	6	98	15.0	4.4	8.7	4	10				
Peridophytes	Dennstaedtiaceae		<i>Pteridium esculentum</i> L.	fern	16.0	1.5	12.5	4.3	3.6	70	39	67	34.3	3.7	10.2	17	149				
Primary peat swamp forest - shallow peat (Kalamanggan)																					
Euastrid II	Aquifoliaceae		<i>Ilex cymosa</i> Bl.	tree	6.0	0.1	5.9	4.7	2.8	126	6	14	4.7	5.1	14.9	7	5				
Euastrid I	Apocynaceae		<i>Dyera costulata</i> (Miq.) Hook.f.	tree	13.9	0.3	9.5	2.8	1.7	93	35	186	10.4	4.7	14.8	20	20				
Euastrid I	Apocynaceae		<i>Dyera costulata</i> (Miq.) Hook.f.	tree	16.1	0.5	7.9	9.1	2.1	42	53	209	18.1	2.2	10.1	13	7				
Ericales	Ebenaceae		<i>Diospyros hermaphrodita</i> (Zoll.) Bakh.	tree	13.7	0.4	17.5	8.1	6.0	166	81	1112	3.9	5.4	14.4	5	41				
Ericales	Ebenaceae		<i>Diospyros hermaphrodita</i> (Zoll.) Bakh.	tree	13.7	0.2	15.8	3.9	3.5	78	16	485	4.4	0.3	16.6	91	28				
Ericales	Ebenaceae		<i>Diospyros hermaphrodita</i> (Zoll.) Bakh.	tree	16.1	0.5	10.7	5.6	3.4	92	92	857	3.9	2.8	14.5	5	30				
Ericales	Ebenaceae		<i>Diospyros hermaphrodita</i> (Zoll.) Bakh.	tree	12.7	0.2	10.2	6.8	4.2	151	35	347	3.6	0.4	9.3	5	9				
Ericales	Ebenaceae		<i>Diospyros hermaphrodita</i> (Zoll.) Bakh.	tree	13.4	0.2	16.0	4.3	3.5	34	35	787	4.2	5.8	17.0	3	34				
Ericales	Ebenaceae		<i>Diospyros hermaphrodita</i> (Zoll.) Bakh.	tree	17.6	0.1	9.8	2.5	1.2	309	20	141	15.4	2.8	5.2	50	3				
Ericales	Ebenaceae		<i>Diospyros hermaphrodita</i> (Zoll.) Bakh.	tree	12.7	0.2	18.2	3.9	2.4	251	22	409	11.8	0.2	7.5	17	3				
Ericales	Sapotaceae		<i>Madhuca molleyna</i> (de Vrc.) Pier ex Dubard	tree	6.5	0.4	5.9	6.8	3.1	58	49	19	4.8	3.3	13.7	63	104				
Ericales	Sapotaceae		<i>Madhuca molleyna</i> (de Vrc.) Pier ex Dubard	tree	7.4	0.5	5.4	6.5	3.3	114	47	14	5.3	3.2	13.1	8	107				
Ericales	Tetrameristaceae		<i>Tetramerista glabra</i> Miq.	tree	7.5	0.1	11.1	4.2	2.5	181	10	12	5.3	6.7	13.2	2	3				
Ericales	Tetrameristaceae		<i>Tetramerista glabra</i> Miq.	tree	9.5	0.2	9.1	3.8	0.9	4	14	12	4.8	3.8	12.5	6	33				
Ericales	Tetrameristaceae		<i>Tetramerista glabra</i> Miq.	tree	8.1	0.3	9.6	5.7	2.7	34	67	6	3.9	14.0	14.1	7	74				
Ericales	Tetrameristaceae		<i>Tetramerista glabra</i> Miq.	tree	9.9	0.2	9.9	2.3	0.5	392	11	19	15.7	7.5	5.8	54	7				
Ericales	Tetrameristaceae		<i>Tetramerista glabra</i> Miq.	tree	15.7	0.1	10.0	1.9	0.5	333	10	25	14.4	1.3	3.0	12	8				
Ericales	Tetrameristaceae		<i>Tetramerista glabra</i> Miq.	tree	11.6	0.1	8.4	2.1	1.0	375	12	6	2.7	1.9	9.9	11	3				
Ericales	Tetrameristaceae		<i>Tetramerista glabra</i> Miq.	tree	11.6	0.2	11.5	7.3	2.3	210	11	5	3.1	1.3	6.7	35	4				
Eurosid II	Dipterocarpaceae		<i>Shorea balangeran</i> (Korth.) Burck	tree	13.2	0.3	6.7	7.0	2.1	150	19	15	3.7	1.2	13.9	4	32				
Eurosid II	Dipterocarpaceae		<i>Shorea balangeran</i> (Korth.) Burck	tree	10.8	0.6	5.8	8.4	1.1	84	129	608	4.0	5.8	12.8	6	25				
Eurosid II	Dipterocarpaceae		<i>Shorea platycarpa</i> Heim.	tree	10.5	0.3	2.4	1.8	1.5	6	19	170	13.1	4.0	14.9	7	22				
Eurosid II	Dipterocarpaceae		<i>Shorea platycarpa</i> Heim.	tree	11.5	0.3	5.0	4.8	0.8	317	16	23	10.4	0.2	4.2	21	11				
Eurosid II	Dipterocarpaceae		<i>Shorea retusa</i> Meijer	tree	14.5	0.3	6.3	4.2	1.4	289	25	94	2.8	2.2	15.1	14	18				
Eurosid II	Melastomataceae		<i>Dactyloctenium aegyptium</i> Oliv., in Hook	tree	12.3	0.3	6.3	4.9	2.9	110	15	17	5.0	0.5	15.1	12	8				
Eurosid II	Myrtaceae		<i>Syzygium borneense</i> (Miq.) Muell. Arg.	tree	8.7	0.2	7.9	10.4	2.4	122	56	9	4.3	8.4	15.2	10	22				
Eurosid II	Myrtaceae		<i>Syzygium borneense</i> (Miq.) Muell. Arg.	tree	9.3	0.1	11.1	10.0	1.3	286	17	21	13.9	4.6	4.8	66	14				
Eurosid II	Myrtaceae		<i>Syzygium carnosifolium</i> (Blume) Merr. & L.M. Perr.	tree	12.8	0.2	2.6	5.3	1.1	266	24	441	6.2	6.4	12.8	49	16				
Eurosid II	Myrtaceae		<i>Syzygium lepidocarpum</i> Wall.	tree	16.4	0.2	3.4	7.5	0.8	214	13	18	4.1	8.2	11.0	4	81				
Eurosid II	Myrtaceae		<i>Syzygium lepidocarpum</i> Wall.	tree	10.5	0.2	14.7	6.8	3.9	156	10	30	5.6	11.3	11.4	66	15				
Eurosid II	Myrtaceae		<i>Tristania obscura</i> (Benth.) P.G. Wilson & Waterh.	tree	8.6	0.2	7.1	10.7	1.1	23	15	42	3.1	0.3	13.9	12	18				
Eurosid II	Rutaceae		<i>Acronychia porteri</i> Hook.f.	tree	16.5	0.2	3.8	6.0	1.0	141	23	7	4.4	1.5	11.6	90	124				
Eurosid II	Sterculiaceae		<i>Sterculia gila</i> Miq.	tree	15.2	0.3	8.0	5.9	1.7	524	28	93	12.4	1.0	4.3	29	2				
Eurosid II	Thymelaeaceae		<i>Gonyolobos bancanus</i> Kurz.	tree	10.0	0.1	8.0	2.1	0.6	338	20	23	5.8	3.8	5.7	21	10				
Eurosid II	Thymelaeaceae		<i>Gonyolobos bancanus</i> Kurz.	tree	9.3	0.3	2.6	5.6	1.5	416	100	257	0.4	1.1	11.9	5	4				
Eurosid I	Anisophylleaceae		<i>Combretocarpus rotundatus</i> (Miq.) Danser	tree	10.2	0.5	8.5	3.5	1.9	161	40	432	12.3	4.5	14.6	8294	28				
Eurosid I	Anisophylleaceae		<i>Combretocarpus rotundatus</i> (Miq.) Danser	tree	10.7	0.3	6.5	4.0	2.5	169	50	323	3.8	0.9	14.7	8578	13				
Eurosid I	Caesalpiniaceae		<i>Koompassia malaccensis</i> Maing. Ex Benth.	tree	11.0	0.3	2.8	6.6	2.6	72	19	3	5.2	3.7	13.8	2	4				
Eurosid I	Caesalpiniaceae		<i>Koompassia malaccensis</i> Maing. Ex Benth.	tree	14.2	0.6	3.2	5.9	2.9	42	37	15	4.9	7.0	15.4	4	4				
Eurosid I	Caesalpiniaceae		<i>Koompassia malaccensis</i> Maing. Ex Benth.	tree	17.0	0.4	9.0	3.9	0.9	62	44	28	3.7	4.2	12.1	8	41				
Eurosid I	Caesalpiniaceae		<i>Koompassia malaccensis</i> Maing. Ex Benth.	tree	13.1	0.2	3.0	6.8	2.9	366	11	132	16.6	2.0	5.0	30	35				
Eurosid I	Celastraceae		<i>Lophopetalum multistriatum</i> Ridl.	tree	12.3	0.1	5.9	5.8	1.4	435	17	26	7.3	0.6	5.3	6	20				
Eurosid I	Celastraceae		<i>Calophyllum boeri</i> Ridl.	tree	6.6	0.2	10.7	2.6	1.7	29	6	2	4.4	0.3	12.4	7	45				
Eurosid I	Celastraceae		<i>Calophyllum boeri</i> Ridl.	tree	9.5	0.2	6.2	10.6	1.9	122	28	28	4.1	3.8	11.7	3	13				
Eurosid I	Celastraceae		<i>Calophyllum inophyllum</i> Linn.	tree	16.3	0.2	7.5	7.1	4.0	72	39	361	10.8	1.4	14.2	7	57				
Eurosid I	Celastraceae		<i>Calophyllum inophyllum</i> Linn.	tree	9.3	0.3	8.6	6.2	6.2	36	48	17	4.5	10.4	13.5	8	24				
Eurosid I	Celastraceae		<i>Calophyllum inophyllum</i> Linn.	tree	16.7	0.6	10.8	4.4	5.2	1	95	431	4.3	12.5	15.9	7	122				
Eurosid I	Celastraceae		<i>Calophyllum inophyllum</i> Linn.	tree	7.9	0.1	5.4	3.0	0.7	296	15	24	9.4	4.4	4.8	3	5				
Eurosid I	Celastraceae		<i>Calophyllum inophyllum</i> Linn.	tree	12.4	0.2	5.9	4.3	0.7	322	15	35	8.9	4.0	5.3	6	7				
Eurosid I	Celastraceae		<i>Calophyllum macrocarpum</i> Hook.f.	tree	12.1	0.2	1.8	5.0	1.2	189	37	284	3.3	6.2	13.6	4	40				
Eurosid I	Celastraceae		<i>Calophyllum macrocarpum</i> Hook.f.	tree	6.3	0.2	4.0	4.3	2.6	151	26	12	4.3	2.9	12.3	6	29				
Eurosid I	Celastraceae																				

Appendix 1. (continued)

Phylogenetic group*	Family	Species	Form	N	P	K	Ca	Mg	Na	Fe	Mn	Zn	Cu	Mo	Al	Si
				(g kg ⁻¹)					(mg kg ⁻¹)							
Pteridophytes	Polypodiaceae	<i>Polypodium hirsutum</i> G. Forst	fern	11.3	0.6	12.1	5.5	4.5	379	53	161	8.0	3.3	12.4	37	24
Pteridophytes	Polypodiaceae	<i>Polypodium hirsutum</i> G. Forst	fern	12.0	0.3	12.3	4.0	2.2	578	14	51	3.1	2.3	10.7	36	3
Pteridophytes	Polypodiaceae	<i>Polypodium hirsutum</i> G. Forst	fern	12.6	0.3	11.8	5.3	3.2	411	24	59	2.8	1.4	10.6	96	8
Pteridophytes	Polypodiaceae	<i>Polypodium hirsutum</i> G. Forst	fern	12.9	0.8	13.9	2.1	3.1	395	34	38	2.1	5.7	9.2	27	5
Pteridophytes	Polypodiaceae	<i>Polypodium hirsutum</i> G. Forst	fern	13.1	0.8	12.4	1.9	1.9	378	131	41	5.5	8.9	10.8	175	78
Pteridophytes	Polypodiaceae	<i>Polypodium hirsutum</i> G. Forst	fern	13.9	0.7	11.0	1.1	1.3	294	49	14	7.1	4.1	8.7	194	19
Pteridophytes	Dennstaedtiaceae	<i>Peridium esculentum</i> L.	fern	11.4	0.2	8.0	3.7	0.8	522	12	108	6.7	2.1	10.6	37	10
Pteridophytes	Dennstaedtiaceae	<i>Peridium esculentum</i> L.	fern	12.1	0.9	12.0	2.2	2.6	255	31	84	11.4	4.2	9.0	78	9
Pteridophytes	Dennstaedtiaceae	<i>Peridium esculentum</i> L.	fern	12.5	0.5	14.6	2.2	1.5	458	24	6	3.0	0.5	10.8	49	7
Pteridophytes	Dennstaedtiaceae	<i>Peridium esculentum</i> L.	fern	12.5	0.5	11.8	1.8	1.6	502	27	8	2.2	0.6	12.5	52	13
Pteridophytes	Dennstaedtiaceae	<i>Peridium esculentum</i> L.	fern	13.1	0.5	11.3	1.9	1.3	428	24	25	5.1	2.1	11.6	127	8
Pteridophytes	Dennstaedtiaceae	<i>Peridium esculentum</i> L.	fern	13.3	0.2	11.5	5.1	4.6	294	69	112	3.4	2.0	8.4	69	66
Pteridophytes	Dennstaedtiaceae	<i>Peridium esculentum</i> L.	fern	13.3	0.7	14.2	2.6	2.5	396	30	43	4.0	5.2	10.2	20	4
Pteridophytes	Dennstaedtiaceae	<i>Peridium esculentum</i> L.	fern	13.9	0.6	12.1	2.3	1.3	407	17	2	4.7	0.3	10.6	88	17
Pteridophytes	Dennstaedtiaceae	<i>Peridium esculentum</i> L.	fern	20.0	2.5	19.7	2.6	3.4	135	236	71	18.2	3.2	11.3	24	63
Primary peat swamp forest-riverine (Setia Alam)																
Euastrid I	Apocynaceae	<i>Dyera costulata</i> (Miq.) Hook.f.	tree	18.7	0.5	7.9	4.8	1.8	542	27	333	3.5	1.5	13.3	13	6
Ericales	Ebenaceae	<i>Diospyros hermaphrodica</i> (Zoll.) Bakh.	tree	9.2	1.1	5.7	2.9	1.4	1092	13	71	1.0	0.8	13.0	23	18
Ericales	Sapotaceae	<i>Palaquium cochlerifolium</i> P. van Royen	tree	10.1	0.3	3.9	6.0	2.2	1027	29	30	1.1	0.2	11.7	52	4
Ericales	Sapotaceae	<i>Palaquium ridleyi</i> King & Gamble	tree	10.3	0.6	3.7	8.9	5.8	1246	23	1316	2.6	1.5	12.6	27	1
Ericales	Tetrameristaceae	<i>Tetramerista glabra</i> Miq.	tree	9.5	0.3	13.9	2.6	0.9	699	23	25	1.4	3.6	12.7	10	4
Eurosoid II	Anacardiaceae	<i>Mangifera quadrifida</i> Jack.	tree	12.1	1.4	5.7	9.6	1.7	3733	23	127	0.4	1.7	13.7	13	6
Eurosoid II	Bombaceae	<i>Durio beccarianus</i> L.	tree	9.5	0.2	14.3	4.4	4.4	952	35	43	0.6	1.3	11.5	35	5
Eurosoid II	Dipterocarpaceae	<i>Shorea teysmanniana</i> Dyer ex Brandis	tree	12.4	0.5	4.0	5.8	3.3	545	26	47	1.1	2.7	13.2	35	42
Eurosoid II	Dipterocarpaceae	<i>Shorea uliginosa</i> Forst.	tree	15.1	0.6	9.2	4.9	0.8	3328	26	289	0.3	0.7	12.8	27	4
Eurosoid II	Dipterocarpaceae	<i>Dactyloctenium aegyptium</i> Oliv., in Hook	tree	12.8	0.4	3.2	4.1	1.2	1118	35	184	1.3	5.1	14.6	10	2
Eurosoid II	Myrtaceae	<i>Syzygium rugosum</i> Korth.	tree	14.5	0.5	6.3	5.9	2.1	247	38	1414	0.9	3.9	11.7	8	120
Eurosoid II	Myrtaceae	<i>Syzygium rugosum</i> Korth.	tree	10.0	0.7	3.5	6.2	2.3	1546	21	29	0.8	1.2	12.7	14	10
Eurosoid II	Myrtaceae	<i>Syzygium rugosum</i> Korth.	tree	6.5	0.5	8.5	3.5	2.2	863	14	26	0.5	0.7	12.1	6	4
Eurosoid II	Myrtaceae	<i>Syzygium rugosum</i> Korth.	tree	13.3	0.4	13.0	6.7	2.5	1007	25	703	1.1	1.5	12.7	53	5
Eurosoid II	Myrtaceae	<i>Tristania ovata</i> (Penn.) P. G. Wilson & Waterh.	tree	8.1	0.2	6.0	14.1	2.1	357	24	80	1.7	2.5	15.0	18	4
Eurosoid II	Sapindaceae	<i>Nephelium mangayi</i> Hiern.	tree	16.1	0.6	9.7	3.7	1.5	770	36	474	1.5	3.0	11.9	36	69
Eurosoid II	Thymelaeaceae	<i>Gonystylis bancanus</i> Kurz.	tree	6.9	0.1	2.5	2.8	0.5	2407	11	186	1.7	0.0	13.2	24	4
Eurosoid II	Thymelaeaceae	<i>Gonystylis bancanus</i> Kurz.	tree	8.4	0.2	2.5	3.1	0.6	1248	32	227	1.0	0.8	13.0	23	57
Eurosoid II	Thymelaeaceae	<i>Gonystylis bancanus</i> Kurz.	tree	11.1	1.2	6.8	4.5	1.5	981	27	303	4.9	3.5	13.5	2090	38
Eurosoid I	Anisophylleaceae	<i>Cambesia rotundata</i> (Miq.) Danser	tree	10.6	0.8	7.1	4.6	1.6	557	25	337	4.2	1.7	13.3	2005	7
Eurosoid I	Caesalpiniaceae	<i>Koompassia malaccensis</i> Maing. Ex Benth.	tree	11.5	0.4	2.7	6.4	3.0	303	23	216	8.5	0.3	12.8	26	139
Eurosoid I	Chrysobalanaceae	<i>Parastemon urophyllum</i> (A.D.C.)	tree	10.1	0.3	6.7	7.3	2.8	124	50	352	71.9	3.1	14.5	26	5
Eurosoid I	Clusiaceae	<i>Calophyllum inophyllum</i> Linn.	tree	8.0	0.2	3.0	3.6	0.8	1056	18	32	1.1	0.6	14.7	53	33
Eurosoid I	Clusiaceae	<i>Garcinia dioica</i> Bl.	tree	11.2	0.3	10.0	7.4	4.5	80	48	366	5.3	3.5	12.3	86	8
Eurosoid I	Clusiaceae	<i>Garcinia dioica</i> Bl.	tree	13.8	0.5	6.6	7.8	2.8	889	31	76	0.7	1.6	13.1	28	115
Eurosoid I	Euphorbiaceae	<i>Macaranga pruriens</i> (Miq.) Muell. Arg.	tree	10.9	0.4	5.1	4.8	1.2	981	33	311	2.8	6.2	13.1	79	20
Eurosoid I	Hypericaceae	<i>Cratogeomys arboreus</i> Bl.	tree	6.2	0.3	8.4	5.2	3.0	979	23	39	1.0	0.8	11.7	18	5
Laurales	Lauraceae	<i>Alseodaphne coriacea</i> Kosterm.	tree	12.3	0.4	4.9	5.9	3.3	974	74	62	1.0	0.5	13.4	68	99
Laurales	Lauraceae	<i>Dehaasia caesia</i> Blume	tree	12.3	0.4	4.9	5.9	3.3	974	74	62	1.0	0.5	13.4	68	99
Primary peat swamp forest-dense pole (Setia Alam)																
Euastrid II	Aquifoliaceae	<i>Ilex cymosa</i> Bl.	tree	13.9	0.3	14.8	9.8	4.3	111	31	1128	0.6	4.9	14.7	14	5
Euastrid I	Apocynaceae	<i>Dyera costulata</i> (Miq.) Hook.f.	tree	16.0	0.4	8.4	7.8	1.4	733	35	435	0.4	0.9	11.1	12	5
Euastrid I	Rubiaceae	<i>Timonius flavescens</i> (Jack) Backer	tree	8.5	0.2	5.1	15.5	1.5	396	21	95	1.1	0.8	14.8	12	17
Ericales	Ebenaceae	<i>Diospyros hermaphrodica</i> (Zoll.) Bakh.	tree	9.8	0.5	3.8	5.6	3.0	541	36	99	0.3	2.0	9.7	6	58
Ericales	Sapotaceae	<i>Madhuca molleana</i> (de Vries) Pier ex Dubard	tree	11.1	0.4	6.9	11.5	3.0	501	46	522	0.4	3.9	11.2	30	5
Ericales	Sapotaceae	<i>Palaquium ridleyi</i> King & Gamble	tree	8.5	0.4	9.2	19.6	5.1	50	37	44	4.5	4.5	9.6	38	4
Ericales	Tetrameristaceae	<i>Tetramerista glabra</i> Miq.	tree	9.7	0.3	9.6	4.8	1.1	792	15	38	0.4	3.0	10.8	17	5
Eurosoid II	Anacardiaceae	<i>Canosperma coriacum</i> (Jack.) Hall.f.	tree	12.5	0.5	7.3	17.3	2.8	239	43	132	1.5	3.3	15.3	79	15
Eurosoid II	Caesalpiniaceae	<i>Koompassia malaccensis</i> Maing. Ex Benth.	tree	10.6	0.3	8.3	8.5	2.5	407	27	164	2.9	1.4	14.6	22	36
Eurosoid II	Dipterocarpaceae	<i>Shorea teysmanniana</i> Dyer ex Brandis	tree	10.9	0.3	8.0	5.4	1.3	303	35	52	0.9	2.9	15.3	19	13
Eurosoid II	Myrtaceae	<i>Syzygium rugosum</i> Korth.	tree	8.9	0.3	9.1	10.0	2.1	234	15	31	1.0	3.0	10.9	26	4
Eurosoid II	Myrtaceae	<i>Syzygium rugosum</i> Korth.	tree	9.2	0.2	7.0	28.7	2.2	570	23	27	0.1	1.1	14.4	280	4
Eurosoid II	Myrtaceae	<i>Syzygium rugosum</i> Korth.	tree	7.4	0.2	6.0	13.6	3.9	31	16	30	1.1	1.0	13.0	30	15
Eurosoid II	Myrtaceae	<i>Syzygium rugosum</i> Korth.	tree	9.0	0.3	6.1	5.5	4.2	175	22	29	0.9	2.6	13.4	30	23
Eurosoid II	Sapindaceae	<i>Nephelium mangayi</i> Hiern.	tree	10.0	0.8	6.0	6.5	1.8	555	35	24	0.4	4.3	11.2	10	6
Eurosoid II	Thymelaeaceae	<i>Gonystylis bancanus</i> Kurz.	tree	9.2	0.2	4.8	5.2	1.3	709	18	330	0.4	2.7	11.0	19	5
Eurosoid I	Anisophylleaceae	<i>Cambesia rotundata</i> (Miq.) Danser	tree	12.4	0.4	3.9	5.6	1.6	28	32	185	1.7	2.1	13.5	691	6
Eurosoid I	Clusiaceae	<i>Calophyllum inophyllum</i> Linn.	tree	10.5	0.2	5.1	3.1	1.0	401	29	28	1.0	3.3	15.9	17	19
Eurosoid I	Clusiaceae	<i>Calophyllum inophyllum</i> Linn.	tree	7.9	0.2	4.3	4.5	2.0	58	21	132	0.3	2.0	13.1	10	3
Eurosoid I	Clusiaceae	<i>Garcinia dioica</i> Bl.	tree	10.7	0.4	8.9	24.3	6.7	566	38	770	15.0	2.9	9.4	27	4
Eurosoid I	Hypericaceae	<i>Cratogeomys arboreus</i> Bl.	tree	9.7	0.3	2.7	5.4	1.1	247	23	223	1.3	4.8	15.7	8	32
Eurosoid I	Moraceae	<i>Ficus microcarpa</i> L.	tree	11.5	0.6	3.8	9.2	2.0	111	38	44	0.4	1.7	11.2	14	4
Magnoliales	Annonaceae	<i>Xylocarpus fuscus</i> Maing.	tree	12.9	0.5	10.6	12.8	6.6	533	41	388	0.3	2.9	10.3	16	5
Magnoliales	Myristicaceae	<i>Horsfieldia subglobosa</i> (Miq.) Warb.	tree	6.8	0.4	2.4	17.5	11.2	433	42	1064	6.5	2.0	11.1	2	76
Laurales	Lauraceae	<i>Alseodaphne coriacea</i> Kosterm.	tree	8.5	0.2	5.6	13.6	2.6	799	17	38	0.5	1.6	10.3	32	13
Laurales	Lauraceae	<i>Cinnamomum sintoc</i> Bl.	tree	9.2	0.4	6.2	8.0	2.3	558	28	400	0.4	1.2	11.3	26	68
Laurales	Lauraceae	<i>Dehaasia caesia</i> Blume	tree	12.6	0.4	8.4	9.2	3.5	852	34	78	0.4	0.5	8.3	18	181

* Classification used following the Angiosperm Phylogeny Group (APG, 1998) and Soltis *et al.* (1999), and Pteridophytes group was also added.