The Chinese Species and the World Genera of Vermileonidae (Diptera)

Akira NAGATOMI, Chi-kung YANG & Ding YANG

CONTENTS

Family Diagnosis	2
List of the Genera and Species	4
1. Genus <i>Lampromyia</i> Macquart	4
2. Genus <i>Leptynoma</i> Westwood	12
3. Genus <i>Vermileo</i> Macquart	16
4. Genus <i>Vermilynx</i> Stuckenberg	25
5. Genus <i>Vermiophis</i> C. Yang	26
6. Genus <i>Vermipardus</i> Stuckenberg	51
7. Genus <i>Vermitigris</i> Wheeler	54
Fossil Genus and Possibly Related Fossil Family	69
Geographical Distribution	69
Citations from the Literature on Immature Stages	75
Citations from the Literature on General Biology	92
Difference in Behavior between Vermileonids and Pit-making Myrmeleontids	123
Biological Observations on <i>Vermiophis ganquanensis</i> ..	124
Biology of some <i>Vermiophis</i> Species	128
Discussion	129
Index	153

Vermiophis wudangensis

TROPICS

熱帯研究

Published by
the Japan Society
of Tropical Ecology

The TROPICS is issued quarterly with one volume a year. It is open either to the results of original research, experimental or descriptive, or significant reviews of various aspects of tropical regions. The journal publishes papers in all aspects of ecology and/or other fields of science related to ecology. It aims not only to act as a valued means of communication between Japanese ecologists and other field scientists but also to stimulate the research activities of young ecologists/scientists of tropical nations especially in Southeast Asia. All the members of the society can contribute to TROPICS, and invited articles may be accepted at the request of editorial board. Special issues may be published on occasion by permission of editorial board, as monographs or reviews exceed approximately 100 pages in length.

Editor-in-Chief: Mitsuru HOTTA (Department of Earth and Environmental Sciences, Faculty of Science, Kagoshima University)

Editorial Board

Takuya ABE (Kyoto University)	Yusho ARUGA (Tokyo University of Fisheries)
Akio FURUKAWA (National Institute for Environmental Studies)	Toshitaka IWAMOTO (Miyazaki University)
Takashi KOHYAMA (Hokkaido University)	Naoki KOYAMA (Kyoto University)
Narifumi MAEDA (Kyoto University)	Misa MASUDA (Tsukuba University)
David W. ROUBIK (Smithsonian Tropical Research Institute)	Katsutoshi SAKURAI (Kochi University)
Yo-Ichiro SATO (Shizuoka University)	Eizi SUZUKI* (Kagoshima University)
Isamu YAMADA (Kyoto University)	Takuo YAMAKURA (Osaka City University)
Seiki YAMANE (Kagoshima University)	Tsuyoshi YONEDA* (Osaka Kyoiku University)
	* secretaries of editorial board

Instructions to authors Two copies of manuscript, preferably in English (or Japanese), to be sent to the editors, typewritten on one side of paper, double-spaced, and with a wide margin. Underlinings to indicate letter types should be left to the editors. In addition to type-script copies, authors using word processing equipments are requested to submit a diskette or floppy disk with their text file. Preference is files in Microsoft Word for Macintosh, on double-sided 3.5 inch (DD, HD) diskettes. For texts processed on MS-DOS computers (preferably in MS-Word or Word Perfect), please submit double-sided (DD or HD) 3.5 inch floppy disks with the text saved as ASCII file.

The editors may ask the opinion of one or two reviewers before accepting or refusing the manuscript that has been submitted.

Summary. Each paper should be provided with a summary.

Citation of literature. In the text as follows: (Primack, 1991) or Ashton (1991). In the 'References' as follows:

Primack, R. B. & Hall P. 1991. Species diversity research in Bornean forests with implications for conservation biology and silviculture. *Tropics* 1: 91-111.

Jacobs, M. 1974. Botanical panorama of the Malesian archipelago. In UNESCO, *Natural Resources of Humid Tropical Asia*, 263-294. Paris.

Illustrations should be professionally drawn and lettered, or laser printed on smooth white paper. Photographs and color slides are also acceptable, but the author(s) will be charged for printing them in color.

Reprints. For articles with one author 50 copies free.

Manuscripts and correspondence concerning editorial matters should be addressed to **Editorial Office of the Japan Society of Tropical Ecology**, c/o Department of Earth and Environmental Sciences, Faculty of Science, Kagoshima University, Korimoto 1-21-35, Kagoshima 890-0065, Japan. Phone: (099) 285-8166 Fax: (099) 285-8172 E-mail: hotta@sci.kagoshima-u.ac.jp

General Office of the Japan Society of Tropical Ecology addressed to c/o Dr. T. Yamakura, Faculty of Science, Osaka City University, Sumiyashi-ku, Osaka 558-0022, Japan. Phone: (06) 605-2582 Fax: 605-2522

The Chinese Species and the World Genera of Vermileonidae (Diptera)

Akira NAGATOMI

4-30-7, Murasakibaru, Kagoshima 890-0082, Japan

Chi-kung YANG & Ding YANG

Department of Entomology, China Agricultural University, Beijing
100094, P. R. China

ABSTRACT This paper revises the 2 Chinese genera and 8 species, lists the 7 world genera and 55 species, gives many citations from the literature on immature stages and general biology, and discusses several characters from the taxonomic and phylogenetic points of view as well as geographical distribution, a fossil genus and possibly a related fossil family concerning the Vermileonidae.

Key Words: Vermileonidae / world genera / Chinese species / phylogeny / biology

One of the aims of this paper is to introduce the Chinese species of Vermileonidae described by C. Yang (1979a, b, 1988) and C. Yang & Chen (1986, 1987, 1993) in Chinese. A comparative study has been made here between *Vermiophis* C. Yang and other known genera of the world. Besides the Chinese ones, we have seen directly the following genera and species: *Leptynoma namaquaensis* Stuckenberg, 1960 (from South Africa); *Vermileo comstocki* Wheeler, 1918 (California); *Vermileo opacus* (Coquillett, 1904) (California to New Mexico and Colorado); *Vermipardus intermedius* Stuckenberg, 1955 (South Africa). Our knowledge of the genera *Lampromyia*, *Leptynoma*, *Vermilyn*x and *Vermipardus* given in this paper is entirely due to Stuckenberg (1960, 1995a, b, 1996a, b, 1997, 1998). We hope that the relationships among the genera will be clarified in greater detail in this paper through the presentation of *Vermiophis* which is the most generalized genus.

A list of the genera and species has been prepared partly after Kertész (1908), Verrall (1909), James (1965, 1968), Nagatomi (1975, 1982), Stuckenberg (1980) and Majer (1988). Some important changes have been made to Nagatomi (1982) and Majer (1988) by Stuckenberg (1960, 1965b, 1980).

This paper also gives many citations from the literature on immature stages and general biology. It discusses several characters from the taxonomic and phylogenetic points of view, a fossil genus (= *Protovermileo* Hennig) and geographical distribution. Thus it replaces Nagatomi (1997) whose content was short.

Some head structures and female terminalia of the Chinese species will be studied by some of us in the near future.

Except for *Lampromyia* s. lat. studied by Stuckenberg, the fauna of Vermileonidae is poorly

known. *Vermileo* may possibly contain more than one genus. The knowledge of the immature stages is scanty. Few studies have been made from the aspects of modern ecology and ethology. Everything remains an open field for study.

It should be noted that the quotation marks are omitted in several places where the extent of the cited sentence is apparent.

FAMILY DIAGNOSIS

General appearance resembling *Dialysis* (Coenomyiidae), *Systropus* (Bombyliidae) and *Leptogaster* (Asilidae). Abdomen slender and hind leg long. Hairs on body and legs short and inconspicuous, except for those on cheek, occiput and in *Vermitigris* side of abdominal tergum 1. Pleuron bare, except for propleuron and anterior part of metapleuron.

Head: Wider than high (in anterior view) and higher than long (in lateral view) (except for, e. g. *Vermipardus intermedius* where head is roughly as high as long, because face is protruding conspicuously). Occiput swollen in *Vermileo comstocki* and *Vermileo opacus*, swollen above neck in *Vermiophis* spp. and *Vermitigris sinensis*, rather swollen (in *Vermipardus intermedius*) or nearly flat (in *Lampromyia namaquaensis*, *Vermileo vermileo*, *Vermileo dowi*, etc.). Eyes separated and with facets of uniform size in both sexes. Frons and face narrower than one eye (in anterior view) and the latter longer than distance from antenna to median ocellus. Antennal scape longer than wide; pedicel roughly as wide or sometimes wider than long; flagellum 3-8 segmented, varies in shape according to genus, with or without a slender arista; flagellum gradually tapering apically in general. Proboscis short and fleshy or long and sclerotized according to genus. Palpal segment 2 widest around middle and then gradually tapering apically in *Vermiophis* and *Vermitigris*, swollen in *Vermileo comstocki* and *Vermileo opacus* or cylindrical and more or less swollen beyond middle in *Lampromyia*, *Leptynoma*, *Vermilyn* and *Vermipardus*.

Thorax: Prosternum widely separated from propleuron. Scutellum flattened dorsally, plank-like and with posterior margin rounded. Postscutellum well developed. Subscutellum absent.

Wing: Distance from humeral crossvein to wing base [= (1)] long, and that from humeral crossvein to base of vein R_s longer than (1) (except for *Leptynoma namaquaensis*). Apical portion of vein R_{2+3} gently or strongly curved, and wing margin between R_1 and R_{2+3} longer than that between R_{2+3} and R_4 (except for *Lampromyia namaquaensis*). Veins R_4 and R_5 ending at wing margin above and below wing apex respectively. Crossvein r-m situated before middle of discal cell. Vein M_4 arising from 2nd basal cell or sometimes at the juncture between discal cell and 2nd basal cell. Anal lobe narrow and long. Alula is absent, but *Vermitigris* has a narrow alula at least in ♀ *sinensis*. The calypters not developed. The narrowest and more or less paler point of halter may be the boundary between stem and knob, and if so the knob would not be shorter than stem.

Legs: Hind leg longer than fore or mid leg, and longer than thorax + abdomen. Apical portion of hind femur is swollen or rather so and it is often so in that of hind tibia. Tarsomere 1 in each leg longer than tarsomeres 2-5. Tibial spur formula 1: 2: 2 (but in *Leptynoma sericea* and *Leptynoma appendiculata*, it is 0: 2: 2). Apex of ventral (= anterior) side of hind coxa with an elongate process in *Vermileo* (at least in *V. comstocki* and *V. opacus*) but absent in other genera.

Abdomen: Slender and narrower than thorax; narrowest in segments 2, 3 or 2-3 and widest in

segments 4-5, 5-6, or 6-7 according to individual.

For face, abnormal wing crossveins, pulvilli and empodium, male and female genitalia, sexual dimorphism, etc., see DISCUSSION section.

The Vermileonidae is easily distinguished from the Rhagionidae, Pelecorhynchidae, Athericidae and Tabanidae by having the hind leg markedly long; scutellum flattened dorsally and plank-like; etc. It is also separated from the Xylophagoidea (= Rachiceridae, Xylophagidae, Coenomyiidae, Heterostomidae and Exeretonevridae) by having the clypeus which is not concave and dish-like but flat, or with lower portion protruding forward.

Key 1: Genera Based on External Characters

(couplets 2-5 are after Stuckenberg, 1995b and 1996b)

1. Proboscis long and sclerotized, and much longer than height of head 2
 - Proboscis short and fleshy, and much shorter than height of head 6
- 2(1). Antennal flagellum 8-segmented and with last joints much shorter than preceding 3 joints together; proboscis rigid, directed downward, relatively short and distinctly less than twice as long as fore femur 3
 - Antennal flagellum 3-8 segmented and with last joint long, thin, arista-like and longer than rest of flagellum or that minus basal joint; proboscis "capable of posterior flexure between the legs when the fly is at rest" (1996b: 249), "very elongate, clearly more than twice as long as fore femure" (1995b: 219) 4
- 3(2). "Labella elongate tubular, dorsoventrally flattened in dried specimens, without setae" (1995b: 219); proboscis 1.5-1.7 times as long as fore femur *Vermilyn*
 - "Labella laterally compressed, short, subtriangular, bearing stiff setae" (1995b: 219); proboscis 0.5-1.4 times as long as fore femur *Vermipardus*
- 4(2). "Hind tibia not markedly swollen apically, clearly more slender than hind femur" (1996b: 249); wing shape unspecialised (1996b: 249 & figs. 29-31) as quoted under generic diagnosis of this paper *Lampromyia*
 - "Hind legs appearing robust because tibia is swollen apically as much as femur" (1996b: 249); wing shape apomorphic (1996b: 249 & figs. 14-19) as quoted under generic diagnosis of this paper *Leptynoma* 5
- 5(4). Mouth of marginal cell wide and as long or longer than vein R_5 ; [for other distinguishing characters, see 1996b: couplet 4 on p. 250] Subgenus *Leptynoma*
 - Mouth of marginal cell narrow and shorter than vein R_5 Subgenus *Perianthomyia*
- 6(1). Antennal flagellum gradually tapering apically and with or without arista; hind coxa without ventral (= anterior) process 7
 - Antennal flagellum with an arista and with an abruptly widened base (= apparently flagellomeres 1+2) which is widest around middle and about as long to nearly twice as long as wide; hind coxa with an elongate ventral (= anterior) process at least in *V. comstocki* and *V. opacus* *Vermileo*
- 7(6). Vein R_4 almost vertical to vein R_5 and mouth of 1st submarginal cell less than 1/2 as long as mouth of 2nd submarginal cell; antennal flagellomere 1 (which is distinctly longer than wide) with a long arista (which is 3-segmented) *Vermitigris*

- Vein R₄ making an acute angle with vein R₅ and mouth of 1st submarginal cell over 1/2 as long as mouth of 2nd submarginal cell; antennal flagellomeres 2-5 gradually tapering apically and last joint (= flagellomere 8 or sometimes 6) shorter than rest of flagellum *Vermiophis*

Key 2: Genera Based on Male Genitalia

(excluding *Vermitigris* whose male genitalia are not yet studied)

1. Ejaculatory apodeme reduced in size and situated inside aedeagus (= trumpet tube) 2
 - Ejaculatory apodeme long and extruded anteriorly from aedeagus (= half-open umbrella plate or sclerotic structure) 4
- 2(1). Aedeagus without lateral extension 3
 - Aedeagus (= trumpet tube) with lateral extension (except for *cylindrica* group of *Lampromyia* which is degenerate in *Vermilynx jasoni*) *Lampromyia*, *Leptynoma* and *Vermilynx* (see DISCUSSION section)
- 3(2). "Epandrium [= tergum 9] elongate, projecting well beyond apex of synsternite" (after Stuckenberg, 1998); Figs. 34-35, 44, 48 *cylindrica* group of *Lampromyia*
 - Tergum 9 relatively much shorter *Vermipardus*
- 4(1). Ejaculatory apodeme with lateral ejaculatory process, or with posterior part bifurcate and without lateral ejaculatory process; gonocoxal apodeme (on dorsal bridge) relatively short 5
 - Ejaculatory apodeme without lateral ejaculatory process and without bifurcate posterior part; gonocoxal apodeme (on dorsal bridge) relatively long (Figs. 132, 144, 157, 164, 172, 181, 188) *Vermiophis*
- 5(4). Ejaculatory apodeme with lateral ejaculatory process and its posterior part (= endoaedeagal process) not bifurcate; fused gonocoxites without mid-posterior concavity ventrally
 - *Vermileo comstocki* from North America
 - Ejaculatory apodeme conspicuously long and without lateral ejaculatory process, and its posterior part bifurcate and long (a third to a half of the whole length), fused gonocoxites with mid-posterior concavity ventrally *Vermileo* (4 spp.) from southern Europe, Crete and Sudan

LIST OF THE GENERA AND SPECIES

1. Genus *Lampromyia* Macquart

Lampromyia Macquart, 1835, Hist. nat. Ins., Dipt., 2: 660. Type species: *Lampromyia pallida* Macquart, 1835, by monotypy.

Stuckenberg (1971, 1996b, 1998) revised *Lampromyia* from Canary Islands, southern Africa and the Palearctic region (southern Europe, north-west Africa and Canary Islands) respectively. Stuckenberg (1996b, 1998) prepared a cladogram for *Lampromyia* and recognized three species groups: *pilosula*-group (3 species from southern Africa: *pilosula*, and *flavida* + *rebecca*); *canariensis*-group (3 species from Canary Islands: *canariensis*, and *fortunata* + *hemmingseni*); *cylindrica*-group (6 species from southern Europe and north-west Africa: [*lecerifi* + *nigripennis*] + [*cylindrica* + *funbris*],

Fig. 1. *Lampromyia canariensis*. Left (A), ♀; right (B), ♂ abdomen. From Wheeler (1930).

Figs. 2-3. *Lampromyia flavida*. 2, ♂; 3, ♀. "Protrusion of ♀ sternites near the end of the abdomen is a consequence of postmortem shrinkage." From Stuckenberg (1960, 1996b).

and *pallida* + *iberica*).

Generic diagnosis (based on Stuckenberg, 1960, 1965a, 1971, 1995a, b, 1996b; quoted from 1996b:249). Head: Face produced forward beyond eye margin in association with a prolonged proboscis. The degree of facial protrusion varies with species. Antennal flagellum 3-5 (or 6) segmented, with basal segment much longer than wide, and with apical segment long, thin and arista-like. Palpal segment 2 clavate and swollen beyond middle. "Proboscis very elongate and slender, 2.5-4.5 times mesonotal length (usually more than 3.0), capable of posterior flexure between the legs

Figs. 4-17. Male genitalia of *Lampromyia* (*pilosula* group). 4-9, *L. flavida*; 10-14, *L. pilosula*; 15-17, *L. rebecca*. 4, aedeagus, dorsal view; 5, 13, aedeagus, posterior view; 6, 12, 17, aedeagus, lateral view; 7, 14, 16, tergum 9, cerci and sternum 10, ventral view; 10, 15, hypopygium, dorsal view; 11, same, ventral view; 8, cerci and sternum 10, dorsal view, 'indicating the course of the rectum'; 9, gonocoxites and gonostyli, ventral view. ae, median intromittent tube; ad, ejaculatory apodeme; cr, cercus; db, dorsal bridge; dt, dististyle (=gonostylus); ex, extension of gonocoxite; p, lateral accessory process or sclerotised projection from aedeagus; pr, proctiger (=sternum 10); r, upper edge of aedeagus base; va, ventral aperture. From Stuckenberg (1960).

Figs. 18-33. Male genitalia of *Lampromyia* (*canariensis* group). 18-22, *L. canariensis*; lettering as in Figs. 7, 9-14; cd, mediodorsal projection or 'condyle on which the articulating surface of the proctiger engage.' 23-27, *L. fortunata*; 23, fused gonocoxites, gonostyli, and aedeagus, dorsal view; 24-25, fused gonocoxites and gonostyli, ventral and lateral views; 26, aedeagus, lateral view; 27, tergum 9, cerci and sternum 10, ventral view. 28-33, *L. hemmingseni*; 31, tergum 9, dorsal view; 33, tergum 8, dorsal view, basal margin below; ae, aedeagus. From Stuckenberg (1960, 1971, 1998).

Figs. 34-43. Male genitalia of *Lampromyia* (*cylindrica* group). 34-38, *L. iberica*; 34-35, dorsal and lateral views; 36, fused gonocoxites and gonostyli, dorsal view; 37-38, aedeagus, lateral and posterior views; lettering as in Figs. 10-14. 39-43, *L. funebris*; 39-40, hypopygium, dorsal and ventral views; 41, cerci, sternum 10 and dorsal bridge, dorsal view; 42, aedeagus, lateral view; 43, 'apex of ♂ abdomen, showing form of tergum 9; T8 is largely withdrawn beneath T7 and only its posterolateral corners are exposed' (see Fig. 44); lettering as in notes under Figs. 44-55. From Stuckenberg (1960, 1998).

when the fly is at rest.a bifid pseudotrachea in each labellum." Proboscis distinctly over twice as long as fore femur.

Wing: "Wing shape unspecialised (figs. 29-31); costal and posterior wing margins more convexly curved, costal cell relatively broad; apex of wing almost symmetrically rounded; stalk relatively short, about one-sixth to one-seventh of total wing length; radial veins unspecialised." Mouth of 2nd submarginal cell shorter than vein R₅; 4th posterior cell petiolate, closed at wing margin or narrowly open; anal cell usually open but mouth of anal cell not longer than m-cu crossvein. Apical portion of

Figs. 44-55. Male genitalia, etc. of *Lampromyia* (*cylindrica* group). 44-47, *L. cylindrica*; 44, ♂ abdomen; 45, lateral view (T9 incomplete; see T9 in Fig. 44); 46, synsternite (gonocoxites) and gonostyli, ventral view; 47, aedeagus, lateral view. 48-55, *L. pallida*; 48, ♂ abdomen; 49, 'apex of ♂ abdomen, showing form of epandrium (=T9) in posterodorsal view, posterior margin of T7 adjacent to basal margin of T9, only the posterolateral corners of T8 visible'; 50, ♂ T8, oblique dorsolateral, posterior margin on left; 51-53, hypopygium, dorsal, lateral and ventral views; 54, aedeagus, lateral view; 55, tergum 9, cerci and sternum 10, ventral view; ae, aedeagus; ag, apical guides; cd, mediodorsal projection or condyle; db, dorsal bridge; ea, ejaculatory apodeme; gs, gonostylus; p, lateral accessory process or sclerotised projection from aedeagus; pr, proctiger (=sternum 10); sn, synsternite (=fused gonocoxites); va, ventral aperture. From Stuckenberg (1998).

vein R_{2+3} weakly or strongly curved upward according to species [and possibly individual too]. Distance between h-crossvein and base of R_s 2.5-2.7 times (in 3 species from southern Africa) as long as basal section of R_s .

Legs: "Hind tibia not markedly swollen apically, clearly more slender than hind femur." Hind coxa without ventral (= anterior) process.

Male genitalia (see many figures in Stuckenberg, 1960, 1971, 1996b, 1998) (= Figs. 4-55): Ejaculatory apodeme situated inside aedeagus (= trumpet tube) which has a pair of lateral processes (except for *cylindrica* group). See DISCUSSION section.

Verrall (1909: 256) wrote, "in *L. pallida* [if correct in species identification] the frons is wide at the vertex, being a quarter the width of the head, but contracts down to the transverse suture just above the antennae, where it is still more than half its width at the vertex, while below the antennae the produced face gradually widens again but not quite to the width of the vertex;"

Lampromyia is distinguished from *Leptynoma* by having the wing shape, hind tibia, and a bifid pseudotrachea in each labellum as quoted above.

1. 1. (1). *Lampromyia canariensis* Macquart (Figs. 1, 18-22, 286, 301, 323, 345-346)

Lampromyia canariensis Macquart, 1839, in Webb & Berthelot, Hist. nat. d'Iles Canaries, Ent., 2 (2): 105. Type locality: Canary Is. (Tenerife: Orotava).

Distribution. Canary Is. (Tenerife; La Palma; La Gomera; El Hierro).

Stuckenberg (1960: 247-250; 1971: 81-82; 1998) redescribed this species.

1. 2. (2). *Lampromyia cylindrica* (Fabricius) (Figs. 44-47, 291)

Empis cylindrica Fabricius, 1794, Entom. Syst., 4: 403. Type locality: Algeria (Mascara) [based on Neotype designated by Stuckenberg, 1998].

Distribution. north-west Africa (Algeria).

Stuckenberg (1998) redescribed this species. Stuckenberg (1971: 78) wrote, "Becker's (1921) record of *cylindrica* and *pallida* from the [Canary] islands is rejected." Stuckenberg (1998) said, "Fabricius described this species on an unknown number of specimens in the collection of R. L. Desfontaines. According to Zimsen (1964), Desfontaines travelled in Tunisia and Algeria around 1790, where he collected plants and insects. His collection of insects was placed in the Paris Museum in 1828, but now appears to be lost. A series from Algeria, identified as *cylindrica* by J. Villeneuve, is in the IRSN [Brussels]; it matches the description by Fabricius and is here accepted as of this species. Designation of a neotype from this series is warranted."

1. 3. (3). *Lampromyia flavida* Engel & Cuthbertson (Figs. 2-9, 274, 281, 300)

Lampromyia pilosula var. *flavida* Engel & Cuthbertson, 1937, Trans. Rhodesia Sci. Assoc., 35 (1): 2. Type locality: Zimbabwe (Umtali district: Vumba Mountains).

Distribution. Zimbabwe.

Stuckenberg (1960: 243-247) redescribed this species and gave a full specific new status for it.

1. 4. (4). *Lampromyia fortunata* Stuckenberg (Figs. 23-27, 287, 323, 334-336)

Lampromyia canariensis Macquart, Hemmingsen, 1963, Vidensk. Meddr dansk naturh. Foren.,

125: 237-267.

Lampromyia fortunata Stuckenberg, 1971, Vidensk. Meddr dansk naturh. Foren., 134: 82. Type locality: Canary Is. (Gran Canaria: Barranco del Guiniguada).

Distribution. Canary Is. (Gran Canaria).

1. 5. (5). *Lampromyia funebris* Dufour (Figs. 39-43, 290)

Lampromyia funebris Dufour, 1850, Ann. Soc. ent. Fr., (2)8: 152. Type locality: Spain (Madrid). Distribution. Europe (Spain; France).

Stuckenberg (1998) wrote, "It [= *L. funebris*] was wrongly synonymised with *cylindrica* by Osten Sacken (1883), and this was accepted by subsequent authors.....The few external characters distinguishing *funebris* from *cylindrica* are given under that species. Hypopygial characters (figs. 21, 28-31) [= Figs. 39-43]: aedeagus, male cerci and gonostyli obviously of different shape. Spermathecae like those in *lecerfi* (fig. 12)" [= Fig. 302].

1. 6. (6). *Lampromyia hemmingseni* Stuckenberg (Figs. 28-33, 288, 296-297, 323)

Lampromyia hemmingseni Stuckenberg, 1971, Vidensk. Meddr dansk naturh. Foren., 134: 85. Type locality: Canary Is. (Lanzarote: Puente de la Vega de Gualiza).

Distribution. Canary Is. (Lanzarote; Fuerteventura).

1. 7. (7). *Lampromyia iberica* Stuckenberg (Figs. 34-38, 289)

Lampromyia iberica Stuckenberg, 1998, Bonn. Zool. Beitr., 48: 87. Type locality: Spain (Zaragoza).

Distribution. Spain.

"Rare in collections, the specimens available date from more than half a century ago. On present evidence, *iberica* seems confined to north-eastern Spain; there are no records supporting the statement by Séguy (1926) that it occurs in the south. It may be significant that most specimens come from the Zaragoza region; this includes the Monegros," (Stuckenberg, 1998: 87).

1. 8. (8). *Lampromyia lecerfi* Séguy (Figs. 292, 298-299, 302)

Lampromyia lecerfi Séguy, 1928, Bull. Soc. ent. Fr., 1928: 45. Type locality: Morocco (High Atlas).

Distribution. Morocco (High Atlas).

The male of this species is unknown. "This species is known only from the western end of the High Atlas Range, in the region of the Toubkal, Massif" (Stuckenberg, 1998: 83).

Stuckenberg (1971: 86) wrote, "In external features *hemmingseni* is most similar to *lecerfi* Ség., which is based on a female holotype, kindly loaned for comparison by Dr. L. Tsacas of the Paris Museum; *hemmingseni* is distinguished, *inter alia*, by having the median mesonotal vitta divided in the midline (undivided in *lecerfi*), the wing membrane much darker grey, the stigma darker, and the apical part of R_{2+3} curiously flexed [see fig. 4] (this evenly curved forward to costa in *lecerfi*)." For distinguishing characters between *lecerfi* and *nigripennis*, see key to species of NW Africa in Stuckenberg (1998).

1. 9. (9). *Lampromyia nigripennis* Séguy

Lampromyia nigripennis Séguy, 1930, Mém. Soc. Sci. nat. phys. du Maroc., 24: 68. Type locality: Morocco (Middle Atlas).

Distribution. Morocco (Middle Atlas).

The male of this species is unknown. For redescription of this species, "known only from its type locality at the north-eastern end of the Middle Atlas Range", see Stuckenberg (1998).

1. 10. (10). *Lampromyia pallida* Macquart (Figs. 48-55, 293, 303)

Lampromyia pallida Macquart, 1835, Hist. nat. Ins., Dipt., 2: 662. Type locality: Algeria (Oran).

Lampromyia miki Marchal, 1897, Mém. Soc. zool. Fr., 10: 10. Type locality: Tunisia (Tunis).

Distribution. north-west Africa.

"Not *Lampromyia pallida* Macquart, sensu Lindner, 1925, pp. 6, 7; Séguy, 1926, p. 93; 1934, p. 20; Stuckenberg, 1960, pp. 250-2, figs 42-45 (Misidentifications of *L. iberica* sp. n.).....The species [= true *pallida*] is thus known from coastal and hinterland districts of Tunisia and Algeria, from Tunis to Oran; it probably occurs westward in Morocco." (Stuckenberg, 1998).

1. 11. (11). *Lampromyia pilosula* Engel (Figs. 10-14, 279)

Lampromyia pilosula Engel, 1929, Ann. Transvaal Mus., 13: 172. Type locality: South Africa (Eastern Cape Province: Aliwal North).

Distribution. South Africa; Lesotho.

Stuckenberg (1960: 241-243) redescribed this species.

1. 12. (12). *Lampromyia rebecca* Stuckenberg (Figs. 15-17, 280)

Lampromyia rebecca Stuckenberg, 1996, Ann. Natal Mus., 37: 263. Type locality: Harare, Zimbabwe.

Distribution. Zimbabwe.

2. Genus *Leptynoma* Westwood

Leptynoma Westwood, 1876, Trans. ent. Soc. Lond., 1876: 517. Type species: *Leptynoma sericea* Westwood, 1876, by monotypy.

Leptynoma was separated from *Lampromyia* by Stuckenberg (1996b) who gave a key to Afrotropical genera of *Lampromyia* et al. (*Lampromyia*, *Leptynoma*, *Vermilynx* and *Vermipardus*) and a cladogram showing relationships among the four genera and among the four species of the subgenus *Leptynoma* respectively. Stuckenberg (1996b) also made a key to species of *Leptynoma*.

Leptynoma (s. str.): *sericea*, and *appendiculata* + (*hessei* + *namaquaensis*); *Leptynoma* (*Perianthomyia*): *maculata* and *phantasma*.

Generic diagnosis (based on Stuckenberg, 1960, 1995b, 1996b and on *L. namaquaensis* we have seen directly; quoted from 1996b: 249-251). Similar to *Lampromyia* except as follows.

Head: Antennal flagellum 6-8 segmented, with basal segment more robust and as long or somewhat longer than wide. "A trifold pseudotrachea in each labellum."

"Wing shape apomorphic (figs. 14-19), stalk long, about one-fifth to one-quarter of total wing length, form narrower, with more nearly parallel fore and hind margins; costal cell narrow, costal

Fig. 56. *Leptynoma* (*Leptynoma*) *hessei*, ♂.
From Stuckenberg (1960).

Fig. 57. Wing venation in *Leptynoma* (*Leptynoma*) *namaquaensis*, ♂.
Darkened parts of wing membrane are omitted. Refer to pp. 131, 141. From Stuckenberg (1960).

margin almost straight; wing apex more bluntly curved, asymmetrical, its most distal part posterior to median longitudinal axis of wing; radial veins usually with conspicuous modifications." Mouth of 2nd submarginal cell as long or longer than vein R₅ in the subgenus *Leptynoma*; vein R₂₊₃ angulated at apical portion in the subgenus *Leptynoma* and often so at basal portion. Distance between h-crossvein and base of R₅ about 1.5-2.0 times (Stuckenberg, 1996b: figs. 14-19) as long as basal section of R₅. In *L. namaquaensis*, vein R₂ always present.

Legs: "Hind legs appearing robust because tibia is swollen apically as much as femur."

Male genitalia: See Figs. 58-77 and DISCUSSION section.

In the subgenus *Leptynoma*, "frons narrower, clearly longer than wide, usually narrowing towards ocelli."

In the subgenus *Perianthomyia*, "frons subquadrangular, about as long as wide, wider towards ocellar tubercle."

L. (L.) namaquaensis: Frons at median ocellus less than 1/2 as wide as one eye on a mid line, less than twice as wide as ocellar triangle, and distinctly (♂) or somewhat (♀) narrower than that just above antenna (space between ocellar triangle and eye narrow especially in ♂; face gradually wider towards proboscis and wider than frons).

Verrall (1909: 256) wrote, [under *L. (L.) sericea*], "the frons is narrower but still with parallel sides, and the face is twice as wide."

Leptynoma is separated from *Vermilynx* and *Vermipardus* by having last joint of flagellum much

Figs. 58-69. Male genitalia of *Leptynoma* (*Leptynoma*). 58-60, *L. appendiculata*; 61-63, *L. hessei*; 64-66, *L. namaquaensis*; 67-69, *L. sericea*. 58, 61, 64, 67, fused gonocoxites and gonostyli, dorsal view; 'inset is the lefthand gonostylus in the horizontal plane.' 59, 62, 65, 68, aedeagus, posterior view. 60, 63, 66, 69, aedeagus, lateral view. For lettering, see notes under Figs. 4-17. From Stuckenberg (1960).

longer than preceding 3 joints together.

Subgenus *Leptynoma* Westwood

Mouth of 2nd submarginal cell as long or longer than vein R₅. For key to species of this subgenus, see Stuckenberg (1996b: 254).

2. 1. (13). *Leptynoma* (*Leptynoma*) *appendiculata* (Bezzi) (Figs. 58-60, 278)

Lampromyia appendiculata Bezzi, 1926, Ann. S. Afr. Mus., 23 (2): 301. Type locality: South Africa (SW Cape Province: Ceres Division: Matroosberg).

Distribution. South Africa.

Stuckenberg (1960: 238-239) redescribed this species.

2. 2. (14). *Leptynoma* (*Leptynoma*) *hessei* (Stuckenberg) (Figs. 56, 61-63, 275)

Figs. 70-77. Male genitalia of *Leptynoma* (*Perianthomyia*). 70-73, *L. maculata*; 74-77, *L. phantasma*. 70, 74, tergum 9, cerci and sternum 10, ventral view; 71, 75, hypopygium, dorsal view; 72, 76, aedeagus, lateral view; 73, 77, fused gonocoxites and gonostyli, ventral view. ag, apical guide; db, dorsal bridge; sn, synsternite (=fused gonocoxites); va, ventral aperture. From Stuckenberg (1996b).

Lampromyia hessei Stuckenberg, 1960, S. Afr. anim. Life, 7: 233. Type locality: South Africa (SW Cape Province: Oudtshoorn dist.: Zebra).

Distribution. South Africa.

2. 3. (15). *Leptynoma* (*Leptynoma*) *namaquaensis* (Stuckenberg) (Figs. 57, 64-66, 277)

Lampromyia namaquaensis Stuckenberg, 1960, S. Afr. anim. Life, 7: 228. Type locality: South Africa (SW Cape Province: Van Rhynsdorp clist.: Knersvlakte).

Distribution. South Africa.

2. 4. (16). *Leptynoma* (*Leptynoma*) *sericea* Westwood (Figs. 67-69, 337-338, 347-348, 354-356, 391-

394, 400-401)

Leptynoma sericea Westwood, 1876, Trans. ent. Soc. Lond., 1876: 517. Type locality: South Africa (SW Cape Province).

Lampromyia argentata Bigot, 1885, Bull. ent. Soc. Fr., (6)5: 68. Type locality: South Africa (SW Cape Province: Hex River [after Stuckenberg, 1960: 236]).

Distribution. South Africa.

Stuckenberg (1960: 236-238) redescribed this species and confirmed *L. argentata* as a synonym of *L. sericea*. Bezzi (1926: 299-301) also redescribed a male of *L. sericea*.

Subgenus *Perianthomyia* Stuckenberg

Perianthomyia Stuckenberg, 1996, Ann. Natal Mus., 37: 257. Type species: *Lampromyia maculata* Stuckenberg, 1960.

Mouth of 2nd submarginal cell much shorter than vein R₅. For key to species of this subgenus, see Stuckenberg (1996b: 257).

2. 5. (17). *Leptynoma* (*Perianthomyia*) *maculata* (Stuckenberg) (Figs. 70-73, 276)

Lampromyia maculata Stuckenberg, 1960, S. Afr. anim. Life, 7: 239. Type locality: South Africa (SW Cape Province: Beaufort West dist.: Merweville, Koup Karoo).

Distribution. South Africa.

2. 6. (18). *Leptynoma* (*Perianthomyia*) *phantasma* Stuckenberg (Figs. 74-77)

Leptynoma (*Perianthomyia*) *phantasma* Stuckenberg, 1996, Ann. Natal Mus., 37: 260. Type locality: Uguchab River nr / Aurusberg, SW Africa Namibia.

Distribution. SW Africa Namibia.

3. Genus *Vermileo* Macquart

Vermileo Macquart, 1834, Hist. nat. Ins., Dipt. 1: 428. Type species: *Vermileo degeerii* Macquart, 1834 (= *Vermileo vermileo* Linnaeus, 1758) by monotypy.

Pheneus Walker, 1852, [In Saunders, W. W. ed.], Insecta Saundersiana. Diptera. London. p. 155. Type species: *Pheneus tibialis* Walker, 1852 by monotypy.

Arthrostylum Williston, 1895, Kansas Univ. Quart., 4: 108. Type species: *Arthrostylum fascipennis* Williston, 1895 by monotypy.

Verrall (1909: 258) wrote, "In 1840 Blanchard is said to have proposed the name *Psammorycter* for the same genus [= *Vermileo*] (in spite of an existing *Psammoryctes* Poepp. Mamm. 1836, and subfamily Psammoryctinae), and that name was adopted by Loew, who founded on it in 1874 (Berl. Ent. Zeitschr., xviii, 381) the subfamily *Psammorycterina*; it is certain that Loew only adopted the name *Psammorycter* because he refused to accept Macquart's generic name of Latin derivation. The same genus was again described by Perris in 1852 under the name of *Apogon*, but there was already a prior genus of that name."

Generic diagnosis (based on *V. comstocki* and *V. opacus* from North America).

Head: Frons at median ocellus roughly twice as wide as that just above antennae and over twice as wide as ocellar triangle, and distinctly over 1/2 as wide as one eye on a mid line. Space between

Figs. 78-85. *Vermileo comstocki*. 78-82, ♀; 83-84, ♂. 79, Head, anterior view; 80, antenna; 81, apical portion of tarsus; 82, wing with abnormal venation; 83-84, genitalia, dorsal and lateral views; 85, eggs. Refer to pp. 91, 138. From Wheeler (1930).

ocellar triangle and eye roughly as wide as ocellar triangle and not concave but flat. Face (except wider lower portion) parallel-sided and not wider than frons. Occiput swollen. Antennal flagellum with a slender 3-segmented arista, and with a widened base which is apparently composed of flagellomeres 1+2, widest around middle and about as long or not over twice as long as wide. Palpal segment 2 widened or swollen in a lateral view, and without an elongate portion gradually tapering apically. Proboscis short and fleshy.

Wing: Mouth of 2nd submarginal cell shorter than vein R₅; vein M₃ often incomplete; 4th posterior cell (when vein M₃ complete) petiolate or wide open (see fig. 29E in Wheeler, 1930); mouth of anal cell often longer (in *V. comstocki* and *V. opacus*) or shorter (in *V. vermileo* [= *V. degeerii*]) than m-cu crossvein (Verrall, 1909: figs. 184-185; Séguy, 1926: fig. 248). In *V. comstocki*, apical portion of vein R₂₊₃ either almost straight or curved upward according to individual.

Legs: Apex of ventral (= anterior) side of hind coxa with an elongate process.

Male genitalia (see Nagatomi, 1984: figs. 143-146 on *V. comstocki*) [= Figs. 87-91]: Ejaculatory apodeme large and extending far beyond anterior margin of aedeagus (= 'half-opened umbrella plate'). Endoaedeagal process about 1/2 as long as ejaculatory apodeme.

Note on face of *Vermileo*

Wheeler (1931: 167-168) wrote as to *Vermileo dowi* from Cuba, "[Female] face and front narrow, linear, parallel-sided, of equal width.....[Male] face and front not narrower than in the female,"

DeLeon (1938: 170-171) wrote, "[In *Vermileo willetti* from Mexico], Male.....inner margin of

Fig. 86. Wing of *Vermileo tibialis*, ♂.
From Teskey (1981b).

compound eyes along front nearly parallel up to vertex, ocelli set on a distinct prominence which is narrowed and shining behind and below, lateral ocelli closer to inner margin of compound eyes than the width of the anterior ocellus; occiput flat.....; Female.....; similar to male but; inner margin of compound eyes along front more distinctly divergent towards the apex, lateral ocelli scarcely more distinct from inner margin of compound eyes than the width of anterior ocellus;....." Verrall (1909: 256) wrote, "in *Vermileo* [probably *vermileo* from Europe] the frons is about one-sixth the width of the head and has parallel sides, the face being if anything narrower than the frons and also continued downwards with parallel sides; back of the head very little inflated even on the lower part"

Thus in *V. comstocki* and *V. opacus* frons is wider at vertex and occiput is swollen, but in *V. dowi*, *V. vermileo* and *V. willetti* frons is parallel-sided and occiput is flat (this is not certain in *V. willetti*).

Note on the male genitalia of the Palearctic *Vermileo*

The male genitalia of the Palearctic *Vermileo* (including 4 species) are markedly different from those of *Vermileo comstocki* as shown in Key 2 and in Figs. 87-91, 97-113. There are two terms used by Stuckenberg (1965b), that is, (1) sclerotic structure and (2) aedeagus. They are interpreted here as (1) aedeagus (= half-opened umbrella plate sensu Nagatomi, 1984) and (2) ejaculatory apodeme (= anterior bar of aedeagus sensu Nagatomi, 1984) + endoaedeagal process (= posterior part of aedeagus sensu Nagatomi, 1984).

Key 3: New World species of *Vermileo* (*V. willetti* is not included)

(from Pechuman, 1938)

1. Wings without dark markings; fourth posterior cell usually open although sometimes much narrowed at margin 2
- Wings with dark markings; fourth posterior cell usually closed 3
- 2(1). Ground color of mesonotum and scutellum dull grayish brown; the three thoracic stripes are a dull brown, the middle one split by a grayish band which is usually as wide as either half of the stripe; wings with a brownish tinge. (California) *V. comstocki*
- Ground color of mesonotum and scutellum yellowish brown becoming very pale in region of humeri; the three thoracic stripes are a shining brown, the middle one split by a fine yellowish line which sometimes may be almost invisible; wings practically hyaline. (Calif., Nevada, Utah, New Mexico) *V. opacus*
- 3(1). Face bare; apical spot becomes somewhat paler at extreme tip of wing. (Mexico) *V. fascipennis*

Figs. 87-96. Male genitalia. 87-91, *Vermileo comstocki*; 92-96, *Vermipardus intermedius*. 87, 94, lateral view; 88, 90, 92, 95, dorsal view; 89, 91, 93, 96, ventral view. aap, anterior bar of aedeagus (=ejaculatory apodeme); ads, aedeagal dorso-anterior sclerite (=lateral ejaculatory process; endophallic sclerite); b, basistyle (=gonocoxite); bdp, basistylar dorso-medial anterior process (=gonocoxal apodeme); c, cercus; d, dististyle (=gonostylus); db, dorsal bridge; h, half-opened umbrella-like plate (=aedeagus); pa, posterior part of aedeagus (=endoaedeagal process); t, trumpet-like tube (=aedeagus). Refer to p. 136. 87, from Teskey(1981b); 88-96, from Nagatomi (1984).

Figs. 97-101. Male genitalia of *Vermileo*. 97-99, *V. ater*; 100-101, *V. nigriventris*. 97, hypopygium, dorsal view; 'inset to the left the sclerotic structure enclosing the aedeagus in lateral view, to the right some of the hairs on the outer surface of the membranous sheath.' 98, 100, tergum 9, cerci and sternum 10, dorsal view; 99, 101, same, posterodorsal view. ae, aedeagus (=correctly ejaculatory apodeme + endoaedeagal process); db, dorsal bridge; dst, dististyle (=gonostylus); ms, membranous sheath; s, sclerotic structure (=correctly aedeagus); 9sb, fused gonocoxites. Refer to p. 136. From Stuckenberg (1965b).

- Face densely white pollinose; apical spot of uniform density 4
- 4(3). First antennal segment about one and one half times as long as broad; posterior half of abdominal tergites not infuscated but first tergite bearing, antero-dorsally, a black transverse band; cross-band of wing narrow and somewhat irregular. (Jamaica) *V. tibialis*
- First antennal segment more than one and one half times as long as broad; posterior half of abdominal tergites brownish; first tergite without a transverse band; cross-band of wing broader and with relatively straight margins. (Cuba) *V. dowi*

Key 4: Palearctic species of *Vermileo* based on male genitalia

(prepared from Stuckenberg, 1965b)

1. Each posterior corner of tergum 9 with two lobes; Figs. 99, 112 2
- Each posterior corner of tergum 9 with one lobe; Fig. 101 3
- 2(1). Fused gonocoxites without desclerotised part ventrally (Fig. 109); gonostylus stouter; dorsal bridge, gonocoxal apodeme, and aedeagus (= sclerotic structure beneath dorsal bridge) as in Fig. 108 *V. vermileo*
- Fused gonocoxites with a large semi-membranous, median area ventrally (Fig. 104); gonostylus slender; dorsal bridge, gonocoxal apodeme, and aedeagus as in Fig. 105 *V. nigriventris*
- 3(1). Two lobes at each posterior corner of tergum 9 greater in length; gonostylus with a basal dilation extended into a finger-like projection; ventral hind margin of fused gonocoxites irregular in shape; dorsal bridge with postero-lateral process; bifurcate endoaedeagal process (= bifurcation of the aedeagus sensu Stuckenberg, 1965b) over apical half of ejaculatory apodeme + endoaedeagal process; Figs. 97-99, 192 *V. ater*
- Two lobes at each posterior corner of tergum 9 relatively much shorter; gonostylus long, narrow and strongly sinuous (Fig. 107); ventral hind margin of fused gonocoxites almost straight; dorsal bridge without postero-lateral process; endoaedeagal process appears to be bifid for only the

Figs. 102-109. Male genitalia, etc. of *Vermileo*. 102-103, *V. ater*; 104-106, *V. nigriventris*; 107, *V. niloticus*; 108-109, *V. vermileo*. 102, fused gonocoxites and gonostyli, posteroventral view; 104, 109, hypopygium, ventral view; 105, 108, same, dorsal view (with ejaculatory apodeme + endoaedeagal process partly omitted); 103, 106, basal portion of ♂ abdomen, lateral view. Refer to p. 136. From Stuckenberg (1965b).

apical third of ejaculatory apodeme + endoaedeagal process *V. niloticus*

3. 1. (19). *Vermileo ater* Stuckenberg (Figs. 97-99, 102-103)

Vermileo ater Stuckenberg, 1965, Ann. Mag. nat. Hist., 13 (8): 498. Type locality: Greece (Crete: St. Nicolo).

Distribution. Greece (Crete).

"Known only in the male, similar to *nigriventris* in that sex in the intensity and extent of the dark markings, *ater* being distinguished from that species by the more uniformly coloured antennae, anteriorly coalescent mesonotal stripes, presence of basal and apical yellowish markings on tergites 2-4 only, and the form of these tergites in profile; *nigriventris* has the third antennal segment distinctly darker than the others, the mesonotal stripes clearly defined and separated anteriorly, yellowish markings on tergites 2-5 or 2-6 at the base only, and tergites 2-4 only slightly undulant in profile." (from original description).

3. 2. (20). *Vermileo comstocki* Wheeler (Figs. 78-85, 87-91, 258-262, 295, 331, 343-344, 351-353, 358, 383-390, 395-396, 413)

Vermileo comstocki Wheeler, 1918, Proc. New Engl. Zool. Club, 6: 84. Type locality: California (Alta Meadow).

Distribution. North America (California).

3. 3. (21). *Vermileo dowi* Wheeler

Vermileo tibialis var. *dowi* Wheeler, 1931, Psyche, Camb., 38: 167. Type locality: Cuba (Trinidad Mts.).

Distribution. Cuba.

Pechuman (1938: 85) wrote, "Finally Wheeler (1931, p. 167) described *Vermileo tibialis* (Walker) var. *dowi* from Cuba. However, on the receipt of the specimen of *V. tibialis* from Jamaica mentioned above, Wheeler decided that *dowi* was probably subspecifically distinct. The writer is inclined to regard them as distinct species."

3. 4. (22). *Vermileo fascipennis* (Williston)

Arthrostylum fascipennis Williston, 1895, Kansas Univ. Quart., 4: 109. Type locality: Mexico (Guerrero: Xucumanatlan).

Distribution. Mexico.

3. 5. (23). *Vermileo nigriventris* Stroble (Figs. 100-101, 104-105)

Vermileo vermileo nigriventris Stroble, 1906, Beiträg. Mem. Sos. Españ. Hist. Nat., 3: 271. Type locality: Spain (Cercadilla and Los Molinos).

Distribution. Spain; southern Portugal ("Praia de Rocha").

Stuckenberg (1965b: 495) wrote, "After a study of the British Museum material described by Edwards, including the preparations of genitalia, I have concluded that *nigriventris* is a valid species. There are small but definite differences in the genitalia (which have been confirmed in new preparations), and the differences in colouration are considerable and not bridged by intermediate specimens. At present I have no records which indicate that *vermileo* and *nigriventris* are sympatric; the latter is known only from Spain, whereas *vermileo* is recorded also from there but from different localities in addition to its wide range in southern Europe generally. Nevertheless, the differences mentioned are considered to be sufficient to warrant the recognition of *nigriventris* as a full species and not a subspecies of *vermileo*."

Edwards (1935) wrote, "Mr. Main's specimens from Praia de Rocha, South Portugal, show the following differences in colouring from typical *V. vermileo* as described above. (1) Third antennal segment all dark. (2) Palpi blackish. (3) Pronotum dark. (4) Middle dark stripe undivided and (except in some ♀♀) reaching back to scutellum. (5) Scutellum blackish, at least in ♂. (6) Dark area in middle of pleurae extending on to pteropleura. (7) Hypopleura largely dark. (8) Hind coxae blackish, other coxae also largely darkened. (9) Hind femora and tibiae blackish on at least the distal two-thirds. (10) Wings with a small smoky area towards outer end of upper basal cell. (11) Abdomen mainly black; in ♂ and in darker ♀♀ with the bases of segments 3-6 yellow, 2 and 7-9 entirely black; in lighter ♀♀ also with posterior margins of segments 2-7 yellowish."

3. 6. (24). *Vermileo niloticus* Edwards (Figs. 107, 112-113)

Vermileo niloticus Edwards, 1935, Proc. R. ent. Soc. London, 10: 56. Type locality: Sudan.

Distribution. Sudan.

Stuckenberg (1965b: 498) wrote, "*V. niloticus* Edwards is similar in colouration to *vermileo* but resembles *ater* in the structure of the hypopygium, differing from the latter in that the two projections on each posterior corner of the ninth tergite are relatively much shorter; the dististyles are long, narrow and strongly sinuous (fig. 6, inset) [= Fig. 107]; the hind margin of the synsternite is almost straight; the dorsal bridge has a different shape and is simpler, consisting of a median sclerite

supported by long, narrow struts and supporting in turn the sclerotic sheath [= half-opened umbrella plate] embracing the aedeagus; the aedeagus [= ejaculatory apodeme] appears to be bifid for only the apical third of its length." Edwards (1935: 54) wrote, "Perhaps the record of *V. vermileo* from Egypt may refer to this new species [= *V. niloticus*] rather than to the true *V. vermileo*."

"In general very similar to *V. vermileo*, but showing many small differences. Palpi yellow, but shorter than in *V. vermileo*; third antennal segment also shorter. Face slightly narrowed below, which is not the case in *V. vermileo*. Pronotum yellow. Middle dark stripe of mesonotum undivided, but not reaching back to scutellum. Scutellum blackish; a large blackish patch in middle of pleurae, which, however, does not extend on to pteropleura; hypopleura dark. Hind coxae all dark; hind femora and tibiae not very dark. Abdomen with yellowish ground-colour; tergites 2-6 each with large but ill-defined dark dorsal spots, and with less obvious dark lateral spots; 7 all dark." (from original description).

3. 7. (25). *Vermileo opacus* (Coquillett)

Pheneus opacus Coquillett, 1904, [In Baker, C. F.], Invertebrata Pacifica, 1: 21. Type locality: Nevada (Ormsby Co.).

Distribution. North America (California to New Mexico and Colorado).

3. 8. (26). *Vermileo tibialis* (Walker) (Fig. 86)

Pheneus tibialis Walker, 1852, [In Saunders, W. W. ed.], Insecta Saundersiana. Diptera. London. p. 156. Type locality: Jamaica.

Distribution. Jamaica.

3. 9. (27). *Vermileo vermileo* (Linnaeus) (Figs. 108-109, 110-111, 268-273, 329-330, 341-342, 359-363, 381-382, 403, 405-411)

Musca vermileo Linnaeus, 1758, Systema Naturae, Ed. 10: 590. Type locality: southern France (after Wheeler, 1930: 164).

Vermileo degeeri Macquart, 1834, Hist. nat. Ins., Dipt., 1: 428. Unjustified replaced name.

Apogon dufouri Perris, 1852, Ann. Soc. Linn. Lyon, 5: 199. Type locality: France (Grandes Landes).

Distribution. Southern Europe (Spain; France; Italy; southern Tyrol; Dalmatia; Balearic Is.; Bulgaria [after Majer, 1988: 34]).

Hemmingsen (1968: 291) wrote, "In France it occurs as far North as the environs of Paris and Bur-sur-Seine (Séguy, 1926). Szilády (1934, p. 231) mentions a specimen allegedly from Bavaria. Schrank (1781, Ins. Austr.) reports it from Linz, Austria."

Stuckenberg (1965b: 495) wrote, "The species of worm-lion occurring in Egypt was recorded by Efflatoun (in Wheeler, 1930: 148) on larval material as *vermileo*, and that name was applied to the species discussed by Hafez & El-Moursy (1956a, b, 1964), but, as Edwards surmised, the species is more likely to be *niloticus*." Stuckenberg (1971: 78) also wrote, "While it is not impossible that a *Vermileo* might occur there [= Canary Is.], Brauer's (1883) record must be considered as doubtful in view of the absence of this genus in the extensive material collected by Dr. Hemmingsen."

Wheeler (1930: 164) wrote, "I have not been able to compare my specimens with material taken

Figs. 110-113. Male genital parts of *Vermileo*.
110-111, *V. vermileo*; 112-113, *V. niloticus*.
110, 112, tergum 9 and cerci, dorsal view;
111, 113, fused gonocoxites and gonostyli,
ventral view. Refer to p. 136. From Edwards
(1935).

in Southern France, the type locality, in Tyrol or Italy, but the published descriptions indicate that the specimens from the Balearic Islands [Mallorca and Menorca] probably represent a distinct variety, for which I propose the name *balearicus* var. nov. The general yellow color of the body and appendages seems to be paler or duller. Other slight differences will probably be revealed by careful comparison with typical continental specimens."

Edwards (1935: 55) wrote, "*V. v. nigriventris* is clearly not the same as *V. v. balearicus* Wheeler from Majorca; the description of the latter does not indicate any well-marked distinction from *V. v. vermileo* L."

Hennig (1967: fig. 31) [= Fig. 301] prepared a distribution map of the genus *Vermileo* in which Turkey and Syria are included. Majer (1988: 34) cited Syria in the distribution of *V. vermileo*. It is unknown to us where their original records are.

"(1) Third antennal segment mostly yellowish, darkened on upper edge only. (2) Palpi yellow. (3) Pronotum yellow. (4) Middle dark stripe of mesonotum divided by a pale area except in front and not reaching back beyond middle of thorax. (5) Scutellum yellowish. (6) Dark area in middle pleurae not extending on to pteropleura. (7) Hypopleura yellowish. (8) Hind coxa with a dark patch behind, fore and mid coxae wholly yellowish. (9) Hind femora and tibiae indefinitely darkened towards tips. (10) Wings practically clear. (11) Abdomen largely yellow; in ♀ with median and lateral dark spots on each tergite; in ♂ with these spots larger and fused into dark bands on each of tergites 2-4, but leaving the anterior and posterior margins of these tergites, as well as most of tergites 5-7 yellow" (from Edwards, 1935).

"The features of colouration which distinguish *Vermileo vermileo* (Linn.) from *nigriventris* are listed by Edwards [1935], but it should be noted that the 'yellow' markings he describes vary from pale luteous to pale yellowish-brown" (from Stuckenberg, 1965b: 498).

3. 10. (28). *Vermileo willetti* DeLeon

Vermileo willetti DeLeon, 1938, Entomol. News, 49: 170. Type locality: Mexico (between Mexico, D. F., and Cuernavaca, Morelos).

Distribution. Mexico.

"This species seems to be allied to *V. fascipennis* (Will.) but differs from it by the front and face being grey-pollinose whereas the front of *fascipennis* is brown and the face opaque yellow; by the first two joints of the antennae being yellow, while in *fascipennis* they both are partly black; by the pleura

being yellow-brown instead of black; and by the abdomen having five transverse yellowish bands instead of being reddish and without any bands" (from the original description).

4. Genus *Vermilynx* Stuckenberg

Vermilynx Stuckenberg, 1995, African Entomology, 3: 30. Type species: *Lampromyia* (*Vermipardus*) *vansoni* Stuckenberg, 1965, by original designation.

Generic diagnosis (based on Stuckenberg, 1960, 1965a, 1995a, b). Similar to *Lampromyia* except as follows: Head: Antennal flagellum as in *Vermipardus* (Stuckenberg, 1965a: fig. 1) [= Fig. 116].

Proboscis 1.5-1.7 times as long as fore femur or about 1.3-1.6 times as long as mesonotum. "One trifold pseudotrachea in each labellum" (1996b: 249) [as in *Leptynoma*].

Wing: Distance between h crossvein and base of R_s about 2.0-2.5 times as long as basal section of R_s .

Legs: At least in *V. vansoni*, pulvilli and empodium reduced in size. Pulvilli and apical portion of empodium triangular in shape (Stuckenberg, 1965a: fig. 2) [= Fig. 117]. In *Leptynoma*, "the hind tibia is considerably swollen apically and is as enlarged as the hind femur; this condition also occurs in [*Vermilynx*] though it is not so marked in the female" (1996b: 245).

Male genitalia (Figs. 118-124): see Stuckenberg (1996a: 198) or DISCUSSION section.

Stuckenberg (1965a: 107) wrote as to *V. vansoni*, "Female:.....Face unusually prominent [see Fig. 116].....Frons slightly broader than high,.....ocelli large, close together, in a triangular with base slightly longer than sides."

Vermilynx is distinguished from *Vermipardus* by having "Labella elongate tubular, dorsoventrally flattened in dried specimens, separated over their entire length though usually held pressed together, without setae [as in *Lampromyia* and *Leptynoma*]; one trifold pseudotrachea in each labellum [as in *Leptynoma*]; aedeagus trifold, prominent lateral extensions present [in *V. vansoni*; this is degenerate in *V. jasoni*] [as in *Lampromyia* (except *cylindrica* group) and *Leptynoma*]" (1996b: 249).

It is separated from *Leptynoma* by having last segment of flagellum not longer than preceding two segments together.

Stuckenberg wrote to one of us (A. N.) (pers. comm., 19. iii. 1997) that "The characterisation of *Vermilynx* given in my latest paper on the genus will have to be amended. Two new, closely related species which I reared from larvae this summer show that antennal dimorphism is only a group character of the *vansoni-jasoni* species pair. The two new species have the same, slender antennae in both sexes."

4. 1. (29). *Vermilynx jasoni* Stuckenberg (Figs. 121-124, 284-285)

Vermilynx jasoni Stuckenberg, 1996, African Entomology, 4: 198. Type locality: South Africa (Namaqualand: Goegap Nature Reserve near Springbok).

Distribution. South Africa.

4. 2. (30). *Vermilynx vansoni* (Stuckenberg) (Figs. 114-120)

Lampromyia (*Vermipardus*) *vansoni* Stuckenberg, 1965, J. Ent. Soc. S. Africa, 28: 107. Type locality: South Africa (Northern Cape province: Richtersveld, C. P.: Brandkaross).

Distribution. South Africa; Namibia.

Figs. 114-117. *Vermilynx vansoni*. 114-115, ♂; 116-117, ♀. 117, 'Tarsal claws and pretarsus of middle leg, ventral view.' Refer to pp. 129, 131, 137. From Stuckenberg (1965a, 1995a, 1996b).

5. Genus *Vermiophis* C. Yang

Vermiophis C. Yang, 1979, Entomotaxon., 1: 84. Type species: *Vermiophis ganquanensis* C. Yang, 1979, by original designation.

Generic diagnosis.

Head: Frons at median ocellus not over 1.5 times as wide as that just above antennae, not over twice as wide as ocellar triangle and one eye on a mid line respectively. Space between ocellar triangle and eye distinctly narrower than ocellar triangle and more or less concave. Face (except wider lower portion) parallel-sided, and about as wide as or narrower than frons. Antennal flagellomeres 2-5 gradually tapering apically and flagellomeres 6-8 (or sometimes only 6) forming or not forming arista according to species. Palpal segment 2 widest around middle and then gradually tapering apically.

Wing: Mouth of anal cell shorter than m-cu crossvein. According to individual within the same

Figs. 118-124. Male genitalia of *Vermilynx*. 118-120. *V. vansoni*; 121-124, *V. jasoni*. 118, 121, hypopygium, dorsal view; 'aedeagus of *V. jasoni* exceptional in lacking lateral extensions, although these may be represented by an obtuse subtriangular projection on each side of aedeagus towards its base (see Fig. 202, the projection on right side of aedeagus is shown).' 119, 123, tergum 9, cerci and sternum 10, ventral view. 122, gonostyli and apex of fused gonocoxites, ventral view. le, lateral expansion of aedeagus; md, mediadorsal projection of upper margin of gonocoxite; pr, cerci + sternum 10; vf, ventral flange of tergum 9; for other letterings, see notes under Figs. 44-55. From Stuckenberg (1995a, 1996a).

species, vein R_{2+3} weakly or rather strongly curved upward apically, mouth of 2nd submarginal cell roughly as long as or shorter than vein R_5 , and 4th posterior cell petiolate or open.

Legs: Hind coxa without ventral (= anterior) knob.

Male genitalia: Ejaculatory apodeme large and extending far beyond anterior margin of aedeagus

(= 'half-opened umbrella plate').

Details of male genitalia

Fused gonocoxites: As wide as long or roughly so; paired gonocoxites completely fused with each other at ventral surface and sternum 9 absent; a pair of desclerotized dorsomedial corners fused with each other through dorsal bridge [whose lateral boundary was indistinct in several individuals examined]; a pair of dorsomedial posterior projections varied in size and shape and sometimes absent; ventral fused gonocoxites (= synsternite) with a deep mid-posterior concavity and the posterolateral part thus formed varied in size and shape; dorsal bridge with paired long gonocoxal apodemes.

Gonostylus: Varying in shape with visual angle; usually pointed at apex; in lateral view, usually widened basally; [the specific character not studied here carefully].

Aedeagus: Temple bell like in shape, and with posterior part curved or protruded dorsally.

Ejaculatory apodeme: Large, and with widened anterior part which is elliptical or rounded in lateral view.

Tergum 9: Arched dorsally; apart from the degree of bend, somewhat wider than long or sometimes as wide as long, and usually more or less narrower posteriorly; with a mid-posterior concavity which is V or U-shaped; some species with a large mid-posterior ventral plane.

Sternum 10: Triangular, semicircular, rectangular, or rather pentagonal probably according to species.

Cercus: Varying in shape with visual angle; elliptic in dorsal or ventral view; [the specific character not studied here carefully].

Remarks on antennal segmentation, some parts of male genitalia, etc.

Except for *V. tibetensis*, the segmentation of the antenna is not clearly seen unless the specimen is washed with alcohol. Not many specimens are thus treated, so individual variation within the same species is not necessarily certain in this character. However, the segmentation of antenna is apparently of specific importance as shown in Key 5, although this character is not repeated in the descriptions of respective species.

It should be noted that the ejaculatory apodeme varies considerably in length and width with the visual angle. Within the aedeagus behind the ejaculatory apodeme, one or two pairs of longitudinal weak lines are seen (may not be visible due to specimen condition) and these lines may represent parts of a membranous tube and may be homologous with endoaedeagal tube defined by Sinclair *et al.* (1994). In *V. tibetensis*, there are paired strong longitudinal sclerites which may be parts of the above endoaedeagal tube. In the illustrations, the lines above are omitted, except for *V. tibetensis*.

The exact understanding of the shape of sternum 10 was not necessarily easy in our study and individual variation within the same species was not tested for this character, although it is probably of specific value in some species.

There is presence or absence of the circle at the apex of the aedeagus in dorsal view. It may vary with visual angle within the same species.

Further, some parts of the male genitalia remain vague in *V. taishanensis* and *V. tibetensis* as mentioned in the description of respective species.

In the illustrations here presented, the antenna and tarsomeres 1-5 are not straightened out completely and the relative lengths of the segments are not exactly shown.

As shown in the key to species, *V. taihangensis*, *V. taishanensis* and *V. yanshanensis* are difficult to determine by the external characters only.

Key 5: *Vermiophis* species based on antenna and coloration of body

1. Antennal flagellum (not macerated in KOH solution) indistinct in segmentation but last joint apparently not shorter than preceding two joints together 2
 - Antennal flagellum (not macerated in KOH solution) quite distinct in segmentation and last joint shorter than preceding two joints together *tibetensis* (♂) (♀ antenna unknown)
- 2(1). Fore and mid coxae dark brown to black (or largely so) (in ♀ *ganquanensis*, mid coxa sometimes yellowish brown as in ♂) 3
 - Fore and mid coxae yellowish brown (or largely so) (or mid coxa sometimes dark brown, as in ♀, in ♂ *ganquanensis* and ♂ *wudangensis*) 7
- 3(2). Apical roughly 1/2 (or more), of hind femur, yellowish (or reddish) brown (rarely this character may not be clear cut in ♀ *ganquanensis*); antennal flagellum (washed with alcohol) 8-segmented 4
 - Hind femur wholly or almost wholly dark brown to black; antennal flagellum (washed with alcohol) 5-6 segmented 5
- 4(3). Abdomen (except segment 1 in both sexes and 6-8 in ♀) largely yellowish (or reddish) brown *ganquanensis* (♀)
 - Abdomen (except anterior part of segments 3-5) deep black *wudangensis* (♀)
- 5(3). [It is uncertain whether or not the couplets 5-6 are significant]; antennal flagellomere 1 or 1-2 (washed with alcohol) with subsegmentation 6
 - Antennal flagellum (washed with alcohol) without subsegmentation *yanshanensis* (♂, ♀)
- 6(5). Antennal flagellum practically 5-segmented *taishanensis* (♂) (♀ unknown)
 - Antennal flagellum 6-segmented *taihangensis* (♂, ♀)
- 7(2). Basal portion of hind femur dark brown to black, as well as whole surfaces of hind tibia and hind coxa; mesonotum black (in *ganquanensis*, side of mesonotum sometimes yellowish brown as in *minshanensis*); antennal flagellum (washed with alcohol) 8-segmented 8
 - Hind femur, hind tibia (except apical portion and often basal lesser 1/2), and often hind coxa yellowish brown; mesonotum (except three black stripes) yellowish brown; antennal flagellum (washed with alcohol) 6-7 segmented *minshanensis* (♂, ♀)
- 8(7). See upper branch of couplet 4 *ganquanensis* (♂)
 - See lower branch of couplet 4 *wudangensis* (♂)

Key 6: *Vermiophis* species based on coloration of abdomen and legs

1. Abdominal segments 2-4, 2-5, or 3-5 dark brown to black and with *distinct* yellowish brown *anterior* band 2
 - Abdominal segments 2-4 or 2-5 largely yellowish brown and without *distinct* dark brown to black *posterior* band 4
- 2(1). Dark part of abdomen not deep black and segment 2 with anterior yellowish brown band 3
 - Abdomen largely deep black and segment 2 wholly so *wudangensis* (♂, ♀)
- 3(2). Hind femur (often except apical dorsal part) yellowish brown *minshanensis* (♂, ♀)

- Hind femur dark brown, but yellowish brown at apex and at middle portion *ibetensis* (♂, ♀)
- 4(1). Hind femur dark brown to black (in *taishanensis* and *yanshanensis*, apical lesser 1/2 may be more or less yellowish brown)
..... *taihangensis* (♂, ♀), *taishanensis* (♀ unknown), and *yanshanensis* (♂, ♀)
- Apical roughly 1/2 (or more) of hind femur yellowish (or reddish) brown (rarely this character may not be clear cut in ♀) *ganquanensis* (♂, ♀)

Key 7: *Vermiophis* species based on male genitalia

1. Tergum 9 with a mid posterior large ventral plane directed anteriorly 2
- Tergum 9 without the above plane 3
- 2(1). Posterior curved part of aedeagus (in lateral view) with apical process directed anteriorly, which is rectangular (somewhat trapezoid) in dorsal view; gonostylus (in dorsal view) narrow; posterolateral ventral part of gonocoxite wide and short; there is no strong paired longitudinal sclerites within aedeagus; mid-posterior ventral plane on tergum 9 triangular and with mid-anterior process directed dorsally; Figs. 153-158 *taihangensis*
- Aedeagus (in lateral view) with apparently three (at least two) dorsal projections, of which the middle is directed posteriorly; gonostylus (in dorsal view) wide; posterolateral ventral part of gonocoxite long and narrow; within aedeagus behind ejaculatory apodeme there are paired longitudinal strong sclerites; mid posterior ventral plane on tergum 9 rectangular and with anterolateral corner protruding; Figs. 169-173 *ibetensis*
- 3(1). Dorsomedial posterior process of gonocoxite rounded apically; aedeagus with two dorsal projections (in lateral view), of which anterior one represents a transverse band in dorsal view 4
- Dorsomedial posterior process of gonocoxite triangular and pointed apically; posterior curved part of aedeagus with anterior process (or knob) near base (in lateral view) 5
- 4(3). Anterior widened part of ejaculatory apodeme elliptical in lateral view; mid-posterior concavity of tergum 9 narrow; Figs. 161-165 *taishanensis*
- Anterior widened part of aedeagus rounded in lateral view; mid-posterior concavity of tergum 9 wide; Figs. 185-189 *yanshanensis*
- 5(3). Dorsomedial posterior projection fused with rest of gonocoxite and not demarcated basally; anterior process on posterior curved part of aedeagus longer and distinct 6
- Dorsomedial posterior projection conspicuous and demarcated from rest of gonocoxite; anterior knob on posterior curved part of aedeagus smaller and indistinct; Figs. 178-182 *wudangensis*
- 6(5). Anterior widened part of ejaculatory apodeme rounded in lateral view; Figs. 129-133
..... *ganquanensis*
- Anterior widened part of ejaculatory apodeme elliptical in lateral view; Figs. 139-14
..... *minshanensis*

Key 8: *Vermiophis* species based on fused gonocoxites and gonostylus

1. Dorsomedial posterior projection (if present) fused with rest of gonocoxite without any

- demarcation 2
- Dorsomedial posterior projection large and triangular, and its basal border distinctly demarcated from rest of gonocoxite; Fig. 181 *wudangensis*
- 2(1). Gonostylus in dorsal view narrow; ventral posterolateral part shorter and wider 3
- Gonostylus in dorsal view wide; ventral posterolateral part longer and narrower; Figs. 172-173 *tibetensis*
- 3(2). Dorsomedial posterior projection smaller and rounded apically or absent (apart from the presence of a process directed ventrally) 4
- Dorsomedial posterior projection larger and pointed apically; Figs. 132, 144
..... *ganquanensis* and *minshanensis*
- 4(3). Dorsomedial posterior projection rounded apically; Figs. 164, 188
..... *taishanensis* and *yanshanensis*
- Dorsomedial posterior projection absent (but there is a process directed ventrally at dorsomedial posterior corner of gonocoxite); Fig. 157 *taihangensis*

Key 9: *Vermiophis* species based on aedeagus and ejaculatory apodeme (in lateral view)

1. Posterior part of aedeagus with one curved broad part directed dorsally 2
- Posterior part of aedeagus bi-lobed (or sometimes possibly tri-lobed) 3
- 2(1). Posterior curved part large and wide, and with a small knob near base of anterior surface; ejaculatory apodeme rather rounded; Figs. 180, 181 *wudangensis*
- Posterior curved part small and narrow, and with an apical process directed anteriorly [which is rectangular in dorsal view]; ejaculatory apodeme elliptical; Figs. 155, 157 *taihangensis*
- 3(1). Posterior curved part with one anterior process near base 4
- Posterior part with two (or three) dorsal projections (of which the anterior one forms a transverse broad band in dorsal view) 5
- 4(3). Ejaculatory apodeme rounded; Fig. 131 *ganquanensis*
- Ejaculatory apodeme elliptical; Fig. 143 *minshanensis*
- 5(3). Two dorsal projections simple in form; [in dorsal view, paired longitudinal sclerites are weak or not seen] 6
- Posterior dorsal projection may be divided into two, of which one would be longer and directed posteriorly, if so; [in dorsal view, paired longitudinal strong sclerites are seen]; Figs. 171, 172 *tibetensis*
- 6(5). Ejaculatory apodeme elliptical; Fig. 163 *taishanensis*
- Ejaculatory apodeme rounded; Fig. 187 *yanshanensis*

Key 10: *Vermiophis* species based on tergum 9

1. Mid-posterior concavity with bottom transverse; tergum 9 with mid-anterior ventral plane 2
- Mid-posterior concavity V-shaped; tergum 9 without any mid-anterior ventral plane 3
- 2(1). Tergum 9 with a mid anterior rectangular ventral plane of which the anterolateral part is protruding; Fig. 170 *tibetensis*

Figs. 125-128. *Vermiophis ganquanensis*, ♂. Antenna and wing. 127-128, showing abnormal wing venation. Refer to p. 130. Original.

- Tergum 9 with a mid-anterior triangular ventral plane; Figs. 153, 154 *taihangensis*
- 3(2). Mid-posterior concavity larger and wider and posterolateral part of tergum 9 rather narrow posteriorly; Figs. 129, 139, 178, 185
.....*ganquanensis*, *minshanensis*, *wudangensis* and *yanshanensis*
- Mid-posterior concavity smaller and narrower and posterolateral part of tergum 9 rather wide at posterior portion; (sternum 10 rather semicircular); Figs. 161, 162 *aishanensis*

5. 1. (31). *Vermiophis ganquanensis* C. Yang (Figs. 125-133, 312, 324, 339-340, 402)

Vermiophis ganquanensis C. Yang, 1979, Entomotaxon., 1: 84. Type locality: China (Shaanxi: Liulimao, Ganquan).

Distribution. China (Shaanxi).

Antennal flagellum 8-segmented. Mesonotum black but sometimes yellowish brown laterally. Hind tibia and basal portion of hind femur dark brown to black. Fore and mid coxae yellowish brown in ♂ but dark brown in ♀, but sometimes ♂ mid coxa dark brown and ♀ mid coxa yellowish brown. Abdomen (except segment 1 and some terminal segments) largely yellowish (or reddish) brown.

This species is almost the same as *V. minshanensis* in the structure of male genitalia but is easily separated from the latter by the external character as shown in couplet 7 of key 5 and couplet 1 of key 6.

Male. Head: Black and with pale gray pollen which is heavy on face. Antenna (except apical portion of flagellum), palpus and proboscis yellowish brown but often antenna and proboscis dark

Figs. 129-133. *Vermiophis ganquanensis*. Male genitalia. 129-130, tergum 9, cerci and sternum 10, dorsal and ventral views; 131-133, hypopygium, lateral, dorsal and ventral views. Original.

brown probably abnormally. Hairs on ocellar tubercle, occiput, cheek, scape, pedicel, palpus and proboscis short and pale (or chiefly so). Width of frons at median ocellus 1.2-1.6 times that just above antenna, 0.4-0.6 times width of one eye on a mid line in direct frontal view, and 1.4-1.7 times width of ocellar triangle, which is 1.0-1.2 times its length; width of frons just above antenna 0.7-1.1 times width of face at lowest part in direct frontal view, and 0.5-0.8 times distance from antenna to median ocellus; distance from proboscis to antenna 2.1-2.5 times that from antenna to median ocellus; antenna 3.2-3.7 times as long as distance from antenna to median ocellus. (N= 9)

Thorax: Shining black in ground color, and with pale gray pollen (which is absent on mesopleuron and area just below wing base); mesonotum with a pair of narrow heavy pale gray pollinose stripes from opposite anterior part of humerus to opposite wing base; sometimes, side of mesonotum before and behind suture, humerus, postalar callus, paired stripes, posterior border of scutellum, and base of postscutellum yellowish brown. Mesonotum, scutellum, propleuron and anterior part of metapleuron with pale hairs which are short on the former two.

Wing: Membrane with a light brown tinge; a small spot from base of marginal cell to base of

discal cell somewhat darker; veins dark brown. Anal cell open; 4th posterior cell petiolate, closed at wing margin and sometimes narrowly open; 1st posterior cell *sometimes* convergent apically or closed at wing margin (rarely petiolate); vein M_4 arising from 2nd basal cell. Halter brown to dark brown, with stem (before narrowest part) yellowish brown, but sometimes halter apparently almost wholly yellowish brown.

Legs: Yellowish brown but tarsomeres 3-5, hind femur (except apical part), hind tibia (except extreme base and apex), hind coxa, hind trochanter, and sometimes tarsomere 2 and mid coxa dark brown to black. Hairs on legs short and chiefly pale. Relative lengths of segments (excluding coxa and trochanter) of fore leg 132 (123-137): 141 (133-148): 100: 36 (28-38): 26 (24-27): 18 (15-21): 21 (18-24), of mid leg 154 (150-158): 169 (161-179): 94 (91-100): 32 (30-35): 22 (19-24): 15 (14-17): 18 (17-18), of hind leg 238 (235-244): 267 (262-272): 132 (124-140): 45 (41-53): 28 (24-31): 18 (16-19): 21 (19-24) and in hind leg from the side, relative widths of femur, tibia and tarsomeres 1-3, 30 (27-31): 23 (21-27): 14 (13-14): 9 (8-10): 8 (7-9); (N= 7).

Abdomen: Yellowish brown; segments 1 and 8, anterior part of tergum 2 dark brown to black, and segments 6-7 largely so. Hairs on abdomen short.

Tergum 9 (in dorsal or ventral view): Posterolateral protruded part bluntly pointed; mid-posterior concavity wide, U-shaped, and with a mid-anterior projection directed dorsally.

Sternum 10: Apparently triangular.

Fused gonocoxites: Dorsomedial posterior projection rather large and tapering apically; posterolateral ventral part wide and rather trapezoid.

Aedeagus: Posterior curved part with an anterior knob near base (in lateral view).

Ejaculatory apodeme: Anterior widened part rounded (in lateral view).

Length: Body 9.6-13.1 mm; wing 7.4-9.3 mm; fore basitarsus 1.5-1.9 mm; (N= 7).

Female. Similar to male except as follows: Head: Width of frons at median ocellus 0.8-1.4 times that just above antenna and 1.3-2.0 times width of ocellar triangle which is 0.8-1.4 times its length; width of frons just above antenna 0.7-0.9 times distance from antenna to median ocellus; distance from proboscis to antenna 2.2-2.9 times that from antenna to median ocellus; antenna 3.4-4.4 times as long as distance from antenna to median ocellus; (N= 12); [perhaps no significant difference is present in the above values between both sexes].

Wing: Vein M_4 *rarely* arising from discal cell (this may be so in ♂).

Legs: Fore and mid coxae brown to dark brown; mid coxa sometimes yellowish brown as in ♂; yellowish brown apical portion of hind femur rarely indistinct. Relative lengths of segments of fore leg 140 (133-147): 155 (150-160): 100: 30 (27-32): 19 (15-22): 12 (10-13): 17 (14-18), of mid leg 167 (160-176): 181 (176-188): 95 (89-100): 28 (26-30): 17 (15-19): 11 (9-12): 15 (13-16), of hind leg 259 (250-271): 287 (277-290): 139 (130-147): 42 (37-47): 24 (19-27): 16 (14-18): 20 (18-22) and in hind leg from the side, relative widths of femur, tibia and tarsomeres 1-3, 32 (29-35): 23 (20-26): 15 (13-18): 10 (9-12): 8 (7-9); [N= 10].

Abdomen: Segments 1, 6-8 and posterior part of segment 5 (or 4-5) dark brown to black; rest of abdomen largely yellowish brown.

Length: Body 10.1-15.9 mm; wing 8.6-11.0 mm; fore basitarsus 1.5-1.9 mm; (N= 17).

Specimens examined: 5 ♂♂, 16 ♀♀ (holotype and paratypes), Shaanxi (Ganquan: Liulimao), C. Yang leg (the larvae were collected during vi-ix, 1971 and the adults emerged during 9. iv.-7.v., 1972

Figs. 134-138. *Vermiphis minshanensis*. Antenna, wing and abdomen. 134-136, 138, ♂; 137, ♀. 136, 137, showing abnormal wing venation; 138, dorsal view. Original.

and 14-27. iv. 1973 respectively); 2 ♂♂, 1 ♀, Shaanxi (Ganquan), C. Yang leg (emerged during v-vi, 1981).

5. 2. (32). *Vermiphis minshanensis* C. Yang & Chen (Figs. 134-145, 313, 324)

Vermiphis minshanensis C. Yang & Chen, 1993, Entomotaxon., 15: 131. Type locality: China (Gansu: Wenxian County).

Distribution. China (Gansu).

Antennal flagellum 6-7 segmented. Mesonotum yellowish brown, with 3 broad black stripes ending before scutellum and far distant from humeri. Fore and mid coxae, hind tibia (except apical portion), hind femur and often hind coxa yellowish brown. Abdominal segments 2-5 banded with dark brown at posterior part.

This species is very similar to *V. ganquanensis* in the structure of male genitalia but they are easily separated from each other by the external characters (see keys 5 and 6).

Male. Head: Black and with pale gray pollen which is heavy on face. Antenna (except flagellum minus flagellomere 1 or 1-3), palpus and proboscis yellowish brown. [Palpus and proboscis are sometimes dark brown but this may be due to bad specimen condition]. Ocellar tubercle, occiput, cheek, scape, pedicel, palpus and proboscis with pale hairs which are short on scape and pedicel. Width of frons at median ocellus 1.1-1.5 times that just above antenna, 0.4-0.6 times width of one eye on a mid line in direct frontal view, and 1.4-1.9 times width of ocellar triangle which is 0.9-1.1 times its length; width of frons just above antenna 0.8-1.0 times width of face at lowest part in direct frontal view and 0.5-0.9 times distance from antenna to median ocellus; distance from proboscis to antenna 1.8-2.7 times that from antenna to median ocellus; antenna 3.1-4.1 times distance from antenna to median ocellus; [N= 8].

Thorax: Yellowish brown and pale gray pollinose; mesonotum with three broad dark brown to black stripes ending before scutellum, of which lateral stripes far distant from humeri; mesopleuron shining black, and rest of pleuron, scutellum and postscutellum (except base and sides) with a brown to dark brown tinge which may be weak in metapleuron. Mesonotum, scutellum, propleuron and anterior part of metapleuron with pale hairs which are short on the former two.

Wing: Membrane lightly tinged with brown; a somewhat darker spot present from base of marginal cell to base of discal cell; veins dark brown. Anal cell open (with mouth usually narrow); 4th posterior cell petiolate; 2nd posterior cell *rarely* petiolate at base; vein M₄ arising from 2nd basal cell but sometimes from discal cell at the base of m-cu crossvein. Halter yellowish brown [sometimes apex of knob darkened probably abnormally].

Legs: Yellowish brown; tarsomeres 2-5 and apical half of hind tibia dark brown; hind coxa, apical dorsal surface of hind femur and basal lesser 1/2 of hind tibia often with a brown to dark brown tinge. Hairs on legs short, and pale (or chiefly so). Relative lengths of segments (excluding coxa and trochanter) of fore leg 133 (125-144): 144 (142-148): 100: 35 (33-36): 27 (25-29): 17 (16-18): 20 (17-21), of mid leg 147 (143-156): 172 (164-188): 96 (92-100): 35 (33-36): 23 (21-25): 14 (13-16): 18 (17-18), of hind leg 234 (229-240): 272 (261-280): 129 (125-132): 45 (42-46): 30 (25-32): 19 (17-21): 21 (19-22) and in hind leg from the side, relative widths of femur, tibia and tarsomeres 1-3, 29 (25-32): 23 (21-26): 15 (14-17): 12 (11-13): 8 (7-8); [N= 4].

Abdomen: Yellowish brown; segment 1 (sometimes except tergum 1), posterior part of terga 2-5, and sometimes segments 6-8 dark brown; hairs on abdomen short, and pale (or chiefly so).

Tergum 9 (in dorsal or ventral view): Posterolateral protruded part bluntly pointed; mid-posterior concavity wide, V-shaped, and with a mid-anterior projection directed dorsally.

Sternum 10: [The shape remained unknown in our study].

Fused gonocoxites: Dorsal posterior part tapering apically as a whole and pointed; posterolateral ventral part wide and rather trapezoid.

Aedeagus: Posterior curved part with an anterior elongate process near base (in lateral view).

Ejaculatory apodeme: Anterior widened part elliptic (in lateral view).

Length: Body 7.7-9.7 mm (N= 6); wing 6.0-7.7 mm (N= 6); fore basitarsus 1.2-1.4 mm (N= 4).

Female. Similar to male except as follows: Head: No difference was found in the measured values between both sexes [N= 6 in ♀].

Legs: Relative lengths of segments of fore leg 145 (139-150): 165 (161-172): 100: 31 (29-32): 19 (18-20): 12 (11-14): 17 (16-18), of mid leg 168 (161-172): 191 (186-196): 102 (96-107): 31 (29-32):

Figs. 139-145. *Vermiphis minshanensis*. Male genitalia. 139-142, tergum 9 and cerci (sternum 10 is omitted), dorsal, ventral, lateral and posterior views; 143-145, hypopygium, lateral, dorsal and ventral views. Original.

18 (14-20): 11 (11-12): 16 (14-18), of hind leg 261 (250-271): 300 (293-307): 143 (139-146): 45 (43-50): 29 (29): 17 (14-18): 21 (20-21) and in hind leg from the side, relative widths of femur, tibia, and tarsomeres 1-3, 35 (32-36): 26 (25-29): 18 (18): 10 (9-11): 8 (7-9); [N= 4].

Wing: Fourth posterior cell open (this may be so in ♂); 2nd submarginal cell *rarely* petiolate at base (this may be so in ♂).

Length: Body 9.7-10.9 mm (N= 3); wing 7.3-8.4 mm (N= 5); fore basitarsus 1.3-1.4 mm (N= 4).

Specimens examined: 8♂♂, 6♀♀ (holotype and paratypes), Wenxian County, Gansu, C. Yang & S. Zhang leg (the larvae were collected and the adults emerged on 28. iv. - 12. v., 1981-1986).

5. 3. (33). *Vermiphis taihangensis* C. Yang & Chen (Figs. 146-158, 314, 325)

Vermiphis taihangensis C. Yang & Chen, 1993, Entomotaxon., 15: 129. Type locality: China (Beijing: Shangfangshan [500 m]).

Distribution. China (Beijing; Shanxi).

Antennal flagellum is 6-segmented, but the flagellomeres 1-2 each may have an indistinct division. Mesonotum black and with a pair of pale gray pollinose stripes. Coxa, hind femur and hind tibia dark brown to black. Abdominal segments 2-5 largely yellowish brown and segments 2-4 (or 2-5) without

Figs. 146-152. *Vermiophis taihangensis*. 146, 150, 151, ♀; 147-149, 152, ♂. 146, head, anterior view; 147, palpus; 148, antenna; 149, wing; 150, hind tarsus, lateral view; 151, apex of hind tarsus, ventral view; 152, abdomen, dorsal view. Original.

posterior dark brown to black band.

This species is similar to *V. taishanensis* and *V. yanshanensis* but the male genitalia are different markedly from them.

Male. Head: Black and with pale gray pollen which is heavy on face. Antenna, palpus and proboscis brown to dark brown but basal portion of flagellum and pedicel usually yellowish brown. Sometimes palpus and proboscis apparently yellowish brown. Ocellar tubercle, occiput, cheek, scape, pedicel, palpus and proboscis with pale hairs. Width of frons at median ocellus 1.3-1.7 times that just above antenna, 0.4-0.5 times width of one eye on a mid line in direct frontal view, and 1.3-1.9 times width of ocellar triangle which is 0.9-1.3 times its length; width of frons just above antenna 0.8-1.0 times width of face at lowest part in direct frontal view and 0.5-0.7 times distance from antenna to median ocellus; distance from proboscis to antenna 1.8-2.4 times that from antenna to median ocellus; antenna 2.5-3.5 times distance from antenna to median ocellus; [N= 12].

Thorax: Shining black in ground color and with pale gray pollen; mesonotum with a pair of stronger pale gray pollinose stripes starting from or near humerus and ending before scutellum. Propleuron, anterior part of metapleuron, mesonotum and scutellum with pale hairs which are shorter on the latter two.

Wing: Membrane with a light brown tinge; 1st basal cell may have a somewhat darker small subapical spot; veins dark brown. Anal cell open or *rarely* closed at wing margin; 4th posterior cell petiolate, closed at wing margin or *rarely* open; vein M_4 arising from 2nd basal cell but sometimes from discal cell at the base of m-cu crossvein. Halter dark brown, but the basal part before narrowest part yellowish brown.

Legs: Yellowish brown, but coxa, tarsomeres 2-5 or 3-5, in hind leg, trochanter, femur and tibia dark brown; fore and mid coxae sometimes partly yellowish brown in certain lights. Hairs on legs short and pale but those on hind tibia apparently chiefly black. Relative lengths of segments (excluding coxa and trochanter) of fore leg 144 (137-150): 163 (152-171): 100: 36 (33-38): 27 (25-29): 20 (17-21): 22 (21-24), of mid leg 166 (158-172): 193 (181-204): 95 (86-104): 34 (29-39): 24 (21-29): 17 (13-21): 20 (19-21), of hind leg 265 (248-279): 297 (278-308): 130 (119-146): 46 (42-50): 30 (25-36): 20 (17-23): 23 (21-27) and in hind leg from the side, relative widths of femur, tibia and tarsomeres 1-3, 32 (28-38): 26 (21-33): 18 (14-21): 11 (10-13): 9 (7-12); [N= 10].

Abdomen: Yellowish brown; segment 1, 6-8, genitalia and often side of tergum 2 and apical portion of segment 5 dark brown to black; sometimes terga 1 and 6 partly yellowish brown. Hairs on abdomen chiefly black.

Tergum 9 (in dorsal or ventral view): Posterolateral protruded part wide and rounded; mid-posterior concavity wide with the bottom transverse; mid-posterior margin with a large triangular ventral plane having a small mid-anterior projection directed dorsally.

Sternum 10: Rectangular (or possibly rather trapezoid) and wider than long.

Fused gonocoxites: Dorsomedial posterior projection absent but a small elongate process (directed ventrally) present at posterior medial corner; dorsomedial margin folded ventrally; posterolateral ventral part wide and with a small knob protruding posteriorly; dorsal bridge with posterior margin deeply concave.

Aedeagus: Posterior curved part with apical process directed anteriorly which is rectangular or trapezoid in dorsal view.

Ejaculatory apodeme: Anterior widened part elliptical (in lateral view).

Length: Body 7.9-11.0 mm (N= 12); wing 6.0-8.6 mm (N= 14); fore basitarsus 1.0-1.5 mm (N= 14).

Female. No significant difference is found between sexes, besides genitalia. Head: Width of frons at median ocellus 1.2-1.8 times that just above antenna, 0.5-0.6 times width of one eye on a mid line, and 1.4-1.8 times width of ocellar triangle which is 1.1-1.4 times its length; width of frons just above antenna 0.6-1.0 times width of one eye on a mid line in direct frontal view and 0.5-0.8 times distance from antenna to median ocellus; distance from proboscis to antenna 2.0-2.5 times that from antenna to median ocellus; antenna 2.8-3.6 times distance from antenna to median ocellus; [N= 12].

Legs: Relative lengths of segments of fore leg 153 (144-161): 174 (167-184): 100: 33 (31-36): 22 (19-24): 14 (12-19): 20 (17-24), of mid leg 180 (169-193): 207 (193-221): 94 (84-100): 28 (23-32): 18 (16-21): 11 (9-14): 17 (16-20), of hind leg 284 (263-314): 323 (300-357): 135 (119-143): 40 (38-50): 27 (23-31): 16 (12-19): 20 (17-23) and in hind leg from the side, relative widths of femur, tibia and

Figs. 153-158. *Vermiophis taihangensis*. Male genitalia. 153-154, tergum 9, cerci and sternum 10, dorsal and ventral views; 155-156, lateral and posterior views; 157-158, hypopygium, dorsal and ventral views. Original.

tarsomeres 1-3, 34 (31-36): 26 (23-29): 18 (16-20): 11 (9-12): 9 (8-10); [N= 10].

Length: Body 8.6-12.1 mm (N= 14); wing 7.9-9.7 mm (N= 15); fore basitarsus 1.1-1.6 mm (N= 16).

Specimens examined: 18♂♂, 30♀♀ (holotype and paratypes). The larvae were taken at Beijing (Shangfangshan [500 m]; Miaofengshan [1000 m]; Yanqing County) and Shanxi Province (Wutaishan [1720 m]; Yungang) by C. Yang and the adults emerged during April and May, 1974-1983.

5. 4. (34). *Vermiophis taishanensis* C. Yang & Chen (Figs. 159-165, 326)

Vermiophis taishanensis C. Yang & Chen, 1993, Entomotaxon., 15: 130. Type locality: China (Shandong Province: Taishan: Shanziya [600 m]).

Distribution. China (Shandong).

This species is most closely related to *V. yanshanensis*, but may be separated from the latter in the details of male genitalia. It is also similar to *V. taihangensis* in external characters, although the segmentation of antennal flagellum may possibly be different from that of the latter. The male genitalia are different markedly between *V. taishanensis* and *V. taihangensis*.

Fitting description of *V. taihangensis* except as follows: Head: Palpal segment 2 may be whitish in

Figs. 159-160. *Vermiophis taishanensis*, ♂. Antenna and wing. Original.

certain lights (as in *V. yanshanensis*). Width of frons at median ocellus 1.5 times that just above antenna, 0.4 times width of one eye on a mid line in direct frontal view, and 1.3 times width of ocellar triangle which is 1.2 times its length; width of frons just above antenna 0.5 times width of face at lowest part in direct frontal view and 0.4 times distance from antenna to median ocellus; distance from proboscis to antenna 1.5 times that from antenna to median ocellus; antenna 2.0 times distance from antenna to median ocellus; [N= 1].

Wing: [In 1 specimen (holotype) examined], 4th posterior cell open; anal cell open; vein M₄ arising from discal cell at base of m-cu crossvein (on right wing) and from 2nd basal cell (on left wing); vein M₂ ending just before wing margin (on right wing) and ending at wing margin (on left wing). Over basal 1/2 of halter may be yellowish brown (as in *V. yanshanensis*).

Legs: Apical lesser 1/2 of hind femur is indistinctly yellowish brown (this is sometimes so in *V. yanshanensis*) and may be abnormal in condition. Relative lengths of segments (excluding coxa and trochanter) of fore leg 150: 173: 100: 38: 31: 19: 27, of mid leg 173: 196: 100: 38: 23: 19: 23, of hind leg 265: 277: 127: 42: 31: 19: 27 and in hind leg from the side, relative widths of femur, tibia and tarsomeres 1-3, 38: 27: 19: 13: 13; [N= 1].

Tergum 9 (in dorsal or ventral view): Posterolateral protruded part rather wide; mid-posterior concavity V-shaped, and rather narrow [presence or absence of median projection directed dorsally at bottom of concavity was not confirmed].

Sternum 10: Triangular or semicircular.

Fused gonocoxites: Dorsomedial posterior projection moderate in size and rather rounded apically; posterolateral ventral part trapezoid and with gentle convexity at posterolateral corner; [anterior margin of dorsal bridge was not seen in the specimen examined, so its mid-length remains unknown].

Aedeagus: Posterior part with 2 dorsal projections, of which the anterior one represents a transverse band (in dorsal view) and is Y-shaped (when seen in three dimensions).

Ejaculatory apodeme: Anterior widened part elliptical (in lateral view).

Figs. 161-165. *Vermiphis taishanensis*. Male genitalia. 161-162, tergum 9, cerci and sternum 10, dorsal and ventral views; 163-165, hypopygium, lateral, dorsal and ventral views. Original.

Length: Body 11.8 mm; wing 9.1 mm; fore basitarsus 1.2 mm; (N=1).

Female. Unknown.

Specimen examined: 1♂ (holotype), Shanziya (600 m), Taishan, Shandong Prov., T. Mo leg. From the larva collected, the adult emerged on 7. v. 1986.

5. 5. (35). *Vermiphis tibetensis* C. Yang & Chen (Figs. 166-173, 325)

Vermiphis tibetensis C. Yang & Chen, 1987, Agricultural Insects, Spiders, Plant Diseases and Weeds of Xizang, 2: 157. Type locality: China (Tibet: Bomi Co.: Gyaide).

Distribution. China (Tibet: Bomi Co.; Nyingchi Co.).

This species is characterized by having last segment of antennal flagellum not longer than preceding two segments together (at least in ♂). Mesonotum yellowish brown and with 3 broad black

stripes. Legs yellowish brown, but tarsomere 5, hind coxa, hind tibia, and hind femur at basal and subapical parts brown to dark brown. Abdominal segments 2-6 distinctly or indistinctly banded with yellowish brown at anterior part. The male genitalia of *V. tibetensis* differ markedly from those of other species.

Male. Head: Black and pale gray pollinose; antenna dark brown (except pedicel and base of flagellum) or yellowish brown (except dorsal surface and last two flagellomeres); palpus and proboscis yellowish brown. Ocellar tubercle, occiput, cheek, palpus and proboscis with pale hairs which are short on the ocellar tubercle. Scape and pedicel with short inconspicuous hairs which are pale and black. Width of frons at median ocellus 1.1 times that just above antenna, 0.4 times width of one eye on a mid line in direct frontal view, and 1.1-1.2 times width of ocellar triangle which is 1.1-1.4 times its length; width of frons just above antenna 0.6-1.0 times width of face at lowest part in direct frontal view and 0.9-1.0 times distance from antenna to median ocellus; distance from proboscis to antenna 3.4-3.8 times that from antenna to median ocellus; antenna 5.7-6.5 times distance from antenna to median ocellus; [N=4].

Thorax: Shining black in ground color and with pale gray pollen; mesonotum yellowish brown and with three broad shining black stripes ending before scutellum and distant from humeri. propleuron, anterior part of metapleuron, mesonotum and scutellum with pale hairs which are short on the latter two.

Wing: Membrane tinged with light brown; 1st basal cell with a somewhat darker subapical spot; veins dark brown. Anal cell open, 4th posterior cell open or petiolate; vein M₄ arising from discal cell at base of m-cu crossvein but often from 2nd basal cell. Halter brown to dark brown, but base (before narrowest part) yellowish brown.

Legs: Yellowish brown; tarsomere 5, in hind leg, coxa, trochanter, femur (except middle part and apical portion) and tibia (except base and apex) brown to dark brown. Hairs on legs short and pale but those on hind tibia and posterior surface of hind femur chiefly black. Relative lengths of segments (excluding coxa and trochanter) of fore leg 131 (127-134): 153 (147-159): 100: 37 (34-41): 24 (24-25): 17 (16-17): 19 (18-21), of mid leg 153 (147-159): 178 (172-183): 97 (94-100): 33 (31-34): 20 (19-21): 14 (14): 17 (15-19), of hind leg 251 (242-266): 266 (258-276): 137 (133-141): 41 (38-45): 25 (22-28): 16 (14-17): 19 (18-21) and in hind leg from the side, relative widths of femur, tibia and tarsomeres 1-3, 31 (30-31): 24 (24-25): 15 (14-17): 11 (9-14): 9 (9-10); [N=3].

Abdomen: Dark brown and with anterior part of segments 2-6 yellowish brown, or yellowish brown and with a brown to dark brown tinge in several parts whose boundaries are not well marked (i. e. in general, segments 2-6 banded with yellowish brown anteriorly). Hairs on abdomen chiefly black.

Tergum 9 (in dorsal or ventral view): A square in shape; posterolateral protruded part wide, rather rounded apically and with a transverse anterior black line which disappears in some visual angles; mid-posterior concavity wide and with bottom transverse; mid-posterior margin with a mid-anterior ventral plane which is wider than long, rectangular and has the lateral corner produced anteroventrally.

Sternum 10: Rather semicircular; sternum 10 (plus cercus) situated before ventral plane.

Fused gonocoxites: Dorsal piece relatively short; dorsomedial posterior projection indistinct, small and rounded apically; posterolateral ventral part relatively long, narrow and rather trapezoid.

Aedeagus: Posterior dorsal projection directed posteriorly as a whole in lateral view; anterior dorsal projection represents a transverse band in dorsal view; paired longitudinal strong sclerites

Fig. 166. *Vermiothis tibetensis*, ♂. Slightly modified from C. Yang & Chen (1987).

167

Figs. 167-168. *Vermiothis tibetensis*, ♂. Antenna and wing. Original.

168

present inside aedeagus behind ejaculatory apodeme; [it remains unknown whether or not posterior dorsal projection is divided into two].

Ejaculatory apodeme: Anterior widened part elliptical (in lateral view).

Gonostylus: Wide and with posteromedial corner pointed (in dorsal view).

Length: Body 11.1-11.7 mm (N= 2); wing 8.7-9.1 mm (N= 3); fore basitarsus 1.5-1.7 mm (N= 3).

Female. Apparently no significant difference is found between both sexes (apart from genitalia).

Figs. 169-173. *Vermiothis tibetensis*. Male genitalia. 169-170. tergum 9, cerci and sternum 10, dorsal and ventral views; 171-173, hypopygium, lateral, dorsal and ventral views. Original.

Head (antennal flagellum broken off): Proboscis dark brown (this may be sometimes so in ♂). Width of frons at median ocellus 1.2 times that just above antenna, 0.6 times width of one eye on a mid line in direct frontal view, and 1.4 times width of ocellar triangle which is 1.4 times its length; width of frons just above antenna 0.8 times width of face at lowest part in direct frontal view and 1.1 times distance from antenna to median ocellus; distance from proboscis to antenna 3.6 times that from antenna to median ocellus; [N= 1].

Legs (hind legs broken off): Relative lengths of segments of fore leg 153: 163: 100: 33: 23: 13: 20, of mid leg 177: 200: 100: 33: 20: 13: 20, of hind leg ?; [N= 1].

Length: Body 10.6 mm; wing 9.6 mm; fore basitarsus 1.5 mm; (N= 1).

Specimens examined: Holotype ♂, paratype 1 ♂, Gyaide, Bomi Co., Tibet, 14. vii. 1978, F. Li; paratypes 2 ♂ ♂, Zhamog, Bomi Co., Tibet, 21. vii. 1978, F. Li; paratype 1 ♀, Nyingchi Co., Tibet, 1977, F. Wang.

Fig. 174. *Vermiothis wudangensis* ♂. From C. Yang & Chen (1986).

5. 6. (36). *Vermiothis wudangensis* C. Yang & Chen (Figs. 174-182, 310-311, 324)

Vermiothis wudangensis C. Yang & Chen, 1986, Journal of Huazhong Agricultural University, 5 (4): 321. Type locality: China (Hubei Prov.: Wudangshan: Jinxiandong [900 m]).

Distribution. China (Hubei).

Antennal flagellum 8-segmented. Mesonotum black and with a pair of pale gray pollinose stripes. Apical, approximately 1/2 (or more) of hind femur yellowish brown. Hind tibia dark brown to black. Fore and mid coxae yellowish brown in ♂ and dark brown in ♀ (sometimes ♂ mid coxa dark brown). Abdomen deep black, but segments 3-5 banded with yellowish brown anteriorly.

The male genitalia of *V. wudangensis* are similar to those of *V. ganquanensis* and *V. minshanensis* but are easily separated from them as shown in Key 7 (couplet 5).

Male. Head: Black and with pale gray pollen; antenna dark brown to black, but base of flagellum and usually pedicel yellowish brown; palpus and proboscis yellowish brown. Occiput, cheek, palpus, proboscis, scape, pedicel and ocellar tubercle with pale hairs which are short on the latter three. Width of frons at median ocellus 1.2-1.8 times that just above antenna, 0.4-0.5 times width of one eye on a mid line in direct frontal view, and 1.3-1.7 times width of ocellar triangle which is 0.9-1.2 times its length; width of frons just above antenna 0.6-1.2 times width of face at lowest part in direct frontal view and 0.4-0.7 times distance from antenna to median ocellus; distance from proboscis to antenna 1.9-2.7 times that from antenna to median ocellus; antenna 3.1-3.8 times distance from antenna to median ocellus; [N= 8].

Thorax: Shining black in ground color and with pale gray pollen; mesonotum with a pair of pale gray pollinose stripes which are distinct from humeri and may extend to scutellum. Propleuron and anterior part of metapleuron, mesonotum and scutellum with pale hairs which are short on the latter two.

Wing: Membrane with a light brown tinge; 1st basal cell with a small darker subapical spot; veins dark brown. Fourth posterior cell open, closed at wing margin or rarely petiolate; anal cell open; vein M₄ arising from discal cell at base of m-cu crossvein, but rarely from 2nd basal cell. Halter wholly

Figs. 175-177. *Vermiothis wudangensis*. 175-176, ♂; 177, ♀ abdomen, dorsal view. Original.

yellowish brown.

Legs: Yellowish brown; tarsomeres 2-5, in hind leg, coxa, trochanter, tibia (except extreme base), basal roughly 1/2 (or less) of femur dark brown; fore and mid tibiae, and base of fore and mid coxae often brown to dark brown; mid coxa sometimes wholly dark brown. Hairs on legs short and pale but those of hind femur and tibia apparently partly black. Relative lengths of segments (excluding coxa and trochanter) of fore leg 123 (117-130): 138 (129-145): 100: 36 (34-39): 28 (25-30): 19 (17-20): 21 (20-23), of mid leg 143 (137-149): 163 (156-168): 93 (90-94): 32 (30-35): 24 (22-26): 14 (13-16): 18 (15-19), of hind leg 223 (220-226): 252 (230-268): 132 (126-137): 44 (43-46): [29]: [20]: [20] and in hind leg from the side, relative widths of femur, tibia and tarsomeres 1-3, 27 (26-29): 20 (18-22): 13 (12-14): 9 (8-10): [7]; [N= 7, sometimes 5 or 1 according to segment].

Abdomen: Black; anterior part of segments 3-5 yellowish brown and genitalia partly so. Hairs on abdomen chiefly black.

Tergum 9 (in dorsal or ventral view): Posterolateral protruded part rather narrow and bluntly pointed; mid-posterior concavity wide, V-shaped, and with mid-anterior projection directed dorsally.

Sternum 10: Apparently rather pentagonal in shape.

Fused gonocoxites: Dorsomedial posterior projection conspicuous and triangular; another dorsomedial triangular convexity present around middle of gonocoxite (this convexity may not be visible from some angles); posterolateral ventral part wide and waved at apical margin; dorsal bridge longitudinally thin.

Aedeagus: Posterior curved part wide and with an anterior gentle convexity near base (in lateral view).

Ejaculatory apodeme: Anterior widened part rather rounded in lateral view.

Figs. 178-182. *Vermiophis wudangensis*. Male genitalia. 178-179, tergum 9, cerci and sternum 10, dorsal and ventral views; 180-182, hypopygium, lateral, dorsal and ventral views. Original.

Length: Body 10.1-13.1 mm; wing 7.6-10.0 mm; fore basitarsus 1.6-2.1 mm; (N= 7).

Female. Fitting description of ♂ except as follows: Head: Width of frons at median ocellus 1.1-1.3 times that just above antenna which is 0.7-0.8 times distance from antenna to median ocellus; distance from proboscis to antenna 2.6-3.1 times that from antenna to median ocellus; antenna 4.3-4.4 times distance from antenna to median ocellus; [N= 4].

Wing: Vein M₄ sometimes arising from discal cell apart from m-cu crossvein (this may be so in ♂).

Legs: Fore and mid coxae dark brown. Relative lengths of segments of fore leg 132 (123-138): 148 (144-150): 100: 31 (28-33): 21 (19-23): 14 (13-15): 17 (15-19), of mid leg 157 (153-160): 173 (167-179): 95 (90-100): 27 (26-28): 17 (15-18): 10 (10): 13 (13), of hind leg 244 (228-258): 276 (264-285): 135 (128-140): 40 (38-41): 27 (26-28): 15 (15): 20 (18-21) and in hind leg from the side, relative widths of femur, tibia and tarsomeres 1-3, 30 (28-32): 23 (23-24): 14 (13-15): 8 (8-9): 6 (5-7); [N= 3 or sometimes 2 according to segment].

Length: Body 9.7-12.4 mm; wing 9.6-10.7 mm; fore basitarsus 1.7-2.0 mm; (N= 3).

Specimens examined (all from Wudangshan): 5 ♂♂, 2 ♀♀ (holotype and paratypes),

Jinxiandong (900 m), 29. v. - 1. vi. 1984, C. Yang; 2 ♂♂ (paratypes), 1100 m, 2. vi. 1984, C. Yang; 1 ♀, Taizidong, 2. vi. 1984, X. Wang.

5. 7. (37). *Vermiophis yanshanensis* C. Yang & Chen (Figs. 183-189, 315, 326, 404)

Vermiophis yanshanensis C. Yang & Chen, 1993, Entomotaxon., 15 (2): 127. Type locality: China (Hebei Prov.: Wulingshan [700 m]).

Distribution. China (Hebei).

This species is very similar to *V. taihangensis* and *V. taishanensis* in external characters but may be separated from them by having antennal flagellomeres 1-2 without any subsegmentation. The male genitalia of *V. yanshanensis* are very similar to those of *V. taishanensis* but may be different as shown in the couplet 4 of key 5.

Male. Similar to *V. taihangensis* except as follows: Head: Palpal segment 2 whitish in certain lights in better preserved specimens. Width of frons at median ocellus 1. 2-1. 4 times that just above antenna, 0. 5-0. 6 times width of one eye on a mid line in direct frontal view, and 1.4-1.8 times width of ocellar triangle which is 1. 0-1. 2 times its length; width of frons just above antenna 0. 8-1. 0 times width of face at lowest part in direct frontal view and 0. 5-0. 8 times distance from antenna to median ocellus; distance from proboscis to antenna 1.8-2.4 times that from antenna to median ocellus; antenna 2.8-3.4 times distance from proboscis to antenna; [N= 12]; the above values apparently not different significantly from those of *V. taihangensis*.

Thorax: In halter beyond narrowest part, about half may be rather yellowish brown basally or paler than apical portion.

Wing: Fourth posterior cell closed at wing margin; sometimes petiolate or open; anal cell open; vein M₄ arising from 2nd basal cell or from discal cell at base of m-cu crossvein.

Legs: In fore and mid legs, tibia and tarsomeres 1-2 sometimes brown to dark brown rather than yellowish brown (this may be sometimes so in *V. taihangensis*). Relative lengths of segments (excluding coxa and trochanter) of fore leg 142 (131-150): 160 (148-167): 100: 36 (31-38): 28 (24-30): 19 (17-23): 23 (21-26), of mid leg 160 (148-167): 186 (178-193): 95 (90-100): 31 (28-33): 23 (21-27): 15 (10-20): 20 (17-23), of hind leg 250 (238-267): 282 (266-293): 133 (121-141): 40 (38-44): 28 (23-33): 19 (17-21): 22 (19-24) and in hind leg from the side, relative widths of femur, tibia and tarsomeres 1-3, 30 (28-31): 23 (19-26): 16 (14-17): 10 (9-11): 8 (7-11); [N= 10].

Abdomen: In 1 ♂ (of 13 ♂♂ examined), abdomen largely dark brown, apparently abnormally.

Tergum 9 (in dorsal or ventral view): Posterolateral protruded part rather bluntly pointed; mid-posterior concavity wide, V-shaped, and with mid-anterior projection directed dorsally.

Sternum 10: Apparently rather pentagonal.

Fused gonocoxites: Dorsomedial posterior projection moderate in size and rounded at apical margin; posterolateral ventral part widened at basal part and then gradually narrower toward posterolateral apex; dorsal bridge longitudinally very thin.

Aedeagus: Posterior part with two dorsal projections, of which anterior one represents a transverse U-shaped band in dorsal view and is Y-shaped when seen three dimensionally.

Ejaculatory apodeme: Anterior widened part rounded in lateral view.

Length: Body 9.3-11.4 mm (N= 15); wing 7.4-9.0 mm (N= 14); fore basitarsus 1.4-1.6 mm (N= 14).

Female. No difference is found between sexes, besides genitalia. Head: Width of frons at median

Figs. 183-184. *Vermiphipis yanshanensis*, ♂.
Antenna and wing. Original.

Figs. 185-189. *Vermiphipis yanshanensis*. Male genitalia.
185-186, tergum 9, cerci and sternum 10, dorsal and ventral views; 187-189, hypopygium, lateral, dorsal and ventral views. Original.

ocellus 1.2-1.6 times that just above antenna, 0.5-0.6 times width of one eye on a mid line in direct frontal view, and 1.5-2.2 times width of ocellar triangle which is 1.0-1.3 times its length; width of frons just above antenna 0.7-1.0 times width of face at lowest part in direct frontal view and 0.6-0.8 times distance from antenna to median ocellus; distance from proboscis to antenna 2.0-2.5 times that from antenna to median ocellus; antenna 3.1-3.7 times distance from antenna to median ocellus; [N=12]; the above values are apparently not significantly different from those in ♂ *yanshanensis* and ♀ *taihangensis*.

Wing: Vein M₄ arising from 2nd basal cell but rarely from discal cell apart from m-cu crossvein.

Legs: Relative lengths of segments of fore leg 152 (145-167): 169 (161-177): 100: 32 (31-33): 22 (21-24): 15 (13-17): 19 (17-20), of mid leg 175 (166-193): 196 (186-210): 95 (89-100): 28 (24-30): 18 (16-20): 11 (9-13): 17 (16-19), of hind leg 270 (255-293): 303 (281-322): 135 (128-141): 37 (34-39): 26 (21-31): 16 (14-17): 20 (17-23) and in hind leg from the side, relative widths of femur, tibia and tarsomeres 1-3, 33 (29-37): 24 (23-27): 18 (16-20): 11 (9-13): 8 (6-10); [N=10].

Length: Body 8.4-12.3 mm (N=17); wing 8.3-10.1 mm (N=16); fore basitarsus 1.4-1.7 mm (N=15).

Specimens examined: 13♂♂, 18♀♀ (holotype and paratypes), Wulingshan (700 m), Hebei Prov., S. Liu & C. Yang leg. From the larvae collected, the adults emerged in April and May, 1974-1976.

6. Genus *Vermipardus* Stuckenberg

Vermipardus Stuckenberg, 1960, S. Afr. anim. Life, 7: 227 (as subgenus of *Lampromyia*). Type species: *Lampromyia intermedia* Stuckenberg, 1955, by original designation.

Stuckenberg (1995b) raised *Vermipardus* from a subgenus of *Lampromyia* to a full genus and gave a key to species and a cladogram for this genus. It contains the following 3 species groups: *promontorii* group (*brincki* and *promontorii*); *intermedius* group (*intermedius*, and *univittatus* + [*basuto* + *sylphe*]); *brevirostris* group: (*londti* + *sathon*) + (*brevirostris* + *munroi*).

Stuckenberg (1997) added 3 new species and showed a revised cladogram for *Vermipardus*.

Subgenus 1: *promontorii* and *barraclooughi*. Subgenus 2 (*univittatus* group): *brincki* + [*trisignatus* + *whiteheadi*]; *intermedius*; *univittatus* + [*basuto* + *sylphe*]. Subgenus 2 (*brevirostris* group): 4 species.

Generic diagnosis (based on Stuckenberg, 1955, 1960, 1961, 1995a, b, 1996b, 1997 and on *V. intermedius*; quoted from 1996b: 245, 249). Similar to *Lampromyia* except as follows: Head: Antennal flagellum 8-segmented and gradually tapering apically and with the last joint much shorter than rest of flagellum. [Shape of palpus as in *Lampromyia*; see Nagatomi and Soroida, 1985: fig. 411 [= Fig. 251]. Proboscis 0.5-1.4 times as long as fore femur. "Labella laterally compressed, short, subtriangular, fused over about half their length, bearing stiff setae; about 6-7 pseudotracheae in each labellum." [The degree of facial protrusion varies with species as in *Lampromyia*].

Wing: Distance between h-crossvein and base of Rs about 3-4 times as long as basal section of Rs (Stuckenberg, 1995b: figs. 38-46) [Fig. 190].

Male genitalia (see many figures in Stuckenberg, 1960, 1961, 1995b, 1997): "Aedeagus is primitively tubular without lateral extensions."

V. intermedius (♂): Frons at median ocellus less than 1/2 as wide as one eye on a mid line and not more than twice as wide as ocellar triangle, but not narrower than that just above antenna (space between ocellar triangle and eye not wider than ocellar triangle); face (including lower portion)

Fig. 190. *Vermipardus brincki*,
♂. From Stuckenberg (1960).

parallel-sided and about as wide as frons. [It is unknown whether or not the characters above are applicable to other species of *Vermipardus*].

Vermipardus is distinguished from *Vermilyn*x by having the labella mentioned above, about 6-7 pseudotracheae in each labellum and aedeagus without lateral extensions. It is easily separated from *Leptynomia* by having last joint of flagellum shorter than preceding 3 joints together.

6. 1. (38). *Vermipardus barracloughi* Stuckenberg (Figs. 192, 238-241).

Vermipardus barracloughi Stuckenberg, 1997, Ann. Natal Mus., 38: 175. Type locality: South Africa (Transvaal: Bridal Veil Falls).

Distribution. South Africa.

6. 2. (39). *Vermipardus basuto* (Stuckenberg) (Figs. 195-198)

Lampromyia (*Vermipardus*) *basuto* Stuckenberg, 1960, S. Afr. anim. Life, 7: 260. Type locality: Lesotho (Basutoland: Bokong P. O.).

Distribution. Lesotho.

6. 3. (40). *Vermipardus brevirostris* (Bezzi) (Figs. 199-201)

Lampromyia brevirostris Bezzi, 1926, Ann. S. Afr. Mus., 23: 303. Type locality: South Africa (Natal: Thukela River valley: Mfongosi).

Distribution. South Africa.

6. 4. (41). *Vermipardus brincki* (Stuckenberg) (Figs. 190, 202-205, 249)

Lampromyia (*Vermipardus*) *brincki* Stuckenberg, 1960, S. Afr. anim. Life, 7: 253. Type locality: South Africa (SW Cape Province: Cape Peninsula: Skoorsteenkop mountain).

Distribution. South Africa.

6. 5. (42). *Vermipardus intermedius* (Stuckenberg) (Figs. 92-96, 214-217, 263-267, 283, 294, 304-305)

Lampromyia intermedia Stuckenberg, 1955, J. Ent. Soc. S. Africa, 18: 263. Type locality: South Africa (Eastern Cape Province: Grahamstown).

Distribution. South Africa.

6. 6. (43). *Vermipardus londti* Stuckenberg (Figs. 206-210)

Vermipardus londti Stuckenberg, 1995, Ann. Natal Mus., 36: 234. Type locality: South Africa (Eastern Cape Province: Boesmansriviermond).

Distribution. South Africa.

6. 7. (44). *Vermipardus munroi* Stuckenberg (Figs. 222-226)

Vermipardus munroi Stuckenberg, 1995, Ann. Natal Mus., 36: 232. Type locality: South Africa (Eastern Cape Province: East London).

Distribution. South Africa.

6. 8. (45). *Vermipardus promontorii* (Stuckenberg) (Figs. 227-230, 282, 306)

Lampromyia (*Vermipardus*) *promontorii* Stuckenberg, 1961, Ann. Natal Mus., 15: 120. Type locality: South Africa (SW Cape Province: Cape Peninsula: Table Mountain).

Distribution. South Africa.

6. 9. (46). *Vermipardus sathon* Stuckenberg (Figs. 231-237)

Vermipardus sathon Stuckenberg, 1995, Ann. Natal Mus., 36: 236. Type locality: South Africa (Natal: Ashburton).

Distribution. South Africa.

6. 10. (47). *Vermipardus sylvphe* Stuckenberg (Figs. 191, 211-213)

Vermipardus sylvphe Stuckenberg, 1995, Ann. Natal Mus., 36: 229. Type locality: South Africa (Natal: Drakensberg mountains).

Distribution. South Africa.

6. 11. (48). *Vermipardus trisignatus* Stuckenberg (Figs. 193, 242-245, 307)

Vermipardus trisignatus Stuckenberg, 1997, Ann. Natal Mus., 38: 180. Type locality: South Africa (W. Cape: Marloth Nature Reserve).

Distribution. South Africa.

6. 12. (49). *Vermipardus univittatus* Stuckenberg (Figs. 218-221, 308)

Lampromyia (*Vermipardus*) *univittata* Stuckenberg, 1960, S. Afr. anim. Life, 7: 257. Type locality: South Africa (Eastern Cape Province: Katberg mountain [4000 ft.]).

Distribution. South Africa.

Fig. 191. *Vermipardus sylvae*. Left, ♂; right, ♀. Refer to P. 139. From Stuckenberg (1995b, 1996b).

Figs. 192-194. *Vermipardus*. 192, head of *V. barracloughi*, ♂; 193, antenna of *V. trisignatus*, ♀; 194, antenna of *V. whiteheadi*, ♀. From Stuckenberg (1997).

6. 13. (50). *Vermipardus whiteheadi* Stuckenberg (Figs. 194, 246-248)

Vermipardus whiteheadi Stuckenberg, 1997, Ann. Natal Mus., 38: 185. Type locality: South Africa (W. Cape Prov.: Onrus).

Distribution. South Africa.

7. Genus *Vermitigris* Wheeler

Vermitigris Wheeler, 1930, Demons of the Dust, p. 274. Type species: *Vermitigris fairchildi*

Wheeler, 1930 (based on full grown larva and pupa), by monotypy.

Generic diagnosis (based on female of *V. sinensis*).

Figs. 195-205. Male genitalia of *Vermipardus*. 195-198, *V. basuto*; 199-201, *V. brevirostris*; 202-205, *V. brincki*. 195, 199, 202, fused gonocoxites and gonostyli, ventral view; 199, 'with right gonostylus inset in posterior view.' 196, 200, 203, tergum 9, cerci and sternum 10, dorsal view. 197, 201, 204, aedeagus, lateral view. 198, 205, aedeagus, posterior view. From Stuckenberg (1960).

Head: Frons at median ocellus less than 1.5 times as wide as that just above antennae, less than twice as wide as ocellar triangle, and much less than 1/2 as wide as one eye on a mid line. Space between ocellar triangle and eye distinctly narrower than ocellar triangle and concave. Face almost parallel-sided at upper portion, somewhat wider at lower portion, with narrowest part about as wide as frons just above antenna. Antennal flagellum with a slender 3-segmented arista and with a widened base (= flagellomere 1) which is longer than wide and gradually tapering apically. Palpal segment 2 widest around middle and then gradually tapering apically. Proboscis short and fleshy.

Figs. 206-221. Male genitalia of *Vermipardus*. 206-210, *V. londti*; 211-213, *V. sylvphe*; 214-217, *V. intermedius*; 218-221, *V. univittatus*. 206, hypopygium, dorsal view; 207, ventral aperture and apical guides; 208, left gonostylus in posterior view; 209, 212, tergum 9. 211, 214, 218, fused gonocoxites and gonostyli, dorsal view. 215, 219, tergum 9, cerci and sternum 10, dorsal view. 210, 213, 216, 220, aedeagus, lateral view; 210, with spiculae enlarged. 217, 221, aedeagus, posterior view. For lettering, see notes under Figs. 44-55. From Stuckenberg (1960, 1995b).

Figs. 222-237. Male genitalia of *Vermipardus*. 222-226, *V. munroli*; 227-230, *V. promontorii*; 231-237, *V. sathon*. 222, 227, 231, hypopygium, dorsal view. 228, same, ventral view. 223, 229, 234, tergum 9, dorsal view; 224, fused gonocoxites, ventral view. 226, 230, 235, aedeagus, lateral view; 225, 233, left gonostylus, posterior view; 236, apex of aedeagus, lateral view; 237, abdomen, lateral view. For lettering, see notes under Figs. 44-55. From Stuckenberg (1995b).

Figs. 238-249. Male genitalia of *Vermipardus*. 238-241, *V. barraclooughi*; 242-245, *V. trisignatus*; 246-248, *V. whiteheadi*; 249, *V. brincki*. 238, 242, 246, gonocoxites, gonostyli and dorsal bridge, dorsal view; 239, 243, gonostylus; 240, 244, 247, tergum 9, dorsal view; 245, 248, 249, aedeagus, lateral view, 'the deep incurvature marked by arrows.' From Stuckenberg (1997).

Figs. 250-255. *Vermitigris sinensis*, ♀. 250, head, anterior view; 251, palpus; 252, antenna; 253, wing; 254, hind tarsus, lateral view; 255, abdomen, dorsal view. Refer to p. 131. Original.

Wing: Vein R₄ almost vertical to vein R₅ and mouth of 2nd submarginal cell distinctly longer than vein R₅; vein R₂₊₃ strongly curved upward apically and apical portion of 1st submarginal cell almost vertical to its basal portion; 4th posterior cell open or closed (Edwards, 1932: fig. 1 [= Fig. 257]) and anal cell narrowly open. A narrow alula present.

Legs: Hind coxa without ventral (= anterior) knob.

Male genitalia: Not yet studied.

Key 11: *Vermitigris* species based on external characters

(except *V. fairchildi* whose adult is unknown)

1. Antennal scape, pedicel and flagellum (except arista in *V. sinensis*) pale orange or largely yellowish brown; abdominal segments 2-4 or 2-6 with a trace of light and dark rings 2
- Antennae dark, 3rd segment black, 1st two segments blackish-red; abdomen without a trace of light and dark rings, so conspicuous in *V. orientalis*; (after Oldroyd, 1947); [♀ unknown]; Borneo (Sabah) *V. infasciatus*
- 2(1). Antennal arista yellowish white; pleuron brownish yellow (except mesopleuron mainly blackish brown); hind femur irregularly brownish yellow; (after Brunetti, 1927); [♀ unknown]; Malaya and southern India *V. orientalis*
- Antennal arista dark brown to black; pleuron dark brown to black; hind femur entirely dark brown; [♂ unknown]; China (Guangxi Zhuang Autonomous Region) *V. sinensis*

7. 1. (51). *Vermitigris fairchildi* Wheeler (Figs. 349-350, 395-396)

Vermitigris fairchildi Wheeler, 1930, Demons of the Dust, p. 267 (based on full grown larva and pupa). Type locality: Sumatra (Deli Province: Konde [1500-2000 ft.]).

Distribution. Sumatra.

The adult stage of this species is not yet known.

7. 2. (52). *Vermitigris infasciatus* Oldroyd

Vermitigris infasciatus Oldroyd, 1947, Proc. R. ent. Soc. Lond. (B), 16: 22. Type locality: Borneo (Sabah: Bettotan, near Sandakan).

Distribution. Borneo.

The following are quoted from the original description.

The single male now before me is from Sandakan, N. Borneo, and is specifically distinct from *orientalis* Brunetti. In view of the locality it might be the adult of Shelford's species [a number of larvae and pupae from Mt. Penrissen, Sarawak recorded by Shelford (1900)], which was not named, but which Wheeler placed on record as 'probably *fairchildi*.' We have thus two possibilities: (1) this male may be the adult of Shelford's species, (2) Shelford's species may be the same as *fairchildi* Wheeler.

Neither of these points can be settled until larvae are collected and bred. In the meantime I think this species should be named.

♂. Differs from *V. orientalis* Brunetti — of which the type is in the British Museum — as follows: — Head: Antennae dark, third segment black, first two segments blackish-red.

Thorax: generally darker, especially the pleura, which are entirely black, with a little white

tomentum. In *orientalis* only the mesopleuron is black, the rest extensively yellowish.

Abdomen: first segment black, the rest warm brown. Second segment has yellowish lateral patches, which extend as light longitudinal stripes as far back as fifth segment. No trace of light and dark rings, so conspicuous in *orientalis*. Hairs brown, and uniformly distributed.

Legs: similar to those of *orientalis*, except that the coxae and hind femora are uniformly dark brown.

Wings: with venation characteristic of the genus, but with fore-border heavily coloured brown, and with a brown patch at the tip of the first basal cell.

Length of body 11 mm; wing 9 mm.

Some of the characters mentioned above are clearly subject to individual variation, as other specimens of *orientalis* have dark coxae and a brown costal border to the wing, and more brown on the hind femora. The dark antennae, *completely* dark hind femora, and the unbanded abdomen mark this species as distinct.

7. 3. (53). *Vermitigris orientalis* (Brunetti) (Figs. 256-257)

Lampromyia orientalis Brunetti, 1927, J. fed. Malay St. Mus., 13: 293. Type locality: Malaya ("two ♂♂ from the entrance to dark cave, Batu Caves, near Kuala Lumpur").

Distribution. Malaya; southern India.

Oldroyd (1947: 21) wrote, "A number of specimens bred by the late Mr. H. M. Pendlebury, and described by Brunetti (1927) as *Lampromyia orientalis*. Edwards (1932) examined a puparium of one of these and transferred the species to *Vermitigris* of which he gave the first description of the adult. The locality was Batu Caves, near Kuala Lumpur, and Bukit Kutu (3000 ft.), Selangor. The British Museum has one further specimen from near Petaling, F. M. S. (A. R. Sanderson), and one from Naraikkadu, Tinnevely Dt., S. India, 12th March, 1936 (B. M. - C. M. Exped.). The puparium is specifically distinct from that of *V. fairchildi* Wheeler."

The original description of *V. orientalis* (excepting some words) is quoted below.

Head seen from in front, barely wider than high, distinctly wider than thorax; eyes bare, separated by a nearly parallel sided vertex and frons, one-eighth the width of head at broadest part; ocellar tubercle large, well developed Frons white dusted, two narrow black streaks on upper part and elongate vertically placed black spot above antennae. Antennae; 1st joint long, cylindrical, 2nd short, cylindrical, about one-third as long as 1st, both pale orange; 3rd joint conical, as long as 2nd, a little browner, with long tapering yellowish white style which is as long as 1st joint. Face slightly narrower than frons, white dusted. Proboscis and palpi brown orange, Occiput flat, broadly snow-white dusted at sides; two roundish yellowish white dust spots behind vertex. Thorax brownish yellow, with three conspicuous dorsal blackish brown stripes narrowly separated; median one from front to hind margin, the others from considerably behind humeri and not quite attaining hind margin. Yellowish dust stripes between stripes and a spot of similar nature in front of wing base. Scutellum blackish, narrowly margined with yellow. Pleurae brownish yellow, mesopleura mainly blackish-brown, metapleura whitish dusted.

Abdomen more than twice as long as head and thorax together, cylindrical, at no part as wide as thorax, 1st segment short, yellowish-grey, hind margin and sides shining black; 2nd to 6th segments longer, subequal, about basal half of each bright ochraceous yellow, remainder black, the black more extensive on 5th and 6th segments. Pubescence of abdomen pale yellow, inconspicuous except on

Figs. 256-257. *Vermitigris orientalis*.
Antenna and wing. Refer to p. 131. From
Edwards (1932).

Figs. 258-267. Mouthparts. 258-262, *Vermileo comstocki*, ♀; 263-267, *Vermipardus intermedius*, ♀. 258-259, 263-264, labrum, cibarial pump, etc., ventral and lateral views; 260, 266, palpus and lacinia; 261, 267, labium (including labellum and theca), lateral view; 262, same, dorsal view; 265, showing presence of degenerate mandible. cp, cibarial pump; dc, dorsal cornu; h, hypopharynx; lc, lacinia; le, labellum; lr, labrum; ma, mandible; mi, mid-distal interlabellar process; pp, pharyngeal pump; pl, palpal segment 1; p2, palpal segment 2; sp, salivary pump; th, theca. Refer to p. 137. From Nagatomi & Soroida (1985).

Figs. 268-273. Head and mouthparts
of *Vermileo vermileo*, ♂. 268-269,
head, anterior and posterior views;
270, labrum, cibarial pump, etc.,
dorsal view; 271, palpus and lacinia;
272, prementum (=theca), ventral
view; 273, labellum, dorsal view. Cd
St, cardo-stipes; F, fulcrum (=cibarial
pump); La, lacinia; Lr, labrum; P Ph,
pharyngeal pump; Pp Mx, maxillary
palpus; PT, pseudotrachea; SM,
median sclerite (=mid-distal interla-
bellar process); Ss, sensory hair.
Other abbreviations are omitted in
explanation. Refer to pp. 137, 138.
From Chassagnard & Tsacas (1974).

sides of 1st segments.

Legs; hind pair, irregularly brownish yellow; trochanters and extreme base of femora blackish, hind tibiae becoming dark brown on apical third, hind metatarsus more than half as long as tibiae, much longer than rest of tarsus;

Wings. Very long, 12 mm, uniformly pale yellowish-grey;; halteres yellowish.

Length 12 mm. Expanse about 26 mm. Described from two ♂♂A further ♂ has been sent me for examination by Mr. Pendlebury taken at Bukit Kutu, Selangor, 3500 ft., 20th April, 1926. It agrees closely with the others but the black in the abdomen is a little more extensive and the hind femora are a little darker brown. Length 17 1/2 mm.

7. 4. (54). *Vermitigris sinensis* C. Yang (Figs. 250-255, 327)

Vermitigris sinensis C. Yang, 1988, Entomotaxon, 10 (3-4): 178. Type locality: China (Guangxi

Figs. 274-275. Labellum and pseudotracheae. 274, *Lampromyia flavida*, ♀; 275, *Leptynoma* (*Leptynoma*) *hessei*, ♀. Refer to p. 139. From Stuckenberg (1996b).

Figs. 276-285. Antenna. 276-278, *Leptynoma*; 279-281, *Lampromyia*; 282-283, *Vermipardus*; 284-285, *Vermilynx*. 276, *Leptynoma* (*Perianthomyia*) *maculata*, ♂; 277, *Leptynoma* (*Leptynoma*) *namaquensis*, ♂; 278, *Leptynoma* (*Leptynoma*) *appendiculata*, ♂; 279, *Lampromyia pilosula*, ♀; 280, *Lampromyia rebecca*, ♂; 281, *Lampromyia flavida*, ♀; 282, *Vermipardus promontorii*; 283, *Vermipardus intermedius*; 284-285, *Vermilynx jasoni*, ♂ & ♀ (without scape). Refer to pp. 129, 143. From Stuckenberg (1995b, 1996a, 1996b).

Figs. 286-293. Antenna of the Palaeartic *Lampromyia*. 286, *L. canariensis*; 287, *L. fortunata*; 288, *L. hemmingseni*; 289, *L. iberica* (without scape and pedicel); 290, *L. funebris*; 291, *L. cylindrica*; 292, *L. lecerfi*; 293, *L. pallida*. Refer to p. 143. From Stuckenberg (1998).

Zhuang Aut. Reg.: Langping, Tianlin Co., Bose Pr.).

Distribution. China (Guangxi Zhuang Reg.).

Besides the characters mentioned in Key 11, *V. sinensis* may easily be separated from *V. orientalis* in the following respects: Vein R₄ more vertical (see Figs. 253, 256); vein R₅ shorter (not over 1/2 as long as vein R₄).

Female. Head: Dark brown to black, and with pale gray pollen which is heavy on face; scape, pedicel and flagellomere 1 largely or partly yellowish brown; palpus and proboscis either yellowish brown or dark brown (the latter case probably occurs abnormally); ocellar tubercle, vertex, occiput, cheek, scape, pedicel, palpus and proboscis with black hairs. Width of frons at median ocellus 1.1-1.7 times that just above antenna, 0.2-0.3 times width of one eye on a mid line in direct frontal view, and 1.3-1.7 times width of ocellar triangle which is 0.8-1.0 times its length; width of frons just above antenna 0.6-0.9 times width of face at lowest part in direct frontal view and 0.3-0.5 times distance from antenna to median ocellus; distance from proboscis to antenna 1.9-2.3 times that from antenna to median ocellus; antenna 2.3-3.0 times distance from antenna to median ocellus; [N= 6].

Thorax: Dark brown to black and with pale gray pollen; mesonotum, scutellum, propleuron and anterior part of metapleuron with pale hairs which are short on the former two.

Wing: Membrane faintly tinged with yellowish in certain lights; veins yellowish brown; 4th posterior cell narrowly open, closed at wing margin or with short petiole; anal cell narrowly open; vein M₄ arising from 2nd basal cell; halter dark brown, with basal part (before narrowest part) yellowish brown.

Figs. 294-295. Female terminalia. 294, *Vermipardus intermedius*; 295, *Vermileo comstocki*. A, dorsal view; B, ventral view. Refer to p. 134. From Nagatomi & Iwata (1976).

Legs: Dark brown to black; fore and mid femora, hind tarsus, and basal 1/2 (or so) of hind tibia yellowish brown; fore and mid tibiae sometimes brown rather than dark brown; hairs on legs short, pale and black. Relative lengths of segments (excluding coxa and trochanter) of fore leg 138 (133-144): 136 (133-141): 100: 29 (26-31): 17 (15-19): 9 (9-10): 13 (11-15), of mid leg 167 (158-171): 174 (170-177): 98 (94-100): 26 (24-27): 12 (11-13): 9 (9-10): 12 (11-13), of hind leg 253 (242-258): 261

Figs. 296-303. Female terminalia and spermatheca of *Lampromyia*. 296-297, *L. hemmingseni*; 298-299, 302, *L. lecerfi*; 300, *L. flavida*; 301, *L. canariensis*; 303, *L. pallida*. 296, 298, 'showing shape of cercus and showing position of median swelling on S8 by arrow'; 297, 'S8, showing position of paired median swellings'; 299, 'showing apical emargination of T9' (dorsal view). Refer to p. 135. From Stuckenberg (1998).

(248-274): 123 (109-132): 29 (21-35): 17 (12-20): 10 (9-11): 16 (15-17) and in hind leg from the side, relative widths of femur, tibia and tarsomeres 1-3, 31 (29-32): 25 (24-26): 20 (18-23): 14 (12-16): 11 (10-11); [N= 4].

Abdomen: Dark brown to black; sterna 2-5 (except posterior part on sterna 4-5 or 5) and anterior band on terga 2-4 yellowish brown; hairs on abdomen short and chiefly black but pile on side of tergum 1 longer and pale.

Length: Body 17.1-18.3 mm; wing 12.6-14.0 mm; fore basitarsus 1.6-1.8 mm; (N= 4).

Male. Unknown.

Specimens examined: Holotype ♀ and paratypes 5 ♀♀, whose larvae were collected from Guangxi Zhuang Aut. Reg. (Bose Pr.: Tianlin Co.: Langping) by C. Yang on 28. v. 1982 and emerged on 8. vi. 1983 at Beijing.

7. 5. (55). *Vermitigris* sp. (Fig. 327)

C. Yang & Chen (1987) noted the presence of *Vermitigris* sp. from Tibet (Yadong Co.), based on some larvae collected by Fasheng Li on August 20, 1973.

A note given by C. Yang & Chen (1987) is here newly translated into English by Ding Yang.

Figs. 304-309. Spermatheca of *Vermipardus*. 304-305, *V. intermedius*; 306, *V. promontorii*; 307, *V. trisignatus*; 308, *V. univittatus*; 309, *Vermipardus* sp. Refer to p. 135. From Stuckenberg (1997).

Figs. 310-315. Female terminalia of *Vermipharis*. 310-311, *V. wudangensis*; 312, *V. ganquanensis*; 313, *V. minshanensis*; 314, *V. taihangensis*; 315, *V. yanshanensis*. 311, ventral view. Refer to p. 135. From C. Yang (1979) and C. Yang & Chen (1986, 1993).

The larvae collected below the rock wall near Yadong by Fasheng Li are larger in body size than that of *Vermiophis tibetensis*. They are identified as *Vermitigris* sp. *Vermitigris* was restricted to the Oriental region and had 3 known species distributed from Indonesia to southern India.

Recently, one Palaearctic species, *Vermitigris sinensis*, was found from China. The proleg on the 1st abdominal segment has 6-7 spines in *Vermitigris* sp. from Tibet, which differs from *V. sinensis* (with 4-5 spines) and is more similar to *V. fairchildi* Wheeler (with 6 spines), the type species of this genus. *V. fairchildi* (whose adult is still unknown), established on larva and pupa, and *Vermitigris* sp. from Tibet are different from each other in the structure of cephalo-pharyngeal skeleton and spiracle.

FOSSIL GENUS AND POSSIBLY RELATED FOSSIL FAMILY

The fossil Vermileonidae contains only one genus and species, that is, *Protovermileo electricus*. *Protobrachyceron liasinum*, only one representative of Protobrachyceridae, may be near in position to the Vermileonidae. *Archirhagio obscurus* was placed in Vermileonidae by Rohdendorf (1938) and then in Rhagionidae by Evenhuis (1994) but removed to the Archisargidae by Mostovski (1997) and by Nagatomi & Yang (1998) who deal with several Mesozoic Brachycera including Protobrachyceridae, Archisargidae, Eremochaetidae, etc. For position of these families, see Nagatomi & Yang (1998). Only *Protovermileo* is discussed here.

Genus *Protovermileo* Hennig, 1967, Stuttg. Beitr. Naturkd., 175: 6. Type species: *Protovermileo electricus* Hennig, 1967, by monotypy.

electricus Hennig, 1967: 26. Palaearctic: Baltic Region (Eocene / Oligocene) [Amber].

The above was quoted from Evenhuis (1994: 310). *Protovermileo electricus* is only one doubtless fossil taxon of Vermileonidae known up to the present. Body length is about 10.8 mm.

From Figs. 316-321 (Hennig, 1967: figs. 32-37), it is obvious that *Protovermileo* is most similar to *Vermiophis* but may be separated from the latter by having the apical segment of palpus club-shaped and the antennal flagellum (except basal joint) more slender.

The sister group of *Protovermileo* is difficult to determine. It is not necessarily *Vermiophis*.

GEOGRAPHICAL DISTRIBUTION

Wheeler (1930: figs. 19 and 24) and Hennig (1967: fig. 3 [= Fig. 322]) illustrated the distributional map of Vermileonidae. The record is summarized below.

Stuckenberg (1995b: 249) wrote, "It is notable that species allopatry is evident throughout the range of the [Vermileonidae] in [South Africa]. I know of no locality in South Africa where two species of vermilionids (even of different genera) occur together."

The Vermileonidae is absent from Australia and South America. Nagatomi (1982, 1991) discussed the geographical distribution of the lower Brachycera including Vermileonidae (but excluding Nemestrinoidea, Asiloidea and Empidoidea) and thought that the Vermileonidae were extinct in Australia and South America.

(1) *Lampromyia* (12 species): Europe (France; Spain); northwest Africa (Algeria; Tunisia;

Figs. 316-321. *Protovermileo electricus*, ♂ from Baltic amber. 320, latero-ventral view; 321, ventral view. From Hennig (1967).

Morocco); Canary Is.; southern Africa (Lesotho; South Africa; Zimbabwe).

Spain and France (2 species): *funbris* and *iberica*.

Algeria, Tunisia and Morocco (4 species): *cylindrica* (Algeria: lowlands); *lecerfi* (Morocco: High Atlas); *nigripennis* (Morocco: Middle Atlas); *pallida* (Algeria, Tunisia and probably Morocco).

Canary Is. (3 species): *canariensis*; *fortunata*; *hemmingseni*. See Fig. 323.

Southern Africa (3 species): *flavida*; *pilosula*; *rebecca*.

Stuckenberg (1996b: 240) wrote, "*Lampromyia* is now seen to be a genus having its main centre of diversity in the south-western Palearctic Region, and a clade with three species in eastern Southern Africa."

The three species in Canary Is. (after Stuckenberg, 1971) live separately as follows: *canariensis*,

Fig. 322. Geographical distribution of Vermileonidae. (*Archirhagio* belongs to Eremochaetidae of Nemestrinoidea; *Lampromyia* means *Lampromyia* s. lat.; it is unknown to us where the original record from Turkey and Syria is). From Hennig (1967).

Fig. 323. Geographical distribution of *Lampromyia* (*canariensis* group) from Canary Islands. From Hemmingsen & Nielsen (1971).

Tenerife; La Palma; La Gomera; El Hierro. *fortunata*, Gran Canaria. *hemmingseni*, Lanzarote; Fuerteventura.

The three species in southern Africa (after Stuckenberg, 1996b): *pilosura*, eastern parts of southern highveld in Cape Province, Orange Free State, Lesotho, western Natal; *flavida*, Zimbabwe (eastern highlands near Mutare); *rebecca*, Zimbabwe (Harare district).

(2) *Leptynoma* (6 species): Subgenus *Leptynoma* (4 species), South Africa; subgenus *Perianthomyia* (2 species), South Africa (*maculata*) and Namibia (*phantasma*).

Except for *phantasma* from Namibia, *Leptynoma* occurs in the south-western Cape Province and Namaqualand.

Stuckenberg (1996b: 253) wrote: (i) *sericea* + *appendiculata* — these two species occur in the distinctive 'fynbos' shrubland of the south-western and southern Cape; their wide altitudinal range is

explained by this association, as the fynbos extends from the coastal plain uninterruptedly to the summits of the Cape Fold Mountains. (ii) *hessei* + *namaquaensis* — these species are associated with succulent karoo flora, but in contrastingly different situations; *hessei* occurs in the southern Cape particularly in the intermontane areas and valleys, where rainshadow effects produce xeric conditions and succulent or mixed floras — this species also ranges northwards into Namaqualand, via intermontane valleys where rainshadow effects are strong; *namaquaensis* occurs on the Knersvlakte, a late Tertiary plain bordering the Atlantic coast in southern Namaqualand, and it probably ranges northwards over the coastal hinterland and into valleys in the east that open onto the coastal plain.

(3) *Vermileo* (10 species): southern Europe, 3 species; northern Africa, 1 species; North America, 2 species; Central America, 4 species.

vermileo: Widely distributed in Spain, France, Italy, southern Tyrol, Dalmatia, Balearic Is., and Bulgaria.

nigriventris: Spain and southern Portugal.

ater: Crete (Greece).

Stuckenberg (1965b: 495) wrote, "At present I have no records which indicate that *vermileo* and *nigriventris* are sympatric."

niloticus: Sudan; the species recorded from Egypt may refer to *niloticus* (after Edwards, 1935).

comstocki: "Its known range extends along the Sierra Madre and Sierra Nevada Ranges from Mt. Wilson in Southern to Lassen Peak in Northern California, a distance of some 850 miles" (after Wheeler, 1930: 174) (see Wheeler, 1930: fig. 19; Hennig, 1967: fig. 31 [= Fig. 322]).

"In April 1934 the writer reared specimens of this species from material collected in February in the Coast Range at Pinnacles National Monument at an elevation of about 1600 feet" (after DeLeon, 1938: 167).

opacus: California, Nevada, Utah, Colorado and New Mexico. This species is not sympatric with *comstocki*.

Central American species: *fascipennis* and *willetti* (Mexico); *dowi*, Cuba; *tibialis*, Jamaica.

The type locality of *fascipennis* is "Guerrero, Xucumanatlan" and that of *tibialis* is "between Mexico, D. F., and Cuernavaca, Morelos" (after James, 1968: 2).

(4) *Vermilynx* (2 species): South Africa and Namibia.

jasoni: "..... Springbok District of northern Namaqualand. This locality is less than 200 km south of the Richtersveld sites where *V. vansoni* was found, but the two species [*jasoni* and *vansoni*] are nevertheless remarkably different" (after Stuckenberg, 1996a: 198).

vansoni: "The Richtersveld is a semiarid mountainous terrain in the extreme northwestern corner of the Northern Cape Province, bounded to the north by a large bend in the Orange River. The region is renowned for its wealth of endemic plant taxa. The Rosh Pinah locality is in a topographic and biological extension of the Richtersveld in Namibia north of the Orange River" (after Stuckenberg, 1995a: 33).

(5) *Vermiophis* (7 species): northern China (6 species) and Tibet (1 species). See Figs. 324-326. They are apparently allopatric. This genus is still unrecorded from southern China (including Yunnan and Taiwan) and north-east China (= Manchuria). One of us (C. Y.) failed to find any Vermileonidae in Hainan in June, 1997.

(6) *Vermipardus* (13 species): Stuckenberg (1997: 173) wrote, "This is the most primitive

Fig. 324. Geographical distribution of 3 species of *Vermiophis* from China. Circle, *V. ganquanensis*; triangle, *V. minshanensis*; square, *V. wudangensis*. Original.

Fig. 325. Geographical distribution of 2 species of *Vermiophis* from China. Circle, *V. taihangensis*; triangle, *V. tibetensis*. Original.

Fig. 326. Geographical distribution of 2 species of *Vermiphis* from China. Circle, *V. taishanensis*; triangle, *V. yanshanensis*. Original.

Fig. 327. Geographical distribution of 2 species of *Vermitigris* from China. Circle, *Vermitigris* sp. (whose adult is unknown); triangle, *V. sinensis*. Original.

Fig. 328. Distribution of *Vermipardus* in South Africa. Subgenus 2 comprises the *brevirostris* and *univittatus* species-groups; the *univittatus* group is represented on this map by its three subgroups. Both Subgenus 1 and the *brincki* subgroup of Subgenus 2 have a species restricted to the Cape Peninsula. From Stuckenberg (1997).

afrotropical vermilionid genus. It is endemic to South Africa, occurring primarily in eastern and south-eastern montane regions of higher rainfall, on or seawards of the Great Escarpment (fig. 1). It has not been found on the high plateau terrain of the eastern interior, or in the Karoo, in the relatively dry interior parts of the Eastern and Western Cape Provinces, or in Namaqualand. One species occurs in Lesotho and two on the Cape Peninsula. Most of the known species are in KwaZulu-Natal, but this may be a collecting artefact." See Fig. 328.

(7) *Vermitigris* (5 species): Malaya, Sumatra, Borneo, southern India, southern China and Tibet.

fairchildi (based on larva and pupa; adult unknown): northern Sumatra (Deli Province: Konde [1500-2000 ft.]).

infasciatus: northern Borneo (Sabah: Bettotan near Sandakan).

orientalis: Malaya (near Kuala Lumpur; Selangor; near Petaling) and southern India (Tinnevely Dt.).

sinensis: southern China (Guangxi Zhuang Autonomous Region: Tianlin Co.). See Fig. 327.

sp. (based on larva): Tibet (Yadong Co.). See Fig. 327.

CITATIONS FROM THE LITERATURE ON IMMATURE STAGES

Larva

Our search of the literature on immature stages is not complete, but we hope that the important works

with illustrations are selected below. The head skeleton and spiracle are mentioned separately as well as pupa and egg.

Lampromyia canariensis from Canary Is. (Tenerife): Wheeler (1930: fig. 25: e-f); Fernández (1952: figs. 1-5).

Lampromyia fortunata from Canary Is. (Gran Canaria): Hemmingsen (1963) (fig. 1: a-c).

Lampromyia pallida from Tunisia: Marchal (1897: figs. 6-7, 9-11).

Leptynoma sericea from South Africa: Engel (1929: figs. 1-2); Wheeler (1930: fig. 25: g-h).

Vermileo comstocki from California: Wheeler (1930: figs. 22[A-C], 25[c-d], 30[A-D]); Teskey (1981: figs. 4-5).

Vermileo vermileo from southern Europe: De Geer (1752: figs. 3-6); de Réaumur (1753: figs. 6-8); Wheeler (1930: fig. 25: a-b); Tsacas (1962: fig. 22; plate 2: 9-12).

Vermileo sp. from Egypt (= possibly *V. niloticus* from Sudan): Hafez & El-Moursy (1956a: figs. 7-9).

Vermiophis ganquanensis from China: C. Yang (1979: figs. 7-8).

Vermitigris fairchildi from Sumatra: Wheeler (1930: fig. 25:i-j).

The vermilionid larva is characterized below. Body gradually wider posteriorly and at abdominal segment 7 widest. Abdomen: Segment 1 with a single unpaired mid-ventral pseudopod whose apical portion has several rigid spines; segment 7 with a transverse row of comb-like spines on the raised posterodorsal border; segment 8 "terminating in four flat, broadly finger-shaped lobes,Just anterior to the bases of this latter pair and on the mid-ventral line of the last segment there is a slit-shaped aperture, the anus (fig. 30 D an)" (after Wheeler, 1930: 179). All or almost all of these characters are easily judged as autapomorphic for Vermilionidae. For more detailed family diagnosis, see Teskey (1981: 530). See Figs. 329-340.

The generic and specific characters lie on the pseudopod (see Figs. 341-350), although the number of spines varies with individual. Wheeler (1930: 252-253, 272) wrote:

[*Lampromyia canariensis*]: The pseudopodial papilla (fig. 25, e, f) is deeply cleft transversely into two protruding, beak-like processes, the posterior of which bears on its hinder surface a group of 6 to 8 backwardly directed spines.

[*Vermitigris fairchildi*]: "The most striking of these is the structure of the pseudopod (fig. 25, i, j) on the ventral side of the first abdominal segment. In profile the tip of this appendage is evenly rounded and hemispherical, with a distinctly differentiated and uncleft papilla and with a low median longitudinal thickening bearing a series of straight, rigid, radiating spines, shorter at the ends of the series than in the middle. These spines vary in number, apparently, with the age of the larva, a full grown specimen having six, the smallest four and an individual of intermediate size, five."

The number of spines on pseudopod (on abdominal segment 1) is given below.

Lampromyia pallida: 4 (of which 2 are present outside papilla) (Marchal, 1897: fig. 10).

Vermileo vermileo: 4 (Wheeler, 1930: fig. 25: a-b); 5 or so (Tsacas, 1962: plate 2: fig. 12).

Vermileo sp.: 4 or 5 (Hafez & El-Moursy, 1956a: 292).

Vermiophis ganquanensis: 6 (rarely 8) (C. Yang, 1979: 89).

Vermipardus promontorii: 2 (Stuckenberg, 1961: 123).

Vermitigris orientalis: 3 (Edwards, 1932: 219).

Vermitigris sinensis: 5 (rarely 4); in smallest specimens 3-4 (C. Yang, 1988: 182).

Here, the number of spines (arranged comb-like) on the posterior border of abdominal segment 7

Figs. 329-340. Larva of Vermilionidae. 329-330, *Vermileo vermileo*; 329, dorsal view; 330, abdominal segment 1, lateral view; from Tsacas (1962). 331, *Vermileo comstocki*, lateral view; from Teskey (1981b). 332-333, *Vermileo* sp. from Egypt (= possibly *V. niloticus*); 332, abdominal segment 1; 333, terminal part of fullgrown larva; from Hafez & El-Moursy (1956a). 334-336, *Lampromyia fortunata*; 334, lateral view; 335-336, terminal part, dorsal and ventral views; from Hemmingsen (1963). 337-338, *Leptynoma (Leptynoma) sericea*, lateral view; 338, pseudopod (=ventral papilla) on abdominal segment 1; from Engel (1929). 339-340, *Vermiophis ganquanensis*, lateral view; 340, pseudopod on abdominal segment 1; from C. Yang (1979). ABD. 1, 2, 7, 8, abdominal segments 1-2, 7-8; PD, pseudopod; Ps, pseudopod; PS, SP, posterior spiracle; StA, anterior spiracle; StP, posterior spiracle; TH. 3, thoracic segment 3; TSt, vestige of spiracle (=trace de stigmat). 1, Head; 2-4, thoracic segments 1-3; 5-12, abdominal segments 1-8.

are also introduced below.

Lampromyia canariensis: 18 (Wheeler, 1930: 253).

Lampromyia fortunata: 20 (Hemmingsen, 1963: fig. 1: b).

Vermileo comstocki: 16 (Wheeler, 1930: fig. 30: E; Teskey, 1981: fig. 39.5).

Vermileo vermileo: 11 (Tsacas, 1962: fig. 22).

Vermileo sp.: 11 (Hafez & El-Moursy, 1956a: fig. 9).

Vermiophis ganquanensis: 14 (C. Yang, 1979: 89).

Vermitigris fairchildi: 12 (Wheeler, 1930: 272).

Vermitigris orientalis: 12 (Edwards, 1932: 218).

Vermitigris sinensis: 12 (C. Yang, 1988: 182).

Judging from Figs. 341-348, *Lampromyia* (or *Lampromyia* s. lat.) may possibly be separated from *Vermileo* in the following point: some of rigid spines on pseudopodial papilla present at or near base of posterior surface. In *Vermileo*, rigid spines are present only at apex.

Wheeler (1930: 253) wrote as to *Lampromyia canariensis*: "The four finger-shaped anal lobes seem to be shorter than in the *Vermileo* larva and this is true also both of the hairs along the sides of the thorax and abdomen and of the recurved setae forming the transverse comb on the seventh abdominal segment. These setae, however, are more numerous, there being 18 of them instead of 10 or 12 as in *Vermileo vermileo*".

It is uncertain whether or not the characters above are significant. Further comparison is needed for finding definite distinctions between *Lampromyia* (or *Lampromyia* + *Leptynoma* + *Vermilynx* + *Vermipardus*) and *Vermileo* based on a number of species.

On the other hand, *Vermitigris* is easily separated from *Lampromyia* and *Vermileo* by the shape of pseudopod and arrangement of spines on it as shown in Figs. 349-350. Moreover, Wheeler (1930: 272-273) wrote, "[The comb-like setae on posterior border of abdominal segment 7] are not flattened at the base as in *Vermileo* and *Lampromyia* but cylindrical and alternating with them is a second series of similar and much shorter bristles. There are also series of three stout dark-colored bristles on each side of the middle line and arising from paired transverse swellings or welts on the dorsal and ventral sides of the seventh and the dorsal side of the sixth abdominal segment."

The specific characters of *Vermitigris* are given below.

Vermitigris orientalis (from Edwards, 1932: 219): In the Malayan larval skin examined the pseudopod of the first abdominal segment has only three strong spines in the row, not six as described by Wheeler for full-grown specimens of *V. fairchildi*; also the armature of segments 5-7 of the abdomen is different, as each of these segments in *V. orientalis* carries spines or bristles both dorsally and ventrally; on segment 5 the dorsal bristles are short, the ventral bristles much longer but slender and pale; on 6 the paired groups of stout, dark bristles are placed close together both dorsally and ventrally, each group having 3-4 long and 2-3 short bristles; on 7 the dorsal groups of several short bristles are placed very wide apart, near the sides of the segment, but the ventral groups, each of three long, stout bristles are quite close together;

Vermitigris sinensis (from C. Yang, 1988: 182): "The pseudopod on the ventral side of the first abdominal segment with a longitudinal row of strong spines always 5 in number (rarely 4, the smallest 3-4 only), that of the *fairchildi* is 6 (smallest 4-5), and of the *orientalis* is 3 in number. The comb on [the posterior margin of the seventh] abdominal segment with 12 long and strong spines and a second

Figs. 341-350. Pseudopod on abdominal segment 1. 341-342, *Vermileo vermileo*; 343-344, *Vermileo comstocki*; 345-346, *Lampromyia canariensis*; 347-348, *Leptynoma (Leptynoma) sericea*; 349-350, *Vermitigris fairchildi*. 341, 343, 345, 347, 349, end view; 342, 344, 346, 348, 350, lateral view. From Wheeler (1930).

Figs. 351-353. Larval head skeleton of *Vermileo comstocki*, dorsal, ventral and lateral views. a, uncinat sclerite; b, lateral sclerite; c, small lineal sclerite; g, gula (?); h, head segment of larva; hc, head capsule; l, labium (?); lb, labrum; lg, chitinized ligament; mx, maxilla; m, mandible; p, maxillary palp; t, first thoracic segment; vp, ventral process of head capsule. From Wheeler (1930).

series of shorter ones as in the two known species. The spines of the sixth and seventh abdominal segments both on the dorsal and ventral side are rather similar to those in *orientalis* than to those in *fairchildi*,"

Vermiophis ganquanensis is peculiar among the known Vermileonid larvae: pseudopod with two apical knobs, that is, anterior and middle ones, and behind middle knob "with two longitudinal rows of 6 (rarely 8) strong spines" (after C. Yang, 1979: p. 89 & fig. 8). See Figs. 339-340.

Further, the respective description in some species is given below.

Figs. 354-358. Larval head skeleton of Vermileonidae. 354-356, *Leptynoma* (*Leptynoma*) *sericea*; 357, *Lampromyia* sp. from Spain; 358, *Vermileo comstocki*. 354, lateral view; 355, mandible and labrum; 356-358, dorsal view. ant, antenna (= correctly maxillary palpus); b scl, basal sclerite; cr, cranium; lbr, labrum; md, mandible; mx, maxilla; mx plp, maxillary palpus (= correctly part of maxilla). Refer to p. 142. 354-357, from Engel (1929); 358, from Teskey (1981b).

Fig. 359. Larval head skeleton of *Vermileo vermileo*, dorsal view. A, antenna; ABMVex, basal article of mandible, externally twisted part; ABMVin, basal article of mandible, internally twisted part; Ar a, anterior articulation; Ar p, posterior articulation; B PI D, thickened border of dorsal plate; Cib-ph, cibarium-pharynx; C Md, mandibular hook; EPD, swelling on dorsal plate; FC, fronto-clypeus; J, cheek (gena); L P, posterior margin of prothorax; Lr, labrum; Mx, maxilla; PMx, maxillary palpus; PrP, paraclypeal phragma; SFG, front-gena suture; S Lr, labrum suture; T, tentorium; TBI, inferior tentorium branch; TBS, superior tentorium branch. Refer to p. 142. From Tsacas (1962).

Figs. 360-363. Parts of larval head of *Vermileo vermileo*. 360, head, anterolateral view; 361, external mandible-maxilla complex and anterior part of dorsal plate, lateral view; 362, median longitudinal section of head, lateral view; 363, labium, anterior view. Cib-Ph, cibarium-pharynx; Cl, clypeus; Csal, salivary duct; En, spine; Ga, galea; Hy, hypopharynx; Lac, lacinia; Lb, labium; Lop, ventral lobe of maxillary palpus; Md, mandible; Oe, oesophagus; Osal, salivary orifice; Ph, pharynx; PL, anteroventral protuberance of labrum; PLb, labial palpus; Ps, sensory papilla; Sal, salivarium; SEp, epipharyngeal sclerite. For other letterings, see notes under Fig. 359. Explanation of some letterings is omitted. Refer to p. 142. From Tsacas (1962).

Vermipardus promontorii (from Stuckenberg, 1961: 123): [Very similar to *Leptynoma sericea* whose larva is described by Engel (1929)]. On the ventral surface of both segments 10 and 11 the larva has a pair of small, closely adjacent pseudopods which bear reclinate, hollow spines similar to (though smaller than) those on the dorsal surface of segment 10. On the lateral parts of segments 10

Figs. 364-370. Parts of larval head of *Chrysopilus auratus* (Rhagionidae). 364-365, anterior part of head, lateral and ventral views. 366-367, median longitudinal section of head, lateral view; 367, anterior part. 368-369, two successive cross sections of labium. 370, sclerotic part of hypopharynx and cibarium-pharynx, dorsal view. CE, brain; Csl, lateral superior crest of labrum; GglFr, frontal ganglion; DGa, teeth of galea; DLr, teeth of labrum; Em, membranous spines of mandible-maxilla complex; Ep, epipharynx; ExpT, posterior expansion of tentorium; LoLb, anterior thickened lobe of labium; Pg, comb of labrum; Pl D, dorsal plate; Psal, salivary duct; R Hy, contraction of hypopharynx; Scl, clypeal silk-like hair; SHy, hypopharynx sclerite; SLb, labial sclerite; Ss, sensory silk-like hair; St, stipes. For other letterings, see notes under Figs. 359, 360-363. Refer to p. 142. From Tsacas (1962).

and 11 there are groups of exceptionally long spines. The anal segment does not have a dense lateral fringe of ventrally-directed hairs like that in *sericea*; the four lobes of this segment are well formed and distinct. The pseudopod on the underside of the fifth segment [= abdominal segment 1] consists of a swollen basal part, which bears two narrow, fleshy lobes placed one behind the other, the anterior one being the largest. Behind the second lobe is a group of short, stiff, hollow spines. The mouthparts

Figs. 371-376. Parts of larval head of *Rhagio scolopaceus* (Rhagionidae). 371-372, anterior part, a ventral view; 373, same, anterior view; 374, median longitudinal section of head, lateral view; 375, labium, etc., ventral view; 376, hypopharynx and anterior part of cibarium-pharynx plate, dorsal view. Asc, saw appendix; SMI, additional silk-like hair; Zsc, sclerotic zone of plug in labrum. For letterings, see notes under Figs. 359, 360-363, 364-370. Refer to p. 142. From Tsacas (1962).

are almost identical to those of *sericea*.

Lampromyia fortunata (from Hemmingsen, 1963: 240) (Figs. 334-336): The figure corresponds to that given by Fernández (1952, fig. 2) of [*Lampromyia canariensis*], but for his segment numbers being equal to mine less 1, segments 1-2 being called 1 by Fernández. In addition to the large forward directed comb of flat hairs on the posterodorsal border of segment 11 and the backward directed minor bipartite combs on the anteroventral borders of segments 10 and 11 and the posterodorsal

border of segment 10, also mentioned or shown in profile in Fernández's figure, there is a minor backward directed bipartite comb on the posteroventral border of segment 9. Other groups of mostly lateral hairs are seen. The tips of the bristles of the large comb appear to be less recurved or hooky, and the number of hairs in some of the various combs greater, than in the corresponding comb of a larva of *Vermileo vermileo* from Spain kindly given to me by Fernández. The spiracle of the amphipneustic larva are situated laterally on segment 2 and dorsally on segment 12; and the anus, ventrally on segment 12.

Larval head skeleton

Lampromyia pallida: Marchal (1897: figs. 11-14).

Lampromyia sp. from Spain: Engel (1929: fig. 7).

Leptynoma sericea: Engel (1929: figs. 5-6).

Vermileo comstocki: Greene (1926: fig. 6); Wheeler (1930: fig. 32: A-C); Teskey (1981b: fig. 39.4).

Vermileo vermileo: Tsacas (1962: figs. 24-35; plate 2 [fig. 14]); Ludwig *et al.* (1996: fig. 1 B, C, D).

Vermileo sp. from Egypt (= possibly *V. niloticus* from Sudan): Hafez & El-Moursy (1956a: fig. 10).

Tsacas (1962) gave a remarkably detailed account on *Vermileo vermileo* and made a precise comparison among *V. vermileo* L., *Rhagio scolopaceus* L. and *Chrysopilus auratus* F., showing a deep discrepancy in cephalopharyngeal apparatus between Vermileonidae and Rhagionidae (Figs. 359-376). For phylogenetic significance of head capsule, see DISCUSSION section.

For detailed description of *Vermileo comstocki*, see Wheeler (1930: 195-201) (Figs. 351-353) and Teskey (1981: 530) (Fig. 358).

Lampromyia canariensis (from Wheeler, 1930: 252): [This species] is more like the larva of *V. comstocki* than that of *V. vermileo* in having the head-capsule relatively broad and short, but the latter consists of paler chitin ventrally. The eyes are clearly discernible even in the adult larva, as two small pigmented bodies in the usual position under the anterior end of the head-capsule and behind the labial sclerites.

Vermileo vermileo (from Wheeler, 1930: 196) (Figs. 351-353): [Head capsule] is decidedly more slender in *V. vermileo* than in any of the other species [of Vermileonidae] which I have examined.

Vermitigris fairchildi (from Wheeler, 1930: 273): The mouthparts and cephalopharyngeal apparatus are much like those of *Lampromyia*, the head-capsule being black, robust, heavily chitinized and scarcely more than twice as long as broad, but the ventral tendon, which is very narrow and incompletely chitinized in *Vermileo* and *Lampromyia*, forms a distinct linear sclerite, like the piece which I have called the gula, but shorter. The mandibles are stout, with rather coarsely serrated borders.

Vermitigris orientalis (from Edwards, 1932: 219): The head capsule also seems to be rather longer than in *V. fairchildi* (fully twice as long as broad), and without chitinisation of the ventral tendon.

Vermitigris sinensis (from C. Yang, 1988: 182): "The head-capsule also seems to be like *orientalis* and much longer (about triple as long as broad)".

Judging from Engel (1929: figs. 5 & 7), maxilla is forked in *Leptynoma sericea* (Fig. 356) but not in *Lampromyia* sp. from Spain (Fig. 357) at sclerotized distal part.

Larval spiracles

Leptynoma sericea: Engel (1929: figs. 3-4).

Figs. 377-382. Anterior and posterior spiracles of larva in Rhagionidae and Vermileonidae. 377-378, *Rhagio scolopaceus*; 379-380, *Chrysopilus auratus*; 381-382, *Vermileo vermileo*. 377, 379, 381, anterior spiracle; 378, 380, 382, posterior spiracle. Refer to p. 143. From Tsacas (1962).

Vermileo comstocki: Wheeler (1930: fig. 31: A-D).

Vermileo vermileo: Tsacas (1962: fig. 23; plate 2 [fig. 15]).

Leptynoma sericea (from Wheeler, 1930: 183-184) (Figs. 391-392): All the structures above described for *V. comstocki* are present in these larvae and of essentially the same form, but larger and much clearer. The stigmatic chamber of the posterior spiracle is, however, shorter than in *V. comstocki* and the orifice of the central piece is much larger, more conspicuous and more eccentric. There are 18 to 19 radial tubules in the posterior but only six, as Engel noticed, in the anterior spiracle. The trabeculae of the long stigmatic chamber of the latter are noticeably more distinct than in *V. comstocki*.

Vermileo comstocki (from Wheeler, 1930: 181-182) (Figs. 383-385): [The respiratory orifices, or spiracles] (fig. 30 A, asp, E, psp) are so inconspicuous that they are easily overlooked even in careful preparations of the chitinous cuticle, and when found are not easily interpreted. The larva is amphipneustic, i. e., possesses only two pairs of spiracles, one on the sides of the prothoracic, the other near the mid-dorsal line of the eighth abdominal segment. They form the anterior and posterior terminations of the pair of long tracheal trunks. The posterior spiracle when seen from the surface in cleared specimens presents the appearance of fig. 31 B [fig. 384].The anterior spiracle, as shown in fig. 31 D [= Fig. 386], though much smaller and more difficult to analyze than the posterior, is nevertheless constructed on the same plan. The radial tubules, however, are only about half as numerous (8 to 10), and the cantral piece and its opening are very small. There is a distinct elongated Filzkammer [= stigmatic chamber] traversed by oblique trabeculae.

Figs. 383-390. Larval and pupal spiracles of *Vermileo comstocki*. 383-385, larval posterior spiracle, in profile, from surface and in longitudinal section; 386, larval anterior spiracle in profile; 387-390, pupal spiracle on prothorax, abdominal segment 1, segment 4 and segment 8. Explanation of abbreviations is omitted. Refer to p. 143. From Wheeler (1930).

Figs. 391-394. Larval and pupal spiracles of *Leptynoma sericea*. 391-392, larva; 393-394, pupa; 391, anterior spiracle; 392, posterior spiracle; 393, prothoracic spiracle; 394, abdominal spiracle. Refer to p. 143. From Engel (1929).

Vermipardus promontorii (from Stuckenberg, 1961: 123): Morphologically the larva is extremely like that of [*Leptynoma*] *sericea* [Westwood] as described by Engel (1929). The prothoracic spiracles differ in detail, having 8 equal stigmatic openings in a fan-shaped group. The posterior spiracle has 5-6 stigmatic openings.

Vermitigris fairchildi (from Wheeler, 1930: 273): "The spiracles are large and distinctly chitinized and their radial tubules, which number six in the anterior and ten in the posterior pair, are unusually large and distinct".

Figs. 377-382, reproduced from Tsacas (1962), represents the spiracles of *Rhagio scolopaceus* (Rhagionidae), *Chrysopilus auratus* (Rhagionidae) and *Vermileo vermileo*. For importance and significance of larval and pupal spiracles, see DISCUSSION section.

Pupa

The species, which have been illustrated, are listed firstly.

Lampromyia pallida from Tunisia: Marchal (1897: fig. 8).

Leptynoma sericea from South Africa: Engel (1929: figs. 8-9) (= Wheeler, 1930: fig. 46).

[*Lampromyia* or *Leptynoma*] sp. (from ?) ('leg. et det. Dr. Engel'): Brauns (1954: fig. 35: C).

Vermileo comstocki: Greene (1926: fig. 11); Wheeler (1930: fig. 27).

Vermileo vermileo: Brauns (1954: fig. 35: B).

Figs. 395-402. Pupa of Vermileonidae. 395-396, *Vermileo vermileo*, ventral and lateral views; 397-398, *Vermitigris fairchildi*, ventral and lateral views; from Wheeler (1930). 399, *Vermileo* sp. from Egypt (= possibly *V. niloticus*), ventral view, from Hafez & El-Moursy (1956b). 400-401, *Leptynoma sericea*, from Engel (1929). 402, *Vermiphis ganquanensis*; terminal part of abdomen, ventral view; from C. Yang (1979). ANT, antenna; ABD1, ABD2, abdominal segments 1-2; LB, labium; L. EP, labrum epipharynx; LG, leg; MX, maxilla; MX. P, maxillary palp; SP. 1, 1st abdominal spiracle.

Vermileo sp. from Egypt (= possibly *V. niloticus* from Sudan): Hafez & El-Moursy (1956b: figs. 9-10).

Vermiphis ganquanensis: C. Yang (1979: fig. 9).

Vermitigris fairchildi: Wheeler (1930: fig. 47).

Vermitigris sinensis C. Yang (1988: figs. 2-3).

Full quotation of the description by various authors is given below in some of the species listed above (Figs. 395-402).

Lampromyia canariensis (from Wheeler, 1930: 254-255): The pupa, which is very much like that

of *L. sericea* (fig. 46) [= Figs. 400-401], measures 13-15 mm and has the head and thorax dark brown, the abdomen more yellowish with pale brown segmental annulations. The surface, as in other species of the genus and in *Vermileo*, is covered with agglutinated sandgrains and bears the shrunken last larval skin attached to the terminal abdominal segments. When the larval skin and sand are removed by immersion in caustic potash solution the abdomen is seen to end in two backwardly directed, parallel appendages and each of its segments from the first to the seventh bears, on each side near the middle, a large, pointed spine-shaped tubercle. The abdominal spiracles are also on tubercles and are much as in *V. comstocki* but larger. The integument of the head and thorax is smooth but that of the abdomen is very finely and transversely rugulose, with areas of small rounded tubercles, especially on the sides and more posteriorly also on the intersegmental membranes. The antepenultimate segment bears a pair of posterolateral appendage-like swellings covered with minute, backwardly directed spines.

Leptynoma sericea (from Engel, 1929: 155-156): Head and thorax of pupa are brownish, abdomen yellowish in colour. Total length varying from 11 to 18 mm. Width of thorax, 2.5 to 3 mm; and that of anal segment, 1.25 to 1.5 mm (fig. 8) [= Fig. 400].

The facialia of the head bear at each side a crest of flat spines — three larger ones on the lower and two smaller ones on the upper part. The sheath of the proboscis reaches almost to the end of that of the fore legs. The thorax is in profile a little hump-backed, and at each side bears three flat, spine-like processes — one above the prothoracic spiracle and two others at base of wings (fig. 9) [= Fig. 401].

The abdomen consists of seven segments, the last of which is but imperfectly divided, and is counted as one. Two lighter brown tergites cover scutellum and metanotum; sternites are wanting for these segments, which are covered ventrally by the sheaths of wings and legs. The first six segments show at each side a long and sharp spine and nearer to the hind margin lies a spiracle, below which is a longitudinal row of shorter and smaller spines. The anal end of the last segment has two strong, ventrally turned hooks, which are always hidden in the adhering larval skin. This bursts in the dorsal median line of the thoracic segments, so that the head-capsule and the whole larval skin are preserved with each pupa.

The whole surface, and especially that of the abdomen, is covered by innumerable sand-grains which are glued to it by a secretion. The integument of the inter-segmental parts of the abdomen is transversally rugose; that of the segments shows densely set, minute spines, all directed anally.

Vermileo comstocki (from Wheeler, 1930: 186, 188): The characters of the pupa (fig. 27) [= Figs. 395-396] are easily seen after the sandgrains adhering to its surface and the last larval skin attached to its posterior end have been removed with caustic potash. It measures 6.5-8 mm and is dark amber brown, with the abdomen paler and more yellowish. The integument of the head and thorax are smooth, but the intersegmental regions of the abdominal segments and their borders and sides are covered with minute, rounded papillae, which seem to represent the regular segmental rows of stiff spinules of other Rhagionid genera (*Rhagio*, *Chrysopilus*, etc.).The last abdominal segment terminates in a pair of short, shaply conical appendages.

Vermileo sp. (= possibly *V. niloticus*) (from Hafez & El-Moursy, 1956b: 345-346): The pupa (fig. 9) is typically orthorrhaphous in form. It measures from 5 to 7 mm in length. The body is somewhat arched dorsally and recurved ventrally. When newly formed, the body is yellowish in colour; after

about one day the head and thorax assume a dark brown colour and the abdomen becomes greyish.

The head and thorax are closely joined to each other and are considerably broader than the abdomen. The large compound eyes are situated within their sheaths (fig. 10) [= Fig. 399] on the antero-lateral regions of the head. The bases of the antennae arise between the eyes. The antennal sheaths are elongated, pointed apically, and extend beyond the posterior margin of the eyes.

The sheaths of the mouthparts occupy a mid-ventral position and extend to the sheath of the fore-legs. The sheaths of the maxillary palps (MX. P.) are widely divergent and separated by the sheath of the labrum epipharynx (L. EP.), maxillae (MX) and labium (LB).

The wing sheaths extend ventrally to the first abdominal segment. The inner margins of the wing sheaths enclose a space occupied by the legs closely folded within their sheaths. The sheaths of the legs (LG) are separated by conspicuous sutures. On the dorso-lateral regions of the thorax, near the anterior end, are located the thoracic spiracles. The metathorax is a narrow strip on the dorsal side of the body.

The abdomen is eight-segmented and tapers towards its posterior end. On the inter-segmented skin are two transverse lines composed of very small roughened areas. Near the antero-lateral regions of the segments 1-7, are situated the abdominal spiracles which are carried on backwardly directed protrusions from the sides of the abdomen.

The skin of the last larval instar remains firmly attached to the last abdominal segment of the pupa throughout the pupal stage.

Vermiophis ganquanensis (from C. Yang, 1979: 89): Pupa (fig. 9) [= Fig. 402]: Length of body 10-15 mm, abdominal segments with spine-like processes from 2nd to 7th segments, and the spiracles of 1st to 7th pairs are borne on distinct tubercles, the last pair is only a slit on the side of 8th abdominal segment. The caudal segment terminates in a pair of large conical closing appendages.

Vermitigris fairchildi (from Wheeler, 1930: 274): The pupa (fig. 47) [= Figs. 397-398] measures 15 mm and is dark brown, with thicker, more brittle and smoother integument than in *Vermileo* and *Lampromyia*. Although the crumpled last larval skin firmly envelops the two posterior segments as in those forms, there is no adherence of sandgrains to the surface. This seems, however, to have been covered during the last larval ecdysis with a whitish but non-sticky secretion. The body is long and narrow, the thoracic dorsum very convex and rounded, the abdominal segments somewhat flattened dorsoventrally. The head has a very broad face and the antennae are widely separated, with long arista and enclosed in free sheaths directed downward along the inner borders of the eyes. The latter seem to be much smaller and less convex than in *Vermileo* and *Lampromyia*. The proboscis is short and stout and, therefore, more like that of *Vermileo* than *Lampromyia*. The caudal segment terminates in a pair of long, slender, curved, diverging appendages. There are no traces of tubercles or spines on the abdominal segments and only the spiracles of the second and third segments are borne on distinct tubercles.Outstanding peculiarities of the pupa are the structure of the antennae, the small size of the eyes and failure to acquire an envelope of sandgrains.

Vermitigris orientalis (from Edwards, 1932: 219): Wheeler describes the integument of the pupa of *V. fairchildi* as being quite smooth, without adhering sandgrains, and with only the second and third pairs of abdominal spiracles set on distinct tubercles; the Malayan pupae have a good deal of fine sand adhering to the integument, and each of the first seven abdominal segments have the spiracle set on a small tubercle, and beyond this on the lateral margin of the segment is a short, pale, spine-like process.

Vermitigris sinensis (from C. Yang, 1988: 182): "Pupa (figs. 2-3) about 15 mm long, with larval skin firmly enveloping the two posterior segments as in the other worm-lion forms. The integument of pupa has adhering sandgrains (no adherence of sandgrains to the surface in *fairchildi*), and the first seven abdominal segments each has the spiracle set on a small tubercle, and beyond this on the lateral margin of the segment is a short spine-like process (in *fairchildi* with only the second and third pairs of abdominal spiracles set on distinct tubercles), make the Chinese species allied to *orientalis*. But the pair of long spine-like processes at the tip of the abdomen is close up at the base (not separated as the other two species), and the spines are very twisted. The form of the widely-separated antennal sheaths seems to be the more chief general character for the pupae of this genus only".

Lampromyia s. lat. (including *Leptynoma*, *Vermilyn*x and *Vermipardus*) are easily separated from *Vermileo*, *Vermiophis* and *Vermitigris* by having a long proboscis.

There is a pair of terminal appendages or processes on last abdominal segment, and probably the form of the above paired appendages represents the most important generic character: *Lampromyia* (or s. lat.): longer than in *Vermileo* and more stout than in *Vermitigris*; *Vermileo*: short and sharply conical; *Vermiophis*: connected with each other, large and conical as a whole; *Vermitigris*: long, thin, widely separated from each other and divergent.

In *Vermitigris*, the antennae are more widely separated from each other than in other genera [in Figs. 397-398, the base of antennae is probably omitted].

More comparison is necessary for finding definite generic diagnosis based on a number of species.

For specific characters of *Vermitigris*, see the description of each species quoted above.

Pupal spiracles

The species, which have been illustrated, are listed firstly.

Leptynoma sericea: Engel (1929: figs. 10-11).

Vermileo comstocki: Greene (1926: fig. 11: u, v); Wheeler (1930: fig. 31: E-H).

Lampromyia canariensis (from Wheeler, 1930: 255): The abdominal spiracles are also on tubercles and are much as in *V. comstocki* but larger.

Leptynoma sericea (from Engel, 1929: 156) (Figs. 393-394): The prothoracic spiracles and those of the abdominal segments are, as in the larva, of different construction. Both spiracles are almost of the same size, for the pupa is always hidden in the sand, but the first is embedded in the strongly chitinized walls of the thorax, while the abdominal one stands on a flat cone. The stigmatic plate of the prothoracic spiracle has a strongly chitinized margin, and shows three to four stigmatic openings. The stigmatic scar lies anally, and a strong muscle, directed head-wards, remains preserved even in the cast pupal skin. The plate of the abdominal spiracle is less chitinized, shows six to eight stigmatic openings, and its short trachea originates from one of the two great lateral longitudinal stems. The stigmatic scar is directed towards the median line of the abdomen.

Vermileo comstocki (from Wheeler, 1930: 186-188): The small spiracles (fig. 31 E to H) (Figs. 387-390) are situated on distinct conical elevations and consist of groups of minute, elongated elliptical slits, radially arranged. They differ in number on the different segments, there being seven in each of the prothoracic spiracles (fig. 31 E) [= Fig. 387], which are the largest, and a gradually diminishing number on the more posterior segments, down to three in the last pair, which is on the eighth abdominal segment (fig. 31 H) [= Fig. 390]

Vermileo sp. (= possibly *V. niloticus*) (from Hafez & El-Moursy, 1956b: 346): Near the antero-lateral regions of the segments 1-7, are situated the abdominal spiracles which are carried on backwardly directed protrusions from the sides of the abdomen.

Vermiophis ganquanensis (from C. Yang, 1979: 89): The spiracles of 1st to 7th pairs are borne on distinct tubercles, the last pair is only a slit on the side of 8th abdominal segment.

Vermitigris fairchildi (from Wheeler, 1930: 274): There are no traces of tubercles or spines on the abdominal segments and only the spiracles of the second and third segments are borne on distinct tubercles.

Vermitigris orientalis (from Edwards, 1932: 219): Each of the first seven abdominal segments has the spiracle set on a small tubercle, and beyond this on the lateral margin of the segment is a short, pale, spine-like process.

Vermitigris sinensis (from C. Yang, 1988: 182): The first seven abdominal segments each has the spiracle set on a small tubercle, and beyond this on the lateral margin of the segment is a short spine-like process (in *fairchildi* with only the second and third pairs of abdominal spiracles set on distinct tubercles), make the Chinese species allied to *orientalis*.

Wheeler (1930: 190) wrote: Greene calls attention to another peculiarity of the pupal spiracles when he says that "the spiracular plate in all the species [of Xylomyidae, Xylophagidae s. lat. and Rhagionidae], except that belonging to the genus *Vermileo*, are very similar, and in all the spiracular entrance, the so-called spiracular opening, closely resembles the script capital letter E."

Egg

The previous descriptions and figures available are quoted below.

Lampromyia pallida (from Hemmingsen, 1968: 297): Eggs laid in June 1961 and preserved in Faester's fluid measured November 1964 1.1×0.7 mm. At high magnification the immediate impression formed of the surface of the chorion are refractive points about 1 micron in diameter arranged in lumps about 3 microns in diameter. The mutual distance of the lumps is about 3 microns, and they are seen to be connected by a reticulum possibly consisting of canals. At certain illuminations the lumps are seen to consist of elevations of the reticulum constituting short rounded papillae. Sand particles stick to the egg surface much as shown for *V. vermileo* by Le Faucheux (1961, fig. 2) [= Fig. 406] .

Vermileo comstocki (from Wheeler, 1930: 215): The pale cream-colored, broadly elliptical eggs measure only 0.6 mm in length and 0.4 mm in width and adhere to one another in rows as shown in fig. 29 H [= Fig. 85]. The chorion, or shell, is thin, transparent and, under a high magnification, finely granular.

Vermileo vermileo (from Hemmingsen, 1968: 297): Eggs protruding from the genital opening or dissected out measured 0.7×0.4 mm, agreeing with the figures 0.67×0.43 mm given as averages for length and greatest width by Le Faucheux (1961, p. 137) . At high magnification the surface of the chorion presented a similar picture to that given by Le Faucheux (1961, fig. 3) viz. rather thin papillae about 5 microns apart connected by straight lines. But the straight lines were generally intersected by others so that an essentially more complicated reticulum was seen to exist in the interpapillary areas than that shown by Le Faucheux.

Vermileo vermileo (from Le Faucheux, 1961: fig. 4): see Fig. 405

Vermileo sp. (= possibly *V. niloticus*) (from Hafez & El-Moursy, 1956b: 339-340): The egg (fig. 4) [= Fig. 412] is oval in shape with the narrow end more or less truncate. It measures from 0.58 to 0.78 mm in length, and 0.30 to 0.40 mm in width. When newly deposited, the egg is more or less transparent and pale creamish in colour. After about three days it becomes opaque and assumes a whitish colour. The egg is always surrounded by scattered fine sand particles stuck to its surface. Usually, this sand is not easily removed even if the egg is immersed in water for many hours. Under high magnification, the micro-sculpture of the egg-shell is neatly seen.

CITATIONS FROM THE LITERATURE ON GENERAL BIOLOGY

The review as given here is far from perfect and the reader should consult the original works, that is, De Geer (1752, 1776), de Réaumur (1753), von Siebold (1861), Marchal (1897), Engel (1929), Wheeler (1930), Buchner (1940), Hafez & El-Moursy (1956a, b, 1964), Le Faucheux (1961), Hemmingsen (1963, 1968, 1977), Hemmingsen & Nielsen (1971), Frederiksen & Hemmingsen (1972), Stuckenberg (1961, 1995b, 1996a), Ludwig *et al.* (1996), etc., of which only some portions are selected. Nevertheless, there are so many long passages quoted here, which we must give in full.

ADULT

1. Sex ratio

It seems to be 1: 1, though records vary.

Wheeler (1930: 255) on *Lampromyia canariensis*: The beautiful dark colored flies were very inactive. They rested with overlapping wings and the head and anterior portion of the thorax elevated, the abdomen lowered and the long, needle-like proboscis directed backward and downward between the legs. Both sexes appeared in about equal numbers, but for some unknown reason showed no inclination to mate.

Hafez & El-Moursy (1956b:335-336) on *Vermileo* sp. from Egypt: It was also observed that the number of emerging females exceeded that of males. Thus, of sixty adults emerging under room conditions, 40 were females and 20 were males.

Le Faucheux (1961) on *Vermileo vermileo*: There were 7 males and 8 females emerged in the laboratory from May 29 to June 15, 1959.

Hemmins (1963: 252) on *Lampromyia fortunata*: Most flies emerged from the middle of May to the end of July, females and males in equal numbers (6 ♀ ♀, 6 ♂ ♂) and with no clear-cut chronological difference.

Hemmingsen (1968: 298): Of 43 *Vermileo vermileo* which emerged in the laboratory 26 were females; 17, males. And of 107 *L. pallida*, 45 were females; 62, males.

2. Feeding habit

Teskey (1981: 531) wrote, "Adults feed on nectar. Members of *Lampromyia* insert a long proboscis into deep flowers, whereas members of *Vermileo*, with a short proboscis, are confined to flat, open umbelliferous flowers", apparently based on Hemmingsen & Nielsen (1971: 152-153) who wrote:

Both sexes of all the *Lampromyia* species [*pallida*, *canariensis*, *fortunata*, *hemmingseni* and

funnebris] dealt with in this paper were seen to insert their proboscis into one deep flower after the other (fig. 3) and they lived up to a week — sometimes even two weeks. The species without a proboscis, *Vermileo nigriventris*, was also seen to suck (from umbelliferous flowers). [The flies mentioned above were] confined in glass-walled containers (fig. 2) usually measuring 77 (length) × 30 (width) × 35 (height) cm.....

Hemmingsen (1963: 252-253) on *Lampromyia fortunata*: The imagines lived, with access to honey-water, from a few days to about a week or even a few days longer. In two flies the very long proboscis characteristic of this species was seen being dipped into the sweet fluid, and the flies were apparently drinking, for the two parts of the flattened tip of the proboscis moved relative to one another.

Frederiksen & Hemmingsen (1972: 58) on *Lampromyia fortunata*: There was a negative correlation [in the field] between number and density of [larval] pits and distance from flowers.

Buchner (1940: 129) on *Vermileo vermileo*: Nun nimmt es den Flug nach der Sonne, lässt sich auf Blüten nieder und nippt gesittet an Tautropfen oder an Süßen Sätzen.

On the other hand, there are some species taking no food in the adult stage.

Vermileo comstocki (from Wheeler, 1930: 214): My flies [emerged from pupae in laboratory] declined to imbibe either water or syrup, and since dissection of females showed that their eggs were fully mature in their ovaries and ready to be laid, it is probable that the short-lived imaginal insect requires no food.

[On p. 213] I was able to complete the life-history of *V. comstocki* with data on its mating, oviposition and earliest larval behavior.

Vermileo sp. (= possibly *V. niloticus*) (from Hafez & El-Moursy, 1956b: 337-338): "The adults persistently avoided liquid food with which they were provided, such as syrup, sugar solutions or honey. They did not even show any tendency to sip nectar from flowers when offered to them.Dissection of the abdomen of the female, just after emergence, revealed that the ovaries contained mature eggs. The pre-oviposition period, i. e. the time elapsing from emergence to the beginning of oviposition, usually lasts from about four and a half hours to about seven hours. Shortly after mating, the female rests on the surface of the sand for one or two hours after which it begins to lay eggs."

There may be no flower-visiting record in the open field (not in the laboratory). Stuckenberg (1960: 224; 1995b: 239) wrote: I have been unable to find any references to the feeding habits of the imago in *Lampromyia* [s. lat.] although the biology of the immature stages is so well known.Hardly anything is known about the biology of *Vermipardus* flies. Their elongate proboscis appears to be adapted for nectar-feeding; evidence for this is seen in the occasional presence of pollen grains on pinned specimens, but I am not aware of any records of these flies being seen on flowers.

Stuckenberg (1996a: 200) also wrote, "The Goegap Nature Reserve, where this species [= *Vermilynx jasoni*] was found, was the site of a study of pollination ecology by Struck (1992).Struck collected all insect pollinators visiting selected plant species during his three-year project, but he records no vermilionid flies. Two possible reasons exist: either his fieldwork, which collectively covered the months of late June to late November, terminated before the flies appeared (the type specimens emerged in January); or, the flies may have been present, but were not noticed because they feed at uncommon or inconspicuous plants that were not included in Struck's study. The known adult phenology of *V. vansoni* in the equally arid Richtersveld covers the entire spring and

summer period of September to April, so *V. jasoni* may have a similarly extended period of adult emergence".

The fact above may simply be ascribed to the scarcity of adult vermilionids.

3. Swarming

Swarming was observed in *Lampromyia fortunata* by Frederiksen & Hemmingsen (1972: 53) who wrote: Flies on the wing were seen outside the caves from 17°C on overcast days. The erratic horizontal flight within a small space, characteristic of the swarming of many insects, was seen among plants in one or a few males at 28°C in the free (as also under similar conditions in the containers in the laboratory).

4. Copulation

There are three postures of copulation: (a) male lies on the back of female; (b) male and female take an opposite direction with their abdomens normal in position; (c) male and female take (b)-posture but with male abdomen turned upside down and legs hanging in the air. All of the (a), (a+b) and (a+b+c) above may be seen at least in some species according to the individual pair.

Vermileo sp. (= possibly *V. niloticus*) (from Hafez & El-Moursy, 1956b: 337-338): Under laboratory conditions, it was noticed that adult males and females which emerged about the middle of March did not show any inclination to mate. Mating only occurred between adults which emerged towards the end of March and later, when the temperatures were higher, by about 3-4°C, at the end of this month rather than in the middle of it. This was true for the years 1952, 1953 and 1954. On the other hand, the relative humidity did not show any regular variation.

Mating took place 2 to 6 hours after emergence. The male first hovers around the female for some time, then approaches her and comes to rest over her back curving its own abdomen downwards around the abdomen of the female. The male then grasps the ovipositor of the female from below with its forceps and for this posture it raises its hind legs while at the same time the female inclines the apex of its abdomen slightly towards the male apparatus. The couple remains in this posture for about ten minutes, probably to ensure a complete hold in the right position. Afterwards, the male instead of remaining on the back of the female, rotates horizontally about 180° so as to lie in line with the female (fig. 3).

Mating lasted from about 30 to 150 minutes. During this period the couple, unless disturbed, remains completely motionless on the surface of the sand or on the sides of the breeding cage.

During the process of mating, the wings of either sex form an acute angle with the long axis of the body. In some cases, however, the wings of the mating individuals may overlap during the process. Disturbance does not interrupt copulation, but merely causes the male and the female to fly while attached to each other. In this case the female usually takes the lead.

After copulation, the two sexes separate and the male flies actively in the air while the female rests for some time before starting to oviposit. It was noticed that the male, unlike the female, showed a certain tendency for repeated copulation.

Vermileo comstocki (from Wheeler, 1930: 214): On the day following emergence several pairs of flies mated. During copulation, which lasted about half an hour, both sexes spread their wings somewhat, the male rested on the back of the female, with his abdomen parallel with hers, but with its posterior segments twisted to the left side and the genitalia at the tip turned upward and firmly applied

to those of his mate.A few hours after mating several of the females oviposited.

Hemmingsen & Nielsen (1971: 160): In *Vermileo vermileo* from Italy and *V. nigriventris* from Spain the copulating pairs observed were hanging vertically as pictured by Buchner (1940) and Hafez & El-Moursy (1956b) and as in fig. 5a [= (b)], and lasted from at least 1/4 to 1 3/4 hours. The copulations in the *Lampromyia* species were as in figs. 4 [= (a)] or 5a-b [= (b) and (c)]. They lasted in *L. pallida* up to 3/4 hour; and in the other *Lampromyia* species [= *canariensis*, *fortunata*, *hemmingseni* and *funnebris*], from a few to 15 minutes.

5. Oviposition site in the field

Hemmingsen & Nielsen (1971: 202): [About] 50 ovipositions of *Lampromyia fortunata* were studied throughout May 1971 in the caves of Cenobia deValeron, Gran Canaria, by A. M. H. jointly with Mr. Jørgen Frederiksen. The ovipositions were seen only in dust and in the shade (385-9000, usually 1600-3300, lux), at 18-21.5°C and relative humidities of 50-70 %. As the Canarian species are largely inactive below 18° these observations imply that this species does not consistently oviposit at higher temperatures than *L. canariensis* (cf. p. 196).

Frederiksen & Hemmingsen (1972: 57) on *Lampromyia fortunata*: Studies in the cave complex Cenóbio de Valerón, Gran Canaria, showed a positive correlation between luminosity and number and density of larval pits; with especially few or no larvae in caves into which the sun never penetrated.

6. Oviposition

Hafez & El-Moursy (1956b: 338-339) on *Vermileo* sp. (= possibly *V. niloticus*) (in laboratory conditions): Shortly after mating, the female rests on the surface of the sand for one or two hours after which it begins to lay eggs.

Before oviposition, the female hovers near the surface of the sand while testing it with the hind-tibiae. It then fixes itself to the sand with its fore and hind legs, and digs with its middle legs a small pit of about one millimetre in diameter and two millimetres in depth. At the same time, the female curves its abdomen downwards and anteriorly. During all these proceedings the abdomen vibrates rapidly, probably to help the eggs moving singly from the ovary towards the tip of the abdomen where they lie ready for deposition. Now the female thrusts an egg into the pit by a rapid stroke and, with the help of the fore and middle legs, it fills the pit with sand so as to hide the egg.

The place where an egg is laid is very typical, and assumes the shape of a small heap of sand with a depression in the centre surrounded by the clear tracks left by the legs of the female.

After an egg has been deposited, the female rests for a while (5-20 minutes) either on the sand or on the sides of the breeding cage. While resting, it scratches the extremity of its abdomen several times as if it were doing some massage before the next egg is laid.

The process of egg laying continues for a period ranging between 30 minutes to about four hours, after which the female remains motionless for several hours (about 4 to 19 hours). This latter period is the post-oviposition period.

The number of fertilised eggs laid by a single female varies from 8 to 30. Unfertilised eggs are uncommon, and when laid by unmated females they shrink after about three days.

Le Fauchaux (1961) on *Vermileo vermileo* (in laboratory conditions): The oviposition habit of *Vermileo vermileo* from France differed from that of *Vermileo* sp. from Egypt in the following

Fig. 403. Flying (A, C), testing (B) and landing (D) for the egg-laying in *Vermileo vermileo*. From Le Fauchaux (1961).

Fig. 404. *Vermiophis yanshanensis*. A sketch of ovipositing female. Refer to p. 129. From C. Yang & Chen (1993).

respects: the female touches or tests the sand with the mid legs (not hind legs) during flight (see Fig. 403: B); no leg is used either for pit-digging or for egg-covering before and after oviposition; no heap of sand is seen in the oviposition site.

Vermileo vermileo repeated 2-6 times two movements (see B and A or C in Fig. 403; A and C, about 15 cm high above the ground), then sat on the sand and commenced egg-laying (see D in Fig. 403).

Oviposition behaviour was analysed in detail chiefly under laboratory conditions by Hemmingsen & Nielsen (1971) in the following 2 genera and 6 species: *Lampromyia* (*pallida*, *canariensis*, *fortunata*, *hemmingseni* and *funbris*); *Vermileo* (*nigriventris*). Some of the results are introduced below.

There are 3 phases: (1) pit-digging and egg-laying; (2) egg-covering; (3) pit-covering. The distinction between (2) and (3) is difficult and united into one phase except for *L. pallida* and some individuals of *L. hemmingseni*. The duration of whole ovipository process was 1.5-4.0 minutes in *L. pallida*, 0.1-2.2 in other 4 species of *Lampromyia*, and 0.05 in *V. nigriventris*. For the duration of each phase, see their table 6.

(On p.199): All the 6 species of Vermileonidae studied dig a pit in loose substrate with the simultaneously backward moved pairs of middle legs and lay an egg in the pit [Phase 1].

(On p.193): Usually the insertion of the abdomen into the soil is quick; the withdrawal, which presumably represents the actual extrusion of the egg, slightly slower.

(On p.198): In *L. pallida* there is an egg covering phase 2 and a pit-refilling phase 3, both carried out by the middle legs only. They are, so to speak, merged into one egg-covering and pit-refilling

Figs. 405-411. *Vermileo vermileo*. 405, egg; 406, egg (after hatching) with sand particles; 407, magnified part of egg chorion; 408-411, process of hatching. Refer to pp. 91, 114. From Le Fauchaux (1961).

Fig. 412. Egg of *Vermileo* sp. from Egypt (= possibly *V. niloticus*). Refer to p. 92. From Hafez & El-Moursy (1956b).

phase in the other species, which use both fore and middle legs.

(On p.199): In *L. fortunata* the fore and middle legs may make some jerks (phase 3 [which is not homologous with that of *L. pallida*]) after the pit has been refilled. In *L. hemmingseni* the jerks often constitute part of phase 2.

(On p.200): During phase 1 the downward bent apex of the abdomen is thrust rhythmically forward in the air in *L. pallida* (synchronously with the digging movements), *L. fortunata* and *L. hemmingseni* (in the latter two species less frequently than the digging movements, which are quicker than in *L. pallida*). In *L. canariensis* the thrusts and the digging movements are still quicker, and the thrusts appear to attempt in vain to follow the digging movements so as to apparently cease at all.

Whether or not such thrusts occur in *L. funebris* and *Vermileo nigriventris* was not noted.

(On p.200): In *L. canariensis* and *L. hemmingseni* the female starts digging immediately after settling from interovipository flight. In *L. canariensis* it flies away immediately after phase 2 without exhibiting any jerks with fore and middle legs (with rare exceptions). In *L. fortunata* the fly does not start digging until a short while after settling, and in this species and *L. hemmingseni* she stays on for a while after phase 2 (or 3 when present).

(On p.200): In *L. canariensis* the female quivers her wings throughout the whole process; in *L. fortunata*, only at the egg-laying proper at the conclusion of phase 1 and during phase 2; and in *L. hemmingseni*, during the first part of phase 1. Exceptions occur.

7. Number of eggs per one ovipository site

It is one in 6 species of *Lampromyia* and *Vermileo nigriventris* studied by Hemmingsen & Nielsen (1971:164) who wrote: After a little rest the fly [= *L. pallida*] flies about, for instance for [2/100 minutes] or sometimes more, to settle on another spot and start digging again. This is repeated in different places, one egg being laid each time.

It is also one in *Vermileo vermileo* by Le Fauchaux (1961) and *Vermileo* sp. from Egypt by Hafez & El-Moursy (1956b).

On the other hand, Wheeler (1930: 215) wrote as to *Vermileo comstocki*: A few hours after mating several of the females oviposited [in the laboratory].All the eggs, more than fifty in number, are extruded at one time, and the female dies soon after withdrawing the tip of her abdomen. Of the eggs laid by the females in 1917 only one batch hatched, but in 1930 many batches of fertile eggs were obtained,The pale cream colored, broadly elliptical eggs measure only 0.6 mm in length and 0.4 mm in width and adhere to one another in rows as shown in fig. 29 H [= Fig. 85].

8. Number of eggs produced by one female throughout her life

There is no mention of it by Hemmingsen & Nielsen (1971) concerning 6 species who took food. *Vermileo* sp. from Egypt and *Vermileo comstocki* refused food and had short longevity in the laboratory, but this may possibly be so in the field too in these two species.

It varied with individual from 8 to 30 in *Vermileo* sp. from Egypt (after Hafez & El-Moursy, 1956b: 339) and over 50 in *V. comstocki* (Wheeler, 1930: 215).

9. Sleeping posture

This was noticed in some *Lampromyia* species by Hemmingsen & Nielsen (1971: 168) who wrote: The peculiar 'sleeping posture' of this [= *L. canariensis*] and other species is shown in fig. 10 [where the fly rests perpendicularly on its head and all legs touching upon substratum]. It is uncertain, however, whether this is an entirely normal phenomenon.

10. Dispersal in the field

Nobody knows the actual distance of flight from breeding site to a new sphere where the next generation in some individuals is established.

Vermileo sp. from Egypt (which seems to take no food and has short longevity) (from Hafez & El-Moursy, 1956 b: 336-337): One may assume that they generally do not go far from their breeding

sites. This may be supported by the following. (1) The adults are delicate and unable to fly for long distances. (2) Being short-lived and produced in small numbers, the dispersal of the adults in the wadi [in Egypt] may decrease the chances for both sexes to come across each other for mating. (3) If fertilized females go far from their breeding places and do not succeed in exploring suitable shaded situations for oviposition and for development of the future larvae they will be completely destroyed. The eggs and pupae, as shown by laboratory experiments, can withstand temperature not more than 30-32°C, while the surface temperature of the sand in exposed areas reaches on the average 45°C. (4) Differences between physical conditions in the shade and in the open may act as barriers preventing the wandering of the adults.

During flight, the male usually curves its abdomen downwards while the female extends it. In the laboratory it was observed that both sexes are positively phototactic as they generally congregate near the illuminated side of the breeding cage. At rest the wings lie on the abdomen, either overlapping or widely separated.

Vermipardus species from southern Africa (which seem to visit flowers) (from Stuckenberg, 1955b: 239, 242): Reared captive flies show a restless, energetic behaviour as soon as firming of the exoskeleton and wings has been completed after eclosion. This may be evidence of a dispersal phase that would occur under natural conditions, when mates are located away from the site of larval development, and when oviposition may occur. The scattered nature of larval colonies suggests that such a dispersal episode probably occurs, though it is also probable that some flies oviposit in the same site where they passed through their immature stages. (P. 239)

The flies [= *V. sylphe*] apparently move from one forest patch to another, in search of the sporadic and isolated sites that meet the special needs of their larvae. (P. 242)

LARVA

1. Breeding site

This topic is summarized by Hemmingsen (1977: 186): Caves are common habitats of the worm-lions, because disintegration products in the form of fine sand or dust, partly derived from the cave walls and ceiling, and partly carried by the wind, collect and settle there. Other worm-lion localities, e. g. the foot of palms or walls exhibit similar sheltering properties.

It is described in detail by Hafez & El-Moursy (1956a: 284, 286) on *Vermileo* sp. from Egypt: The larvae live in pits in the sand (fig. 2), and are usually aggregated in groups commonly known as colonies. These colonies are scattered all over the wadi, but they are dense and more numerous in certain localities.

The localities where *Vermileo* colonies are found are very characteristic in shape, structure, direction and sand consistency. It was noticed that the sand harbouring these colonies is mainly of fine nature.

Larval pits are generally found under overhanging cliffs (figs. 1 & 3) that have a northern or north-western direction. They are not found under rocks with southern direction. This may be due to the fact that in the first position the larval pits are always in the shade sheltered from the direct unfavourable effect of the sun. Furthermore, the rocks protect the pits from rain and wind. This is supported by the fact that although a heavy rain fell on the wadi in December 1951, the sand in which the larvae constructed their pits was completely dry, while the rest of the wadi was wet. It seems also

for this reason that the larval pits are usually found on steps of rocks (fig. 3) at a level about one metre from the bottom of the wadi, or close to the bases of cliffs beside pieces of stones or large solid objects, or even in cavities of big fallen stones that happened to be filled with fine sand. They also construct their colonies within large loops of the chain of rocks (fig. 4). All these situations seem to provide protection for larvae from the sun, heat, accumulated rain, and also from the direct effect of wind. These protective measures against adverse physical conditions seem to be a necessity, as the larvae are completely dependent upon their pits in obtaining their food supply.

Sometimes the larval pits occur among patchy vegetation as shown in figure 2.

It is also described by Stuckenberg (1995a: 239-242) on *Vermipardus*, an inhabitant of forest: The requirements of the larvae resemble those of other vermilionid genera, especially in that these should be dry, loose, smoothly-granulated sand or sandy soil, even of soft, dust-like or powdery fineness, for their pit-making. Although this suggests that *Vermipardus* species may favour arid environments where such material tends to be common, the known distribution of this genus indicates that these flies actually require breeding sites that are humid and shaded. The species are all recorded from montane areas or from topographically diversified terrain, in regions of relatively high rainfall. The larval habitat is always under some sort of shelter — typically the dry sandy floor under a rocky overhang or in a cave, or in a recess in an earthen bank. I have never found *Vermipardus* larvae under the sky or in a situation where they are openly exposed to rain. There is also a preference for such sites to be well shaded, particularly as a result of aspect (south-facing especially) and the presence of sheltering vegetation.

Sites with such a combination of features may be relatively uncommon and scattered in any locality. This became apparent during a search I made along more than one kilometre of both banks of the Rensburgspruit, a river in valley-bushveld near Estcourt in Natal. Although numerous recesses were found in the banks, nearly all were occupied by antlions (Neuroptera, Myrmeleontidae). Only one small wormlion population (about 20 pits of *brevirostris*) could be found; this had exclusive occupation of a sandy recess in a south-east facing bank at about one metre above ground-level, in a distinctive site where a thicket of trees and bushes provided exceptionally continuous and complete shade.

An examination of larval populations of *sylphe* proved particularly instructive. The type-locality, Fairy Glen in the Royal Natal National Park, is a small patch of quite dense *Podocarpus* forest in a little valley surrounded by extensive grassland. Within this forest, the larvae were restricted to a permanently shaded, quite dark recess beneath a small sandstone cliff, adjacent to a permanent mountain stream, where the air was continuously cool and humid. A search for this species was also made several kilometres away, along a path traversing a narrow strip of forest above a bank of the Thukela River below Tendele Camp. This forest is enclosed by extensive grassland. A solitary cluster of three larvae in a saucer-sized niche was found in the earthen bank bordering the path; recesses at ground-level all contained antlions. The flies apparently move from one forest patch to another, in search of the sporadic and isolated sites that meet the special needs of their larvae.

The following fact should be noted and is quoted from Hemmingsen (1968: 296-297): "The tendency to excavate the pits near the side of the containers, as illustrated for *Vermileo comstocki* by Wheeler (1930, fig. 39 on p. 225) and for the Gran Canarian species [= *Lampromyia fortunata*] by Hemmingsen (1963, fig. 5 on p. 243), was seen in *Vermileo vermileo* and *Lampromyia pallida*. As noted also for other species by other authors, in nature the pits are similarly placed next to walls,

rocks, etc. For discussions and explanations cf. e. g. Wheeler (1930, pp. 215, 224-226), Buchner (1940, p. 118), and Hafez & El-Moursy (1956a, pp. 285-286)".

There are exceptional open habitats which must be introduced.

Vermileo comstocki (from Wheeler, 1930): Here [= Alta Meadow, 9000 ft, high Sierras of California] were great numbers of *Vermileo* pits in stretches of very fine glacial silt and dust surrounding the scattered boulders. Though the larvae had made their pits close to the edges of the boulders, they were so deeply embedded in the silt that their rounded and receding sides afforded the insects no protection from the wind, rain and sun. [On p. 170]

Thus in Alta Meadow at 9000 ft. the pits are not sheltered under the overhanging rocks but in the open sand on silt where they are fully exposed to the elements, whereas at 4000-5000 ft. in the Yosemite they are always in dust which is rarely or never wetted by rain or for long periods exposed to the rays of the sun. It is difficult to account for this ecological discrepancy except on the supposition that *V. comstocki* is not strongly thermotactic and is therefore best adapted to the lower temperatures of higher altitudes. This would explain its preference for the shadier and cooler patches of dust along the often very warm borders of the valley floor in the Yosemite. In all cases, however, the *Vermileo* larvae, unlike those of *Myrmeleon*, have an unmistakable penchant for making their pits very near, though not often in contact with, boulders or cliffs. [On p. 174-175]

Lampromyia hemmingseni (from Hemmingsen & Nielsen, 1971: 196): In Fuerteventura [Hemmingsen] found the larvae commonly near walls and fences entirely in the open. Presumably in this island they are not so much in need of shelter from rain, because it sometimes does not rain for a whole year. Wet substances are too hard for the larvae to work in.

2. Discrepancy in breeding site between wormlions (Vermilionidae) and antlions (Myrmeleontidae)

It should be noted that the pit makers belong to a minority within Myrmeleontidae (Neuroptera) and of them dwellers in the shade are further limited to a small number of taxa, with which the wormlions must compete. For example, there are 8 genera and 17 species of Myrmeleontidae in Japan where 2 genera and 5 species (of which 4 belong to *Myrmeleon* and 1 to *Hagenomyia*) have been confirmed as pit makers, of which only *Hagenomyia micans* prefers shady places protected from rain and direct sun (after Matsura, 1989, 1997).

The essence of discrepancy is summed up by Hafez & El-Moursy (1963: 127): The micro-climate of both groups differs considerably. Thus, since the myrmeleontids construct their pits mostly in the open, they are subjected to a wide range of environmental fluctuations. Studies on the Egyptian form has shown that the temperature at a depth of 4 cm where the larvae lie in their pits reaches a maximum of 44.4°C, and a minimum of 6.0°C; while in the case of *Vermileo* [sp. from Egypt] the highest temperature recorded was 29.4°C and the lowest was 12.8°C.

"Rain greatly affects myrmeleontid larvae by inhibiting their activity during the rainy season which lasts about four months every year. On the other hand *Vermileo* larvae are completely protected from rain, hence they are active throughout the year".

There are a scattering of notes in the five following quotations from Wheeler (19330) saying that both groups are seen in the same place.

Leptynoma sericea (a letter from Hans Brauns to Wheeler): Sometimes there are funnels of

antlions among the [*Leptynoma*] colonies. [On p. 259]

Vermileo comstocki: In other localities, especially near Mirror Lake, the *Vermileo* pits were mingled with those of antlions, but the latter were always very small. As a rule, the large larvae of the common species of *Myrmeleon* in the Yosemite make their pits in open sunny places. [On p. 174]

Vermileo dowi: Mr. R. P. Dow informs me that he found the larvae in fine dust under overhanging limestone cliffs. In some places their pits were mingled with those of small antlions, a number of which he brought me in separate vials of sand. [On p. 276]

Vermileo vermileo: The wormlions and antlions therefore establish themselves at the foundations of old walls or at the base of certain rocks where they have overhanging or almost horizontal projections. Such spots afford a kind of extensive shelter which protects the sand or loose earth from the rain. The antlion pits commonly occupy the first situations, that is, the outer portions of these little caves, whereas the pit of the wormlions are in the recesses. The conical burrows of the wormlions are distinguished by having a greater depth than those of the antlion, with the same diameter. The apertures of the largest wormlion pits, however, scarcely attain the diameter of those made by half-grown antlions. [On pp. 319-320; translated by Wheeler from de Réaumur, 1753]

Vermitegrus fairchildi: "The larval pit falls were interspersed with those of a small species of antlion and were difficult to distinguish because they were only slightly smaller". [On p. 270]

The key particulars in the notes above will be 'small species of antlions' or [antlions occupy] 'outer portions of these little caves.'

An important account is given by Stuckenberg (1995b: 243-244) on *Vermipardus*: Certain myrmeleontid taxa (in South Africa, species of the genera *Hagenomyia*, *Cueta* and *Myrmeleon* — Mansell 1985) also make pitfall-traps in sand.During my search for wormlions, I sifted out the inhabitants of numerous pit aggregations but almost invariably found only antlions. My experience in Natal has been that the great majority of pits at ground-level in sheltered sites are occupied by antlions. At Fairy Glen there was a particularly interesting situation: larvae of *Vermipardus sylphe* occurred only in pockets of fine sand on small rock-ledges about one metre above ground-level. On these ledges no antlions were found, but they were numerous at ground-level below the ledges, wherever there were patches of sand. Spatial separation was clearly evident in the two occupation levels.

The other Natal Drakensberg species, *Vermipardus univittatus*, differs in utilising accumulations of fine sand at ground-level under rocky overhangs or in caves, provided there is continuous shade. In the riverine forest near Main Cave in the Giant's Castle Reserve, the floor of a large sloping recess, beneath a massive boulder close to the river, is known by me to have been occupied by a substantial larval population of this species for at least 40 years. Much smaller colonies of larvae occur sporadically under sloping rocks nearby within the forest; one colony occupies powdery soil in a gloomy, [highly] humid recess at the side of a waterfall in a small ravine. No pit-building antlions have been found yet in this forest. I did find, however, among the organic debris sampled from the large larval colony of *univittatus*, a very small live myrmeleontid larva. This antlion proved to be a species that does not construct pits; Dr. M. Mansell (*in litt.*) has identified it as *Tricholeon hirtellus* Eshen-Petersen, which he informs me is common in sand accumulations on rocky ledges in the Drakensberg. If more of these antlions are present, presumably they have to live near the surface between the wormlion pits, and will be in competition for space and prey in the shelter.

A different situation was found at the Krantzklouf Nature Reserve near Durban. A search I made of a thickly-wooded gorge revealed a large recess, about 20 m long, in the base of a horizontal sandstone ridge on the bank of a permanent stream. The floor of this recess contained numerous pits irregularly clustered relative to the depth and distribution of sand. Most of the pits were occupied by antlions; only in one small area was there a cluster of pits inhabited by wormlions (species unknown, not yet metamorphosed). Several small caves and overhangs nearby had only antlions present.

"These preliminary observations raise certain questions: 1) Are differences in larval requirements the cause of the apparent rarity of sites shared by wormlions and antlions? 2) Are wormlions excluded from suitable sites by the presence of antlions? 3) Does predation by antlions on wormlions occur, and is it a limiting factor? These are moot points, and further study is warranted. The case of *sylphe* could be an instance of resource-partitioning whereby predation is avoided".

It is unknown, however, which is larger in fighting strength, antlions or wormlions, apart from a supposition that owner of the trap tends to have an advantage irrespective of kind. There has been no investigations for the same or various sized individuals between both groups.

3. Prey

This is discussed at length by Stuckenberg (1995b: 242-243): Are the larvae unselective predators? The old European literature accords *Vermileo* a reputation as a consumer of ants, and there is a belief prevalent that ants feature in the diet of vermilionids in general. Wheeler (1930) considered that the *Vermileo* species in California probably consume mostly ants; he also recorded rearing *Vermileo* and *Lampromyia* larvae on worker termites, aphids, and small caterpillars, as well as ants. Hemmingsen (1968) reared larvae of both genera on ants; other authors have resorted to flies as larval food, including houseflies, calliphorids, drosophilids and mosquitoes. I recorded a larva of *Vermipardus promontorii* overcoming a polydesmid millipede 30 mm long, and also *Dermestes* beetles 6 mm long (Stuckenberg 1961). *Vermipardus* larvae which I am rearing at the time of writing are being fed mostly houseflies; small spiders, isopods, and termite alates and workers, have also been fed to them successfully. Such records suggest that wormlions may consume any arthropod that falls into their pits, provided it is not too large, too heavily sclerotised, well-protected chemically, or formidably aggressive. Nevertheless, the unusual microhabitat occupied by *Vermipardus* larvae must limit the range of potential prey.

I was once surprised to find some empty carcasses of muscid flies next to pits of *univittatus* in the large larval colony at Giant's Castle (described below); Hemmingsen (1968) noted this, but it was not certain that those flies had been prey. In an attempt to get more information, I revisited the site recently and removed about 125 cc of the fine, dusty sand in which many successive generations of wormlions had lived. The sample was shaken up in water, and the abundant debris that floated to the surface was skimmed off, dried, and examined under a stereoscopic microscope. Most of it was old, comminuted plant material.

The following arthropod remains were found: Coleoptera — head of a small elaterid-like beetle; fragments of a small beetle; hindleg from chrysomelid-like beetle; two incomplete carcasses of a mordellid-like species; thorax and leg-parts of a small, metallic-green beetle; pale orange elytra from at least six individuals of a small species of beetle. Hymenoptera — ant head of medium size; four abdomens and various legs of a small species of black ant; mandible of larger ant; small hindwing.

Homoptera — carcase of small cicadellid-like nymph; small hindwing. Diptera — carcase of black sepsid fly; part of wing. Arachnida — carcase of immature ixodid tick; carcase of large terrestrial mite; part of spider's leg. Miscellaneous — some very small brown insect femora; indeterminate fragments of insect cuticle.

If these remains represent prey, and if they constitute a fair sample, the following conclusions can be drawn: 1) The remains are from small arthropods, which is to be expected as the prey must be of a size within the capacity of the wormlions in different instars to overcome them. 2) Coleopteran remains are more numerous and diverse than those of any other insect order; in part, this could reflect the suitability of the surrounding forest as a habitat for many small species of beetles. 3) Ants constitute part of the food resource, but in that montane forest, at an altitude of about 1675 m, ants are inconspicuous and cryptic so may not commonly be available as prey. 4) Opportunism is suggested by the diverse arachnid, dipteran and homopteran remains. The data suggest that these wormlions are indeed impartial predators of small arthropods.

Stuckenberg (1961: 123-124) on *Vermipardus promontorii*: The larvae do not react to light or movement, but are sensitive to vibration to which they respond by retreating beneath the sand. While awaiting prey they frequently lie with the anterior third or so of the body exposed in the bottom of the pit. If a potential victim tries to escape by climbing the sides of the pit, the larva flicks sand out from beneath, causing it to tumble in again. Large insects, or ones too hard to be pierced, are flicked out of the pit, and the remains of prey are ejected in the same way. The larvae are unspecific about food and will feed on any reasonably small, non-toxic insect. They have been fed mainly on the small, black, household ant, *Acantholepis capensis* Mayr, which they capture and consume with ease. That they can deal with larger prey has been clearly demonstrated. A polydesmid millipede approximately 3 cm long was attacked and sucked out, and two specimens of a beetle, *Dermestes* sp., about 6 mm long, were taken also. Flies as large as *Chrysomya* spp. (Calliphoridae) have been readily accepted provided they were rendered incapable of flight.

Ludwig *et al.* (1996: 10) on *Verilleo vermileo*: Die Larven wurden im Juni 1994 in Rovinj (Kroatien) und in Oktober desselben Jahres auf Giglio (Italien) gesammelt. Gehalten wurden sie in einem flachen Karton, der mit feinem Sand von ihrem Fundort gefüllt war. Als Nahrung dienten vor allem Ameisen mittlerer Grösse. Auch andere Beuteinsekten wurden akzeptiert, beispielsweise konnten kleine Wurmlöwen unter Laborbedingungen erfolgreich mit Springschwänzen (Collembola), Doppelschwänzen (Diplura) der Gattung *Campodea*, mit Milben (Acarina) oder auch kleinen Fliegen (z. B. *Drosophila*) gefüttert werden. Ältere Larven frassen auch Tenebrionidenlarven (Reismehlkäfer, *Tribolium confusum*). Fütterungsversuche mit adulten *Tribolium* waren dagegen erfolglos, da deren Integument offenbar zu hart ist.

Some notes are gathered from Wheeler (1930): Older wormlions [= *Vermileo comstocki*], like the very young individuals previously described, reject heavily chitinized insects, such as beetles and Myrmicine ants, and insects with powerful odors (*Acanthomyops* and certain bugs). [On p. 234]

They [= *Vermileo vermileo*] declined small bug-like insects and small hard-shelled beetles, the former probably on account of their disagreeable odor, the latter because of their hard chitinous integument. [On pp. 150-151; originally from von Siebold (1861)]

But not all insects that fall into the worm-lion's [= *Vermileo vermileo*] funnel become victims; for some are its superiors in strength. I have occasionally seen flies, whose wings I had removed, and

large species of ants escape, and sometimes even insects that are not its superiors in strength have this good fortune [on p. 329; translated from de Réaumur (1753)]. These worms [= *Vermileo vermileo*] are as sensitive as the antlions and never attempt to feed on an insect completely deprived of life, though it may die in an instant after seizure. But if the insect placed in the funnel is still able to make a few weak movements, it will stimulate the worm to attack [on p. 330; translated from de Réaumur (1753)].

Further two notes are added: Engel (1929: 153) on *Leptynoma sericea*: Small spiders are not attacked at all; they are either frightened by charges of sand, or the larva retreats instantaneously into the sand. Hemmingsen (1963: 250) on *Lampromyia fortunata*: Both small and larger ants were sometimes refused, the larva disappearing into the sand after a few strokes at the ant.

4. Cannibalism

This actually occurs as indicated below, but its frequency in nature is a question.

Hafez & El-Moursy (1956a: 291) on *Vermileo* sp. from Egypt: As regards the enemies of the wormlion in its natural habitat, it is probable that it suffers from cannibalism, especially as in nature the larvae are aggregated in colonies and are thus within the reach of each other. As was shown from laboratory experiments, a larva that happens to fall into another's pit, a fight occurs between the two predaceous creatures. The victim, in trying to escape, leaps for several times in the air but may be unable to avoid falling away from its neighbour's pit. In this case, the other larva gets a good hold of the sand, tries at first to drive it outside the pit but then captures it, paralyses it, and finally feeds upon it. Even pupae may be eaten by the larvae if they come across them. Also the adults, which during the next hour after emergence are soft-bodied, sluggish and unable to fly, may unfortunately fall in the larval pits and are then devoured by the larvae. With the exception of this voracious nature of its own species, no other enemies of *Vermileo* were observed.

On the other hand, there is a negative observation by Wheeler (1930: 220) on *Vermileo comstocki*: "Nor could any cannibalistic proclivities be detected among the young *comstocki* larvae. When one of them happened to fall into a neighbor's pit, it was not attacked but promptly ejected, as if it were a pebble or a bit of exhausted prey".

If the rate of cannibalism is high as a rule, the colony must be affected.

5. Various postures and movements

It should be noted at first that the larva either rests or moves with upside down (i. e. dorsal surface down). This is the fundamental posture of the vermilionid larva.

A series of the latest studies on them are Hemmingsen (1963, 1968, 1977) from which the following (sections 5a-5h) are quoted.

In spite of discrepant reports of previous authors regarding the behaviour of the larvae of [Vermilionidae] which make antlion-like sand or dust traps, no differences were found in postures, movements, pit construction, prey capture and method of throwing out carcasses and foreign bodies from the pit, in larvae of the three species, *Vermileo vermileo* (L.), *Lampromyia fortunata* Stuckenberg and *L. pallida* Macquart, from Italy, Gran Canaria and Tunisia, respectively. [1977: 186]

5a. Pit construction

Also the construction or reconstruction of the sand traps were essentially the same in all three of the

above species: The larva lies with its ventral side uppermost. About 7 segments are visible; the rest of the body is in the sand. The foremost segments with the head bend alternately the left and to the right, almost so far as to reach the body, and with the *side of the head* sand is thrown up. Usually 1-3 thrusts are made in the air to one side, then a number of throwings of sand, 1-11 have been recorded, to the same side; and then the same is repeated to the other side (several larvae observed in each species). Sometimes the thrusts in the air may be omitted; and the behaviour may vary, esp. the number of thrusts. But the procedure is essentially the same in all three species. A more or less circular circumference of the bottom of the trap is the result. [1968: 293].The discrepant descriptions of pit construction in these species as quoted by earlier authors can be due only to superficial observation and wrong interpretation. [1968: 294]

5b. Watch and wait attitude

Finally, the foremost part of the body is brought to rest across and above the bottom of the pit, curved in an upwardly slightly concave arc and slightly covered with sand particles. The head is sometimes seen to be slightly lifted from the sand wall. The head is at least usually the highest end of the concave arc.the abdomen was found to point obliquely downwards, i. e. neither vertically as described by Marchal for *L. pallida* or horizontally as described by Wheeler for *V. comstocki* and *L. canariensis*. [1968: 294].The foremost part rests as a diagonal bridge across the bottom of the pit; the concealed part pointing obliquely downwards. The foremost part is coiled horizontally only when the larva is disturbed, but this does not affect the position of the concealed part. More vertical or horizontal positions of the concealed part are seen only when the larva has been caused to move during the removal of sand [by human power]. [1968: 295]

5c. Critique to fig. 40 [= Fig. 413] in Wheeler (1930)

Wheeler (1930, p. 231 with fig. 40: *V. comstocki*) says the larva sometimes awaits its prey by merely leaving its head and thoracic segments with their ventral surface exposed, against the wall, but that usually it lies horizontally on its back (his fig. 40c) with its posterior half buried in the sand and its thoracic and first abdominal segments crossing the floor of the pit and covered with a very thin layer of sand. His figs. 40 c and d do not agree: in c the larva is entirely in the sand; in d it forms a bridge over the bottom of the pit, but the head end is apparently in the sand. The description of Buchner (1940: *V. vermileo*) corresponds with much of what I have seen in *L. [fortunata]*: anterior third of larva visible and lies slightly covered with sand like a small bridge across the ground of the funnel (loc. cit., fig. 2a). [1963: 258]

5d. Prey capture

There is universal agreement for all the species that the larva throws itself violently at the prey and coils its anterior segments rounded it, and there is also agreement regarding its behaviour in general. But little attention is paid to the fact that (at least in *L. [fortunata]*) after the victim has been immobilized, the larva may be seen for long periods to be uncovered, exploring or sucking the victim. I did not in *L. [fortunata]* see the larva move the head exploring the surface of the prey until after it had been coiled over or hooked on to the prey for a long time, and not until the prey had become motionless. But Wheeler (1930, p. 232) writes of the larva of *V. comstocki* just having captured the

Fig. 413. Activities of the larva in *Vermileo comstocki*. A, creeping on its back on the surface of the sand and just entering it with the anterior end; B, making the pit by tossing out the sand in jets; C, lying in wait at the bottom of the pit, lateral view; D, pit as seen from the surface with the sand-powdered anterior end of the larva across its floor; E, method of seizing the prey; F, absorbing the juices of the prey after it has been paralyzed and dragged under the sand. From Wheeler (1930). For critique to these figures, see text.

prey: "While the thoracic segments remain fixed the tip of the head moves about, carefully exploring the hard integument of the victim till it finds a suture or an intersegmental membrane," [1963: 260]

When the prey has been immobilized the larva inspects it at length all over, bending over it or just moving the mouth along the skin of the prey (fig. 96). Eventually it sucks from one or perhaps more selected parts, preferably between segments if an insect is concerned, and this may take hours depending on the size of the prey and of the larva. A soft dipterous larva given as prey could be observed shrinking during sucking. One *Lampromyia [fortunata]* larva studied in the evening, was still sucking the prey, an ant, the next morning. [1963: 251]Three ants of the genus *Formica* were taken from *Lampromyia [fortunata]* larvae, 3, 4 and 4 1/2 hours respectively, after the larva had captured them. After 3 hours intestines and Malpighian vessels were still seen. After 4 hours there were still some soft parts left, including intestines. After 4 1/2 hours all soft parts had disappeared. [1963: 252]

5e. Tossing sand in the direction of escaping prey

Usually the walls of the pit are rather steep, so that the prey insect is unable to climb it. But when for some reason the pit has flattened and the prey, say an ant, attempts to climb the slope out of reach of the larva, the latter may be seen to throw up sand by the same method as it uses in constructing the pit. Only, the sand is then hurled up preferably, or exclusively, to the side where the ant is, that is, not alternately to either side in regular fashion as in pit construction. And the throwings of sand to the side are now and then interrupted by blank thrusts in sagittal direction as if the larva tests whether the prey

has fallen down, so as to eventually hit it. Thus, these blank thrusts are not homologous with the blank thrusts to either side seen during pit construction, preceding digging to the respective side; but, with the thrusts with which the larva, when lying in wait, throws itself over the prey which falls down into the pit. [1963: 248]

5f. Ejection of abandoned prey or other foreign bodies

The tip of the anterior portion of the body is bent forwards, and placed so that the cadaver or foreign body touches the back of the anterior end of the larval body. Then it suddenly stretches its ring-like bent anterior segments and thus flips the insect carcass or foreign body away; about the same movement as when with the forefinger we wish to jerk away a bread ball from the thumb. [1968: 296]

[They are] not thrown out of the hole by the sideways jerks of the head employed to throw up sand, as one might perhaps expect. But the anterior part of the body + head is strongly curved forward (relative to its ordinary lying in wait position in the pit), the forward facing (anatomically upper) part of the downward curved tip of the body is placed against the carcass, and then by a snapping stretching of the tip it is jerked out.To throw up particularly heavy objects, e. g. pebbles, it may even bend the head so much that it passes to the side of the abdomen. [1963: 251]

5g. U or S shaped posture in a state of tension or feigned death, when the larva always lies on its side

When removed from its pit the slightly pink larva [= *Lampromyia fortunata*] may sometimes be outstretched, but generally assumes the lordotic S- or U-shape shown in figs. 2a-b. The back is at the concave side. The strongly curved posture in fig. 2a, with the two branches of the 'horseshore' sometimes tending to cross and the anterior segments tightly recurved ventrally, is not always seen, and when seen soon relaxes into the posture of fig. 2b, or an intermediate posture with more or less recurved anterior segments (fig. 3). This U- or S-shape is generally assumed as soon as the larva uncovered, for instance when the sand is removed from the side of the pit so deeply that it becomes suddenly exposed or falls down the resulting slope. Sometimes it does not react to mechanical stimuli; sometimes it may jump about when touched. It may remain motionless for some time, but eventually the head twists with its ventral side uppermost and starts boring into the sand in jerks (figs. 2c-d). The body gradually straightens out and follows, the segments being moved in sequence from behind forward; and by and by the body is turned so that it crawls on the back (figs. 2e-h). Soon the larva has disappeared in the sand. [1963: 240-242].

5h. Trail on sand

According to Wheeler (1930, p. 234 and fig. 41 on p. 233) the larva of the North American *Vermileo comstocki* Wheeler when placed in the dark sometimes emerges from the sand and wanders about on its surface leaving a characteristic trail consisting of a sinuous groove, made by the creeping body, and on each side a series of punctures, the impressions of the exploring head as it moves first to one side and then to the other. On exposure to the light the larva soon buries itself again and may then wander through instead of over the sand leaving no trails on the surface.

Trails of exactly the same appearance were seen in the Gran Canaria species [= *L. fortunata*] and *L. pallida* (fig. 4). As discovered by my coworker, Mr. John Nielsen, and confirmed by myself, they were produced both in daylight and in the dark and not by the larva wandering on the surface of the

sand, but by its creeping immediately below the surface. The groove results from the sand falling down behind the moving larva into the tunnel it has made, and the punctures on the side are made by the head of the larva emerging at the end of blind side tunnels. Neither the main tunnel nor the side alleys always collapse, depending on the nature of the sand. We have not been able to reproduce the behaviour which according to Wheeler produces the trails in *V. comstocki*, and cannot help suspecting that it has been surmised from the appearance of the trail rather than directly observed. [1968: 292-293]

5i. Leaping

This is described in detail by Wheeler (1930: 227-229) on *Vermileo comstocki*: If the immobile body of the larva, lying on its side, is now carefully explored with the point of a needle there may be no reaction at first, though pinching it with the tweezers, especially near the middle, will cause it to wriggle violently but at once resume its immobile S-shaped posture as soon as the pressure is removed. Certain regions of the body seem to be more sensitive than others to gentle contact, especially the basal abdominal segments, the obliquely truncated posterior end and the tip of the head. Touching these parts with the needle causes the larva to leap suddenly into the air and to land on its side 2 to 5 cm away, with precisely the same S-shaped flexure and continuation of its feint, but after it has been stimulated to make a few leaps in quick succession, touching the body or its sense-hairs at any point with the needle will induce the sudden reaction. The threshold of stimulation therefore falls rapidly, probably as a result of the larva's very gradual recovery from its feint. The movements of the leaping larva are quite simple and not too rapid to prevent analysis. They consist of two phases, a sudden, violent contraction of the antagonistic muscles, producing a complete reversal of the S-shaped flexure of the body, and as sudden a return to the original posture. Though the larva is lying on its side, the first contraction throws it off of the substratum, and the second occurs in the air or just as it is landing. This leaping is never spontaneous and cannot be induced by bright light, but only by mechanical contact or pressure, and certain gases, such as chloroform. It is therefore often resorted to when the larva happens to fall into the pit of another larva and is encircled or bitten.I infer that the young larva [of Mediterranean fruit-fly], like the young cheese-skipper, can only crawl. *Vermileo* larvae, however, will leap when they are only 3 to 4 mm long. The leaping of this insect is obviously a pure reflex and of little or no use, except perhaps in enabling it to escape capture by some other larva into whose pit it has chanced to fall, but the greater leaping powers of the mature, unprotected larvae of *Piophilila* [casei; Cheese-skipper] and *Ceratitis* [capitata; Mediterranean fruit-fly] may enable them to reach snug places for pupation either in the soil or under débris much more expeditiously than by crawling.

5j. Activity of the first instar larva

This is quoted from Wheeler (1930: 216-219) for *Vermileo comstocki*, Hafez & El-Moursy (1956a: 288-289) for *Vermileo* sp. from Egypt, Hemmingsen (1963: 239-240) for *Lampromyia fortunata*, and Hemmingsen (1968: 296) for *Lampromyia pallida*.

Vermileo comstocki: The eggs [which formed one cluster]hatched simultaneously April 28 as minute larvae of essentially the same structure and behaviour as the older larvae with which I was familiar. Though only 2 mm long and 0.2 mm broad the just-hatched differed from the older larvae only in being more transparent, with clearly visible alimentary tract and tracheal trunks, of more

uniform width, owing to the proportionally larger head, head-skeleton and thoracic segments, and in having more distinct eyespots and fewer and proportionally longer setae.The larvae had crawled only a short distance through the sand from the spot where the eggs were laid, and the first sign of their existence was a dense cluster, about the size of a silver fifty-cent piece, of minute pits, each only 2-3 mm in diameter and depth and occupied by a minute wormlion.The small cluster of about sixty individuals at first behaved like a colony. They showed no inclination to disperse and their pits were so close together as to be separated only by narrow partitions of sand. The sedentary propensity of the larva was shown also by the fact that within a few days the bottoms of the pits became rounded and bag-like, so that they were broader below than at the rim. From the first the larvae were very alert and active and showed no tendency to avoid the daylight. They often stood perpendicularly in their pits with the anterior half of the body exposed and the posterior half anchored in the sand. They were so sensitive that when an Aphid fell into one of the pits, the larvae in all the others at once squirmed and tossed out sand as if the prey were within their reach. The compact colonial arrangement may therefore be advantageous to the individual larvae since it must increase the chances of the prey's capture by some larva, even if it happens to escape from the one into whose pit it has fallen.Feeding usually occurred in the open, because the larvae in this stage neither coil around the prey nor drag it under the sand.....

When removed from their pits and placed in a watchglass the first stage were much more active than old larvae. They feigned death with the body in the usual S-shaped posture, but the feint was brief and the tiny creatures soon straightened out and began to crawl away. Then an interesting difference in locomotion between the very young and the old larva was disclosed. The latter invariably walks on its back, but the former, after proceeding some distance in this manner, sometimes turns over and walks on its belly. The first-stage larva is therefore ambigrade, or both dorsi- and ventrigrade, the older larva exclusively dorsigrade. Thus for a brief period after hatching the larva retains vestiges of the ancestral mode of locomotion so rigidly adhered to by nearly all other insect larvae.....

By May 4 the first-stage larvae, on a diet of Aphids, had visibly increased in size and were much more like the older larvae in behavior. They now encircled the prey when it was not too bulky, abandoned their upright position, became more sluggish and lay on their backs, with sandgrains sprinkled over their ventral surfaces, across the floors of their pits. They were still too small to drag the Aphids under the sand and therefore continued to imbibe their juices in the open. The pits had also increased in size, so that the whole colony occupied more space on the surface of the sand.

Vermileo sp. from Egypt: The newly hatched larva is about 2.2 mm in length and nearly three times as long as the egg in which it was confined. It is very active and makes several pores on the surface of the sand as if it were searching for a suitable position to construct its pits. One day or so after hatching, the conical pit has been formed by the small larva. The pit measures about 2 mm in diameter and about the same in length. Larvae hatching at higher relative humidity, i. e. 70 and 90 % could not penetrate through the sand, the latter being compact and wet. The larva normally crawls on its back when moving from one place to another.

It is noteworthy that the first stage larvae behave more or less in the same manner as the older ones. They assume the characteristic U-shape posture, feign death, penetrate in the sand, construct pits and make several leaps when mechanically irritated, exactly like the mature larva, except that the

former is relatively more active.

Lampromyia fortunata: I had no first stage larvae, the behaviour of which in *V. comstocki* Wheeler according to Wheeler (1930, p. 217, et seq.) is different from that of the older larvae. Wheeler's first stage larvae often stood perpendicularly in their pits with the anterior half of the body exposed and the posterior half anchored in the sand.

Lampromyia pallida: While older larvae of all the species are dorsigrade, Wheeler (1930, p. 219) in *V. comstocki* found that first stage larvae are both dorsi- and ventrigrade. I confirmed this for *L. pallida*, in which a larva recently hatched from the egg, occasionally walked on its belly when placed on paper.

5k. Renewals of the trap

It may be frequent as shown below.

Vermileo comstocki (from Wheeler, 1930): The wormlion is less sedentary than might be supposed from casual observation in field or laboratory. During the summer months, when food is most abundant, it not infrequently abandons its pit to make a new one some centimeters away. This may not be due to hunger, because it was also observed in the laboratory among larvae that were fed daily. [On p. 234]

The just-hatched larvae, as we have seen, may be regarded as gregarious, but as they grow larger and establish their pits farther from their birthplace and from one another they become solitary insects. [On p. 224]

Vermileo sp. from Egypt (from Hafez & El-Moursy, 1956a: 289): The activities of the larva amount to repairing and cleaning the old pit or digging another if the old one becomes unsuitable, and it is interesting to watch at night, by the help of an electric torch, the laborious larvae in a cluster, actively digging and repairing their pits.

6. Injection of venom and sucking of prey's content

Hemmingsen (1963: 264) wrote, "The antlion keeps its entire body hidden in the sand during preparation of the pit as well as during capture and consumption of the prey, but the anterior segments of the fly larva (at least *L. [fortunata]*) keeps naked out of the sand for long periods during preparation of the pit, capture, paralyzation, investigation, consumption and abandonment of the prey."

It is not necessarily certain to one of us (A. N.) from the literature which is the more preferred place to consume the prey, above or under the surface of the sand. The sucking site on prey's body is different from the venom injection site into prey's body. However they may *occasionally* be the same as shown below.

Wheeler (1930: 218) on *Vermileo comstocki*: "Sometimes it [= first instar larva] succeeded only in seizing one of the Aphid's legs. As soon as a hold had been secured it pumped venom into its victim and then commenced imbibing its juices, as was proved by the stream of green Aphis blood flowing into the larval stomach".

Some notes on injection and sucking are cited below.

Engel (1929: 153) on *Leptynoma sericea*: Generally only a leg of the prey is seized, and this is always treated in the same manner. The maxillae 'finger' the surface of the seized part till a place is found where the act of sucking can begin. When the prey is weakened by the loss of blood-liquor another place of attack is often sought — for instance, the intersegmental parts between the sternites

of the abdomen, or the occiput. After ten to twenty minutes' sucking the victim is generally pulled totally into the sand, and thus hidden from further observation.

Wheeler (1930: 232) on *Vermileo comstocki*: Since medium-sized ants and termites are paralyzed in three to five minutes, the venom can hardly be less powerful than that of the antlion.

Wheeler (1930: 253) on *Lampromyia canariensis*: On another occasion, when I dropped a codling moth caterpillar two centimeters long into one of the pits, there was an interesting demonstration of the gradual paralysis of the prey. The *canariensis* larva happened to embrace the last segment of the caterpillar. It writhed violently and emitted from its mouth great quantities of silk, but the larva held on and soon thrust its mandibles into one of its anal prolegs. The progressive paralysis induced by the injected poisonous saliva could be clearly followed in a posteroanterior direction, segment by segment, till the whole caterpillar was completely immobilized. *This required five minutes.* Observation was possible under the pocket lens because the long body of the caterpillar was stretched out in such a position that it could not be dragged under the sand with the sudden twist always adopted in the case of shorter and smaller prey. It seemed to me that the larvae of *Lampromyia*, perhaps on account of their greater size, were more alert, vigorous and rapacious than those of *Vermileo vermileo* and *V. comstocki*. I have been unable to detect any other differences in behavior between the species of the two genera.

Hemmingsen (1963: 250) on *Lampromyia fortunata*: Often the part of the prey held by the head of the larva is a leg round which the larva curves (as also observed by Engel, 1929, p. 153, for *L. sericea*). And the larva sticks to this part until the prey ceases moving. This may, e. g. with ants (*Formica*) not be until after 20-30 minutes. The larva is able to hold and move the prey with the mouth parts. These apparently also work as hooks, which at least in some cases appear to be able to hold and move the prey without any aid of the pseudopod.

7. Shape of the pit in relation to that of myrmeleontids

de Réaumur (1753) on *Vermileo vermileo*: The conical burrows of the wormlions are distinguished by having a greater depth than those of the antlion, with the same diameter. The apertures of the largest wormlion pits, however, scarcely attain the diameter of those made by half-grown antlions. (Translated by Wheeler, 1930: 319-320).

Wheeler (1930: 224) on *Vermileo comstocki*: The pits of old larvae vary from 2 to 3 cm in diameter and 1.5 to 2 cm in depth, and are smaller and have steeper walls than antlion pits.If it is excavated in fine dust or kaolin it is apt to become broader at the bottom, owing to the activities of the larva, and its cavity may assume the shape of a truncated instead of a pointed, inverted cone. A pit long inhabited may even become broader at the bottom than across its rim and therefore very unlike an antlion's pit.

Hafez & El-Moursy (1956a: 289) on *Vermileo* sp. from Egypt: The diameter of the larval pit varies from a few millimeters to about 2.5 cm, according to the age and size of the larva.

Hafez & El-Moursy (1963: 127) on *Vermileo* sp. from Egypt: "The pit of myrmeleontids (*Morter hyalinus* Oliver, *Cueta puella* Navas and *Cueta longula* Navas) has a steep wall with a diameter reaching up to 5 cm, while that of the wormlion is less steep and does not exceed 2.5 cm in diameter".

In the above statement words that may possibly be mistaken are in italics.

Hemmingsen (1963: 262-263) on *Lampromyia fortunata*: Even the shape of the pits is often

different. The pit of the ordinary European antlion (*Morter formicarius*) is funnel-shaped with a wide opening, whereas that of the fly larvae (Vermileonidae) is usually more cylindrical above, with steeper walls, and has a more or less circular bottom. Still, it may in some cases be more funnel-shaped especially, as pointed out for *V. vermileo* by Buchner (1940, p. 119), because the steepness of the walls decreases with rising sand grain size. In the free I was usually able to tell the species by the funnel shape, but not always. Some Myrmeleontidae construct accessories to their pits (Biró, 1898, p. 131; Wheeler, 1930, p. 84), e. g. a steep-sided pit with its center prolonged downward as a narrow tube serving as protection against the hot surface of the desert sand.

8. Role of pseudopod on first abdominal segment and a row of stout spines on posterior segments of abdomen

Hemmingsen (1963: 248) on *Lampromyia fortunata*: When a prey falls into the trap the larva throws itself over it like lightning and usually remains bent over it, the head jerking into some part on the other side of the prey (fig. 9a). Whether the pseudopod found on the ventral surface of the first abdominal segment (cf. fig. 1) plays any part in grasping or holding has been impossible for me to see.

Hemmingsen (1968: 295) on *Lampromyia pallida*: The larva throws its anterior segments over the prey. Though it is usually impossible to see the larva from the side during this, I succeeded in one case in good light to see in *L. pallida* how the bipartite pseudopod on its 5. (= 1. abdominal) segment rested against the body of the prey, in this case an ant.

Hemmingsen (1963: 250) on *Lampromyia fortunata*: The position of combs of spines on the abdomen (cf. fig. 1) may serve to stabilize the hind part of the larva, also when it throws the free part onto the prey. A tendency on the part of segments 8-11 to be pulled out will be counteracted by the large comb of forwardly recurved spines on the lower (true upper) part of segment 11 [= segment 7 of abdomen]; by some other forwardly directed minor spines especially on the sides of some segments; by the thickening of the abdomen towards the end; and possibly also by the four lobes with spines on segment 11, though these may appear to be of more importance in preventing movements of the fixed abdomen in the sagittal plane. A tendency to be moved in opposite direction to the segments 1-7, when they are thrown over the prey, about an axis in segments 7-8, is presumably prevented to some extent by the backward directed combs on segments 9-11.

The role of the pseudopod is discussed by Hemmingsen (1963: 255-256): Engel (1929, pp. 149-150, 153) reports for the South African *L. sericea* Westwood that the same method is used for throwing up sand and carcasses. But according to Engel in bending the anterior part the mouthparts seize a process on the pseudopod on the first abdominal segment. Then when the larva stretches the body a little and the mouthparts let loose their hold, the head acts as the strained end of an elastic spring which has been bowed together by its two ends, for the rest of the body remains firmly anchored in the sand. And he adds that as the anal end is fixed to one point, it is easy to understand how the conical form of the pitfall is constructed. Wheeler (1930, pp. 262-263) cannot accept Engel's interpretation of the function of the pseudopod, which Wheeler believes may act as a strigil for cleaning the mouthparts; though most observers have assumed that it acts in grasping the prey. Wheeler has been unable to witness the behaviour described by Engel, in any of the four species of [Vermileonidae], and he argues that "Engel himself admits that actual observation of the seizure of the papilla is impossible when he says that 'the movement is too quick for one to see more than the effect

of the mechanism.' "

Wheeler's view (1930: 263) on it is introduced below: "But, if we cannot accept Engel's interpretation of the pseudopod as a catch or ratchet for the spring-like movement of the neck, what is the function of the appendage? Most observers of *Vermileo* and *Lampromyia* larvae have assumed that it must act like the human thumb when opposed to the fore finger — represented by the anterior segments of the larva — in grasping and holding the prey. But the only species [= *Vermitigris fairchildi*] in which the organ would seem to be fairly well adapted to such a purpose is the one described in the next chapter (p. 272); in all the others it is too small and delicate to function very efficiently in this manner. It occurs to me, therefore, that it may act as a strigil or rather as a toothbrush, for cleaning the larval mandibles and maxillae. That these organs may occasionally become clogged with particles of sand or dust while the larva is making its funnel or feeding subterraneously on its prey, seems, indeed, highly probable. And certainly the tuft or series of backwardly directed spines, with which the pseudopoidal papilla is furnished would seem to be admirably suited for cleansing the mouthparts. For obvious reasons, however, it will not be easy to observe this act".

It seems that the primary role of the pseudopod is to seize the prey, contrary to the views by Engel (1929) and Wheeler (1930).

METAMORPHOSIS

1. Hatching

Hafez & El-Moursy (1956b: 340-341) on *Vermileo* sp. from Egypt: Under laboratory conditions, it was observed that all the eggs laid on the same day, either by one or many females, usually hatched at the same date.It may also be pointed out that no mortality was recorded among the eggs except in very rare cases, nearly all the eggs succeeded to hatch. Under room conditions, hatching took place at any time of the day. Before hatching, the larva can be seen inside the egg working its way through the chorion by means of its mouth parts and hard head skeleton. A small pore is then made forwards the narrow end of the egg and the dark brown head of the larva appears followed slowly by the rest of the body. When the greatest part of the body becomes free, the larva starts immediately to penetrate the sand while trying gradually to get rid of the egg-shell. Larvae encountering solid surfaces on hatching, always failed to free themselves from their egg-shells.

Le Faucheux (1961: 137-139) on *Vermileo vermileo*: See his fig. 5A-D (= Figs. 408-411).

2. Moulting

Wheeler (1930: 160) on *Vermileo vermileo*: Just before ecdysis, the larva stretches out and becomes quiescent beneath the bottom of its pit, two to three centimeters below the surface of the sand. Its thin but tough cuticle then splits the full length of the body in the mid-dorsal line and is cast off as a flat, beautifully transparent ribbon, to the anterior end of which the sloughed dark brown cephalopharyngeal skeleton is attached. During ecdysis the surface of the larva's body remains dry or at any rate if there is any exuvial liquid secreted between the old and new cuticles it is not sufficiently sticky to cause any adherence of the surrounding sandgrains. As soon as the new cuticle has acquired sufficient consistency, the larva proceeds to repair its pit and while thus occupied tosses the ribbon-

like moultskin out onto the surface of the sand, as if it were the exhausted carcass of a piece of prey.

Wheeler (1930: 259) on *Leptynoma sericea*: Failure to repair their funnels for some days is a sign that they are either digesting, moulting or about to pupate [after a letter from Hans Brauns to Wheeler].

Hafez & El-Moursy (1956a: 291) on *Vermileo* sp. from Egypt: When the larva neglects the pit, this indicates that it is searching for a more suitable place, about to moult or to pupate. When the larva is ready to moult, it neglects its pit and conceals itself in the sand. The old skin then comes out in the form of a white flat opaque structure having the shape of the larva but with a longitudinal slit extending from the head to the last abdominal segment. Attached to the exuvium are the sloughed dark brown cephalo-pharyngeal skeleton and also the tracheae. These moult skins are usually thrown by the larvae outside the pit.

3. Prepupation and prepupal period

Hafez & El-Moursy (1956b: 342) on *Vermileo* sp. from Egypt: Under room conditions, the larvae showed signs of pupation by the end of February. Prior to pupation, the larva generally stops feeding, leaves its pit and conceals itself in the sand. It then curves the anterior part of its body for several times so as to squeeze the contents of the alimentary canal in the hind part of the mid-gut. Then the junction between the mid-gut and hind-gut becomes opened and the contents of the gut are evacuated through the anal opening in the form of one or two hard spherical blackish pellets. After this process, the body-shape of the larva undergoes certain changes. It becomes somewhat reddish in colour, loses its transparency, and becomes opaque. But, the characteristic behaviour of the larva such as leaping in the air when mechanically irritated, feigning death, etc., are still retained. It can also penetrate the sand, but never constructs a pit. The larva changes into a pupa, generally *four to six days* after evacuating the contents of the alimentary canal. The larval skin always remains firmly attached to the last abdominal segment of the pupa in the form of a folded tube.

Hemmingsen (1963: 252) on *Lampromyia fortunata*: Before moulting or pupating the larvae ceased making pits, in some cases for *at least 10 days* prior to pupation.

4. Pupation

de Réaumur (1753; translated by Wheeler, 1930: 330) on *Vermileo vermileo*: Having reached the last stage of their growth, they therefore transform themselves into nymphs and that *often* without leaving their funnels.

Shelford (1901) on *Vermitigris* sp. from Mt. Penrissen, Sarawak: I brought down to Kuching alive several of these larvae, and one or two pupated; shortly before pupation, the larva leaves its pitfall and lies close to the surface of the sand, though completely covered; the anterior segments become much swollen and retracted, until the integument bursts, revealing beneath the brownish pupa; by some convulsive movements the whole pupa now appears at the surface, the larval skin being slowly shuffled off backwards, but never becoming entirely freed, so that the posterior end of the pupa always presents a somewhat ragged appearance. Unfortunately the heat of Kuching proved too much for these pupae, and none came to maturity, but shrivelled up;

Wheeler (1930: 160-162) on *Vermileo vermileo*: When the larva has reached maturity and begins to pupate, however, the last ecdysis is of a very different character. A layer of liquid accumulates

between the old larval cuticle and that of the pupa. The former splits only in the head region and is gradually pushed back to the posterior end of the abdomen, as a tube, thrown into compact, accordion-like folds. It is not thrown off but remains attached to the caudal segments of the pupa as a much wrinkled and contracted tail or appendage. The liquid above mentioned clearly serves two purposes: first, it lubricates the surface of the pupa and facilitates the pushing back of the larval skin, and second, owing to its stickiness it causes the surrounding sandgrains to adhere firmly to the surface of the pupa. I have not been able to ascertain the source of this ecdysial liquid.

At first the pupa lies horizontally in the sand one to three centimeters beneath its surface, but when the time for the emergence of the fly arrives, it wriggles up head foremost and thrusts its head and thorax above the surface of the sand.It is during the wriggling of the pupa to the surface that the sandgrains glued so firmly to its surface subserve, I believe, an important function.

Wheeler (1930: 239) on *Vermileo comstocki*: Hence the pupae of the great majority of insects may be said to have no 'behavior'. This is not altogether true of the wormlion pupa, because it does exhibit a very limited behavior just as it is about to become an imago. As we have seen, it then wriggles up to the surface of the sand and exposes its head and thorax and often more or less of its abdomen. The delicate imago, which could hardly make its way through even a few centimeters of sand, is thus enabled to issue into the air without difficulty. Sometimes the pupa does not cease all activity after it reaches the surface but continues to wriggle, with its hind end fixed like a pivot in the sand, till it produces a funnel-shaped pit precisely like that of the larva though smaller. According to Brauns (p. 260) the pupa of *Lampromyia sericea* may leap like the larva. Since I have never observed this behavior in the species of *Lampromyia* and *Vermileo* which I have reared, I believe that it must be exceptional and perhaps peculiar to very young pupae.

Hafez & El-Moursy (1956b) on *Vermileo* sp. from Egypt: After the larva is transformed into a pupa, the latter makes its way through the sand by a vertical movement with the head directed upwards. Then, it either remains in that vertical position with the head only visible from above, or it completely comes to lie over the surface of the sand (fig. 7). [On p. 343].When the adult is ready to emerge, the pupa exposes its head and thorax and often a part of the abdomen from the sand so as to facilitate the emergence of the delicate imago. The activity of the pupa is not limited to this process, but pupae were often observed standing vertically in the sand with their heads just exposed and making circles with the anterior parts of their bodies, with the result that small funnel-shaped pits similar to those of the larvae were produced. Moreover, some pupae were also able to leap when mechanically irritated. [On p. 345].

Hemmingsen (1963: 252) on *Lampromyia fortunata*: The pupae moved to the surface of the sand before emergence.

5. Emergence

Wheeler (1930: 161-162) on *Vermileo vermileo*: The dorsal thoracic cuticle then ruptures with the usual 'orthorrhaph', or T-shaped fissure and the imago slowly escapes.Unless there is some difficulty or abnormality that delays eclosion, the flies emerge only during the night and are found in the morning resting on the perpendicular sides of the dishes in the same position as other Rhagionids....

Wheeler (1930: 214) on *Vermileo comstocki*: The fly escapes from the pupa at night and is usually found in the early morning resting fully pigmented and mature on the side of the dish in which it lived

so many months as a larva.

Hafez & El-Moursy (1956b: 335-336) on *Vermileo* sp. from Egypt: Under laboratory conditions, the majority of adults emerged in the early morning between 9 and 11 a. m., while none emerged later in the day.The adult leaves the pupal skin through a T-shaped slit found on the head and thorax. When newly emerged, the adult is pale yellow and has a soft body which hardens and darkens in colour within about one hour.

Hemmingsen (1963: 252) on *Lampromyia fortunata*: As in [*Vermileo* sp. from Egypt] (Hafez & El-Moursy, 1956b) a T-shaped slit was left on the upper side of the head and thorax of the pupal skin after emergence.

LIFE CYCLE

1. Adult phenology in the field

First, three general accounts on it are cited.

Hafez & El-Moursy (1956b: 335) on *Vermileo* sp. from Egypt: Adults of *Vermileo* are hardly seen in the field. On several visits to Wadi Digla during the year 1952, no adults were met with. Fortunately, during 1953 on the 22nd of April, four living adults (3 ♀ ♀, 1 ♂), were caught. According to Wheeler (1930), no adult *Vermileo* of any species examined by previous workers was found in the field in its natural habitat with the exception of Rondani.

Hemmingsen & Nielsen (1971: 197) on two *Lampromyia* species from Canary Islands: Reports of adult flies are strikingly scarce compared with reports on the larvae. The senior author has seen flies of *Lampromyia fortunata* collected by Mr. J. M. Fernández in the open from April (El Rincón, Gran Canary) and of *L. canariensis* in May-June (Santa cruz de Tenerife and Puerto de la Cruz, Tenerife). Frey (1937, p. 43) reports 40 flies flying in July in front of the opening of a depression in a rock wall at Orotava, Tenerife.But there is nothing in these various facts to establish ecological differences as regards the temperature at which oviposition takes place in the open in the different species.

Stuckenberg (1960: 220) on some species from southern Africa: Nothing seems to be known of the biology of the adult flies in Southern Africa, which may be due to the fact that nearly all the species are rarely encountered in the field. A feature of the only species which I have observed alive (*Vermipardus intermedius*) has been the weak flight of the adult. There is some evidence that the adults may be found only during certain periods in the year. Dr. E. McC. Callan of Rhodes University, Grahamstown, has obtained a very large series of both sexes of *Vermipardus intermedius* from a verandah window in Grahamstown, and he observed that this species appeared only during the months of September to November (personal communication). Other species of which good series are available are *Leptynoma hessei* and *L. sericea*; every specimen I have seen of the former has been taken in October, while *sericea* has been collected from October to January.

Next, an important account is seen in Stuckenberg (1995b: 239) on *Vermipardus* from southern Africa fully cited here as well as his table 2.

In table 2 [= Table 1 copied here] the data on adult phenology are summarised by the month; any month for which there is only a record based on a reared specimen is indicated, as it is not yet known whether such records always correspond to times of adult occurrence in nature. If no distinction is made between rearing and field-based records, the following conclusions can be drawn: 1) There are

two peak periods in the year: (a) September-November (spring to early summer), with a peak in October (5 species); and (b) December-February (summer), with a peak in December (5 species). The two species of the *promontorii* group appear respectively in these two periods. 2) Both spring / early-summer and late-summer adult populations of *sathon* evidently occur. 3) The records for *univittatus* comprise the October-December period and a rearing record in May, so there may be two populations of adults per year. 4) If *sathon* is omitted, the members of the *brevirostris* group are mid-summer emergent, whereas the *intermedius*-group species are mainly spring to early-summer emergent.

Last, some of the field records are taken from the literature.

Lampromyia pallida: Wheeler (1930: 247-248) wrote that Bequaert records the following interesting data on *L. pallida* in Algeria, '.....At that season (June, 1910) adult flies were frequently seen resting on the rocky ceilings of the excavations.'

Vermileo ater: Stuckenberg (1965b): 1♂, end of May, 1904, Crete.

Vermileo comstocki: Wheeler (1930: 172): 19. vi. 1928, California (Yosemite); Nagatomi & Iwata (1976): 1♀, 4. vii. 1962, California (Nevada Co.: Donner Pass).

Vermileo opacus: Leonard (1930): 1♂, 6. vi. Nevada; 2♂♂, 2 & 5. v. 1902, New Mexico; Pechuman (1938): 6♂♂, 1♀, 19. vi. 1931, California (Arroyo Seco, Pasadena).

Vermileo vermileo: 'In southern France it lasts from the end of May to mid June (Le Fauchaux, 1961)' (after Hemmingsen, 1968: 299).

Vermitigris infasciatus: Oldroyd (1947): 1♂, 18. viii. 1927, N. Borneo (Bettotan, near Sandakan).

Six Palaearctic species of *Lampromyia* from Stuckenberg (1998): *canariensis*, 1♂, 20. vii. 1985, 1♀, 2. iv. 1968, Canary Islands. *cylindrica*, 2♀♀, 26. vi. 1910, Algeria. *hemmingseni*, 1♂, 1♀, 21. ii. 1992, 1♀, 3-20. iii. 1990, Canary Islands. *iberica*, 1♂, 21. vi. 1903, 1♀, 27. iv. 1927, Spain. *lecerfi*, ♀♀ during May to July, Morocco. *nigripennis*, 1♀, 12-13. vi. 1929, Morocco.

Two Chinese species of *Vermiophis*: *tibetensis*, 4♂♂, 17 & 21. vii. 1978, China (Tibet) (from C. Yang & Chen, 1987). *wudangensis*, 3♂♂, 2♀♀, 29. v. - 3. vi. 1984, China (from C. Yang & Chen, 1986):

2. Adult longevity under room conditions

Vermileo vermileo from France (Rennes) lived for 4 or 5 days (after Le Fauchaux, 1961).

Both sexes of 6 species in *Lampromyia* studied by Hemmingsen & Nielsen (1971) visited flowers in the container and lived for one or sometimes two weeks.

On the other hand, *Vermileo* sp. from Egypt studied by Hafez & El-Moursy (1956b: 347) refused food and had short longevity cited below.

Adults emerged under laboratory conditions (17-22°C and 34-56 % R. H.) between 5 and 8 a.m. Adult emergence continued in the laboratory for about one month from the middle of March to the middle of April.Mating took place 2 to 6 hours after emergence and lasted from 30 to 105 minutes. The male, unlike the female, showed certain tendency for repeated copulation. Rise in temperature seems to induce mating. The pre-oviposition period lasts from about 4.30 to about 7 hours. Oviposition begins two hours after mating. The eggs are laid singly in pits made by the female in the sand, then covered by a thin layer of sand. The time elapsing between depositing two eggs successively varies from 5 to 20 minutes. The oviposition period lasts from about thirty minutes to about 4 hours. The post-oviposition period is about 4 to 19 hours. The number of eggs laid by a single

Table 1. Phenology of *Vermipardus* flies, derived from label data and rearing records; any month for which there is only a record based on a reared specimen, is marked with X. From Stuckenberg (1995b).

	JUN	JUL	AUG	SEP	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY
<i>brincki</i>							*	*	*			
<i>promonrorii</i>				*			*					
<i>basuto</i>					*		*					
<i>intermedius</i>				*	*	*						
<i>sylphe</i>					X							
<i>univittatus</i>					X	*	*					X
<i>brevirostris</i>								X				
<i>munroi</i>								*	*			
<i>londti</i>							*	*				
<i>sathon</i>					*						*	

Table 2. Effect of copulation on the longevity of adults in *Vermileo* sp. from Egypt. (average of ten experiments in each case. From Hafez & El-Moursy (1956b).

Temperature in °C	Longevity in hours					
	Female			Male		
	Min.	Max.	Aver.	Min.	Max.	Aver.
Unmated	13	44	27.2	13	48	33.9
Mated	12	29	21.1	20	63	40.0

female varies from 8 to 30. The adult is very short-lived. The longevity of unmated females varies from 13 to 44 hours, while that of unmated males extends from 13 to 48 hours. Mated females lived shorter than unmated ones, while the mated males lived longer than unmated ones. The average longevity of the mated male was nearly twice that of the female [see Table 2; based on 10 instances in each case]. During the whole life of the adult, no food is taken.

3. Incubation period of egg

Wheeler (1930: 215-216) on *Vermileo comstocki*: The eggs, left undisturbed in the sand in which they had been deposited by the female [on April 20, 1917 in a hot-house], hatched simultaneously April 28

Hafez & El-Moursy (1956b: 340) on *Vermileo* sp. from Egypt: The possible effect of temperature and humidity on the incubation period was tested in a series of experiments, from which it was found that the most effective factor is temperature, while humidity has no apparent effect. Thus, keeping the temperature constant and varying the relative humidity, did not produce any appreciable effect on the incubation period. On the other hand, temperature variation showed a remarkable effect on this period. The length of the incubation period decreased with increased temperature. At low temperature, e. g. at 16°C, the incubation period was comparatively long and the eggs hatched after 28 days. At 22°C, they hatched after 11 days, at 27°C after 7 days, while at 30°C it lasted only 6 days. Above 30°C, e. g. at 33°C, the eggs shrank and did not hatch. It was also found that increasing temperature from 27 to 30°C seemed to have no appreciable effect on the incubation period as there was only a slight difference between that period at 27 and 30°C.

Hemmingsen (1968: 298) on *Lampromyia pallida*: Minute pits with recently hatched larvae were

seen at the end of July and beg. of August, 23, 26, 27 and 31 days after females of *L. pallida* had oviposited in the laboratory in the same substratum, so these are maximum figures for length of egg stage in this species at room temperatures in the Danish summer.

4. Duration of larval stage

It is long and the hibernation is made during this period. The account below seems to be applicable to almost every species of vermilionids.

de Réaumur (1753) on *Vermileo vermileo*: It was near the end of June when the first flies were born from the worms which I received at the end of August of the preceding year. At that time many of the larvae were very small and several were of more than medium size. It seems very probable therefore that there is each year only a single generation of these flies, and then only if the worms are well nourished. But if they be subjected to too severe and too protracted starvation during the months when the temperature of the air is at least mild, and if their need of food be not so great as to prove fatal, their transformation into nymphs is probably delayed till the following year, and if they be treated as badly during this second year, their transformation may be postponed till the third year. [Translated by Wheeler, 1930: 332-333].

Several examples are taken below.

Wheeler (1930: 150) on *Vermileo vermileo*: de Romand (1833), in a paper which is largely a confirmation of the earlier observations, was the first to notice the larva's ability to fast for long periods. He says: 'This larva may live a long time without feeding. Year after year I have left a great number of them in a box for six whole months without giving them any food and without their being able to obtain any. Yet when I again began to feed them, they were ready to eat and, after repeatedly changing their skin, attained their first transformation and eventually the stage of the perfect insect.' He fails to mention the time of year of this protracted fast, but it was probably the six months of normal hibernation, and his larvae seem to have pupated in the spring after feeding had been for some time resumed. This seems to follow from von Siebold's and Falcoz's observations mentioned below.

Hafez & El-Moursy (1956a) on *Vermileo* sp. from Egypt: The larval resistance to lack of food and water is very remarkable, and differs mainly according to age. Thus, the first stage larva can resist lack of food for about three months, while older ones for about twelve months. During this period of starvation, the larvae assume their normal activity and are often seen constructing and repairing their pits. [On p. 290]

As to the duration of the larval stage, it seems that the larva requires quite a long period to transform into adult. This is supported by the fact that the first stage larvae kept under room conditions did not exceed 5 mm after 18 months. For this reason it was not possible to include in this work the experiments conducted to compare between the duration of the different larval stage. [On p. 291]

The larva can withstand thirst and starvation for long periods varying from 3 to 12 months according to age and size. [On p. 297]

Hemmingsen (1968: 298): The length of the larval stage seemed to depend much on the amount of food prey offered.

38 larvae of *V. vermileo* collected in Italy (Ischia and Pompeji) 23-26. iv. 1961 had pupated in June and July, and imagines emerged 17. vi. - 15. viii. 1961.A few did not pupate until 1962.

78 larvae of *L. pallida* collected in Tunisia 23. iii. - 18. iv. 1961 had pupated in June-September,

and imagines emerged 1) during the last 10 days of June — and 2) (three times as many) from 10. September to end of October, the first 7 in September being males. Only a few single ones emerged at other times including July and August. Of 11 emerged in the next year in October-November and beg. December 9 were females; 2, males.

Some larvae of *L. pallida* collected in Tunisia in the spring 1961 did not pupate and emerge until 1963 or 1964 (the last ones in Sept.-Oct. 1964; none, in the winter months). Others, that hatched later in 1961 from eggs laid in captivity and not fed frequently, pupated and emerged from Sept. 1964 to Sept. 1966 (none, in the winter months). The larvae were not fed at all in the winter months of some years (at temperatures not below 9°C).

While these various facts may perhaps serve as indications, it is not known whether they reflect similar facts in the natural habitats, for laboratory conditions were neither constant nor was it attempted to vary them in imitation of natural conditions. Temperatures had varied between about 9°C to about 20°C according to season or heating.

The summer room temperatures at which these observations were made were doubtless lower than those obtaining in the natural habitats and more or so for *L. pallida* (Tunisia) than for *V. vermileo* (Italy).

5. Diapause

It is suggested by the accounts below that there is winter-diapause in *Vermileo comstocki*, summer-diapause in *Vermileo* sp. from Egypt, and winter-diapause in *Leptynomia sericea*. However, no experiment has been accomplished concerning the effect of daylength and temperature upon the start and termination of diapause in any vermilionid species.

Wheeler (1930: 234-235) on *Vermileo comstocki*: Towards the end of the summer the older wormlions descended more deeply into the sand and reacted feebly or not at all to prey thrown into their pits. They made no attempt to repair the old or to excavate new pits after the sand in the dishes had been shaken down till its surface was perfectly level. By the first week in October all the wormlions seemed to be hibernating, though summer temperatures still prevailed in the laboratory. If removed from the sand they feigned death and would leap when stimulated. When left undisturbed on the surface of the sand they slowly revived and at once buried themselves. I unearthed them once a week or oftener but except these brief responses they could not be aroused from their lethargy.

Wheeler (1930: 259) on *Leptynomia sericea*: They can go without food for a long time and some of them readily hibernate here during the months from June to October. This is the usual rainy season in Southern Cape Colony. [A letter from Hans Brauns to Wheeler].

Hafez & El-Moursy (1956b: 343) on *Vermileo* sp. from Egypt: Most larvae normally passed the summer in feeding and repairing their pits. Some of them, however, neglected their pits during part of the summer, probably taking a period of rest during which no food was taken, but began their normal activity again by the advent of winter.

6. Number of moults

Data on this are from Wheeler (1930: 213-223) on *Vermileo comstocki* as shown below: The older larvae collected in Alta Meadow [California] during August, 1917, were kept in a cold room [in New England] from November 12 to December 31 and then transferred to a hot-house [in New England].

After making their pits and feeding for some weeks they pupated during March and the first imago, a female, emerged April 1. Thereafter till the end of the month a few flies of one or both sexes emerged each day. That their development had been accelerated by shortening the period of larval hibernation is proved by the fact that the overwintered larvae collected in the Yosemite during May, 1930, did not produce flies till June, which as Essig has shown (p. 172) is the time of emergence in their native environment.

Oviposition, 20. iv. 1918; hatching, 28. iv.; 1st moult, 23. v.; 2nd moult, 13. vi.; 3rd moult, ca. 1. vii.; 4th moult, ca. 25. vii.; 5th moult, ca. 30. viii.; prepupation, 1. x.

Those reared in 1918 lived till late February in a hot-house and were then transferred to an unheated cold frame in the garden. When they were brought into the laboratory, April 8, 1919, it was found that they had not survived the damp and cold of the New England March.

In this connection, Wheeler (1930: 160) on *Vermileo vermileo* is introduced below: This [ecdysis or moulting of the larva] occurs repeatedly during larval growth but apparently at rather irregular intervals, depending on the rate of growth, which in turn depends on the amount of food the larva has been able to capture and assimilate. To determine the precise number of moults and the time of their occurrence would require very careful rearing of a number of isolated larvae from hatching to pupation, and this I have not been able to accomplish in the case of *V. vermileo*.

7. Duration of pupal stage

De Geer (1752) on *Vermileo vermileo*:for on the fifteenth of June it assumed the pupal form,Fourteen days later, or on the twenty-ninth of the same month, the insect abandoned its pupal envelope and revealed itself as a fly.....[Translated by Wheeler (1930: 310)].

Wheeler (1930: 161) on *Vermileo vermileo*: The pupal period varies with the temperature and extends according to my estimates, over two to three weeks.

Hafez & El-Moursy (1956b: 344-345) on *Vermileo* sp. from Egypt: See Table 3 (= table 2 in Hafez & El-Moursy, 1956b)..... It may be pointed out at 27°C, a high mortality of the pupae kept under 70 and 90 % R. H., was recorded. This may be explained by the fact that under these conditions the sand retains larger quantities of water and becomes compact, thus making it difficult for the pupa to make its way through it when the adult is ready to emerge. It may also be due to the fact that the very high moisture content of the sand, which sometimes amounts to saturation, has a direct fatal effect on the pupa.

Hemmingsen (1963: 252) on *Lampromyia fortunata*: Pupation took place indoors, largely from the end of April through May and June. The pupal stage usually lasted from 2 to about 3 1/2 - 4 weeks; in one case, during a warmer period, only 8 days. The variation was presumably correlated with the temperature, which was not regulated, but varied some degrees about 20°C.

Hemmingsen (1968: 298) on *Lampromyia pallida*: Observations not always made daily indicate a duration of the pupal stage of 27-34 days at unregulated room temperatures; 8-15 days at 25-26°C.

Hemmingsen (1968: 298) on *Vermileo vermileo*: They were not always observed daily. But the data indicate a duration of the pupal stage of ca. 18-20 days.

Hemmingsen (1968: 298) wrote: The longer duration of the pupal stage in *L. pallida* than in *V. vermileo* may thus possibly be ascribed to a greater lowering of temperature from the natural habitat temperatures of *L. pallida*.

Table 3. Effect of temperature and humidity on the pupal stage in *Vermileo* sp. from Egypt. From Hafez & El-Moursy (1956b).

Temperature in °C	Mean pupal duration in days under different R.H.			
	30%	50%	70%	90%
16	33.2	32.7	34.3	31.0
22	14.6	14.8	14.4	14.2
27	9.2	9.6	9.4	9.2
33	died	died	died	died

Figs. 414-415. *Hagenomyia micans* (Neuroptera: Myrmeleontidae). 414, distribution of larval pits in the box (35 × 46 cm); 415, percentage of mean numbers of the larvae in fine sand (horizontal axis: number of survivors). From Morisita (1952).

DIFFERENCE IN BEHAVIOR BETWEEN VERMILEONIDS AND PIT-MAKING MYRMELEONTIDS

This topic is discussed by Wheeler (1930), Hemmingsen (1963) and Hafez & El-Moursy (1963). Regrettably no personal study and review of the literature have been accomplished and no further mention is possible by us on the ethology and ecology of myrmeleontids.

Wheeler (1930: 279-280) wrote: It would be easy to make a long list of the detailed resemblances between the antlion and wormlion, including the various taxes and sensory reactions of the larva, its normally biennial life-span, the excavation of the pitfall, lying in wait, the poisoning, burial and extra-intestinal digestion of the prey, the occlusion of the posterior end of the stomach, death-feigning, the ability to remain for long months in [physiological] stupor, pupation in the sand, the wriggling of the pupa up to the surface to permit eclosion of the imago, etc. On the other hand, certain striking differences are to be noticed between the two insects, especially those in locomotion (retrograde in the antlion, dorsograde in the wormlion), in pit-excavation (circuitous method absent in wormlion), in the

lying in wait posture and in the method of capturing prey, in ability to leap (absent in antlion), in the making of a cocoon (absent in wormlion), etc.

Only two articles on myrmeleontids, which we happened to see, are introduced below.

Matsura (1997) on *Myrmeleon bore* and *Myrmeleon formicarius*: The renewals of pit took place due to either collapse of ground or abundant rain but almost none of them did due to the absence of prey. The life span without food was 84 days in *M. bore* and 147 days in *M. formicarius* as the average in 3rd instar larva, but no pit-reconstruction occurred during these periods in almost all of the individuals examined, apparently due to a marked decline of the respiration capacity.

In this connection, Wheeler (1930: 224) wrote, "A wormlion [= *Vermileo comstocki*] that is frequently fed may occupy the same pit for several days."

Morisita (1952) on *Hagenomyia micans* MacLachlan whose larvae live in the shaded place as in vermilionids: The larvae used in experiments were chiefly 8-12 mm in length. When a population of the antlion was kept in a sand box, they made the pits which were distant from one another. Thus, there was an interference among them. See Fig. 414.

One, two or more larvae were placed in a box (10.6×8.3×3.4 cm) which contained fine sand in one half side and coarse sand in the other for pit-making. Then, the numbers of larvae settled in each part were counted.

The antlion certainly prefers the fine sand for pit-making, when the population density is low. It is obliged to settle in coarse sand because of the increase of density. See Fig. 415.

There is no such experiment on any vermilionids.

BIOLOGICAL OBSERVATIONS ON *VERMIOPHIS GANQUANENSIS*

(from C. Yang, 1979; newly translated by D. Yang)

1. Habitat

The larval stage of vermilionids is long. The larvae make pitfall trap in sand in order to catch prey, so they must select the living environment. Dry and loose thin sands or dusts are necessary for the construction of traps. They always live under a steep cliff or an old building of man which is free from the rain and wind and from the direct sunlight and where small insects easily drop into the traps as possible prey. The larvae of *V. ganquanensis* are found in such environments. Ganquan is located near Yanan City, Shaanxi. There is a large cliff near the cave dwelling where I stayed for several days. Under the cliff, there were many small traps. I checked over 30 traps, and found only 1-2 traps of antlion at the outer margin. Later, I found similar traps at several other places. They are all located under the cliff. Later, I also found the traps in the discarded old cave-dwelling of Donggaotai, Qingquan Town.

2. Prey habit

The larvae prey on small living arthropods of various kind. The dry dead bodies of prey can be found in or near the traps and contain insects, spiders and crustaceans. The insects include 5 orders, 7 families and over 10 species such as ants and adults of the following: asilids and other flies, crickets, tenebrionids, noctuids and pyralids. We reared them successfully with insects of 7 orders and 20 families in the laboratory. So the larvae show no special selection of prey. But they cannot prey on

solid hard beetles. The phenomenon of cannibalism is rare among the individuals of the same species. It does not matter even if over 10 larvae are placed in the same bottle under the rearing conditions. This is different from the antlions. Two or more larvae of antlions cannot be placed in the same bottle.

The predaceous habit is highly successful. If the prey falls into the trap, it is difficult to escape. The larva lies at the bottom of the trap with the anterior half of the body exposed to the air, which may act as a bait for attracting the predaceous species which come to attack the wormlion (and then become prey). It is therefore not unusual that adult asilids are found in the traps. From this viewpoint, the vermilionids are more advanced than antlions, which only wait to prey on insects falling into the traps. So the traps of vermilionids are more numerous than those of antlions in a restricted area. But from the viewpoint of total long-term development, antlions are dominant in the number of species and in their geographical distribution. This is because the preying method of using itself as a bait is not safe, and primitive. Therefore, extant species of vermilionids are rare with a limited distribution.

As soon as the prey falls into the trap, the larva rolls up it with considerable speed. When it fails to catch the prey, it quickly whips around to make the sand fall so that the prey cannot climb up and is finally caught. Larvae pull the small prey caught to the bottom of the trap into the soil to suck. When it rolls up the larger or struggling prey, it raises the prey in the air to prevent it from escaping. After some time the prey is anaesthetized, then its body liquids are slowly sucked.

It takes a longer time for the larva to prey and eat. Sometimes after 6 hours, it still continues to suck. Based on the complete observation I made once, it takes 13.5 hours from capture to discarding the dead body of prey.

3. Larval activity and trap construction

The larva in the pit is very fierce predator. When moved off the pitfall, it becomes mild. The slender body often curves in a U-form. It can move forward on the ground surface through wriggling and soon dives into the soil with its head leaving a small hole in the soil surface. It differs from the antlions in constructing a pit. It dives obliquely into the soil, then pushes the soil to form the soil heap before the hole, and stretches the anterior part of its body out, throwing the soil regularly from left to right to form the pit. It bends its head back near the body, then strongly stretches to throw soil off the pit. If the soil particles are large, it throws them many times until eventually thrown out. When the old pit is repaired after eating prey, there is an action for clearing accumulated soil. Each time the wormlion always curves its head toward the dorsal surface of the body, touches the soil gently and then throw the soil out. It seems that this action is to make the soil of the pit wall harder because it becomes loose during the catching of prey.

The pits made by larger larvae are 20-30 mm in diameter, up to 40 mm. The larva keeps the ventral surface of its body upward at the bottom of the pit with 6-7 anterior segments exposed in the air and the rest held in the soil. In winter, they go down deeper into the soil for hibernation, then begin to make the pit again when the weather gets warmer. But at about 10°C in the laboratory, there is no diapause. When the prey falls in, the larva instinctively rolls it up. But if large animals invade, the larva quickly swings around and escapes into the soil.

4. Moulting and pupation

It is not easy to observe the process of moulting because the larvae live in the soil. The larvae dive into the soil from the bottom of the trap before moulting. They moult in the soil. Exuviae are milky

white, flat and complete. It is obvious that the skin divides along the mid-dorsal line from the head to just before the row of spines on the 7th abdominal segment. The larvae always keep the ventral surface of body upward, so that the exuvia, which is separated from dorsal surface, covers the larva dorsally and then the larvae escape from the exuvia. The exuviae may be used to study the characters of the larvae.

Pupation occurs in the soil. The pupae often climb to the soil surface where they leave the curved line due to crawl. Only the anterior part of the skin is divided along the mid-dorsal line (usually reaching to before the first abdominal segment). The skin moves backward, then contracts with wrinkling and finally slips over the abdominal apex. The pupa is covered with a layer of thin sand as the body has sticky liquid on the surface when it pupates. The pupa can move its abdomen freely. This enables it to move in the soil and, before emergence, to move to the surface to expose the head apex (sometimes half of head) in the air. When the pupa is touched, to be taken off the soil, it can jump by moving its abdomen, like a larva.

5. Adult emergence

The first adult appeared on April 19. It was placed in a small glass tube *without any food* and kept alive for 8 days. This is a record of longest life-span. It was known in the past that the adult usually died on the day of emergence. The life-span of adults in several species reared never exceeded 36-48 hours or 2-3 days (Wheeler, 1930). The adults emerged in late April and early May, and were kept alive for 1-8 days in our laboratory. Based on observation recorded for 24 adults, most of them can live for 3-4 days (62.5 %), some for one day (12.5 %), some for 5-8 days (12.5 %).

6. Do the adults feed?

Wheeler made several experiments and found that the adults of *Vermileo* and *Lampromyia* do not eat any liquid foods offered them such as honey and syrup. So he inferred that adults do not feed. The mouthparts of vermilionids is of the lapping-sucking type, so we can infer that they cannot prey on other animals but are therefor restricted to liquid food. In our experiments, when a little molasses was placed on the inner side of the stopper, adults quickly came to suck. When a ball of cotton-wool mixed with syrup or water is placed into the bottle, they also come to suck. The action of feeding is similar to that of muscids. They suck for a while and then raise the proboscis, move forward a little and then suck again. [The effect of nourishment on adult longevity and on number of the eggs laid per female throughout her life remain unknown in *V. ganquanensis* or any other genera and species of vermilionids].

7. Copulation and oviposition

This is rarely recorded in the literature. The copulation for adults reared was observed several times. One couple of adults are placed in the culture dish (6 cm in diameter) for three days without any food. At 3.00 p. m. of May 6, they are copulating. But they separated when the culture was moved for observation. Before long they copulated again for 15 minutes. After 3 minutes they copulated again for 18 minutes. Totally they copulated three times with some intervals. During copulation, the female carries the male dorsally. The male bends the tip of the abdomen laterally downward and turns out to meet the female while the male expands abdominal apex dorsoventrally to grip the female 8th sternite

in order to copulate. During copulation, the female vibrates her wings and the male constantly stretches out and draws back his abdomen. After copulation, the female was moved to a box with the sand on one side. She quickly climbed over the sand and turns the abdomen and slightly vibrated her wings. When the honey water is given, she quickly came to eat. After 10 minutes she left off and flew low or had a rest. After about 1 hour, she stopped over the sand and put her abdomen in it to oviposit. Another female copulated again after oviposition, and then layed a total of 24 eggs. They lay either one egg or two eggs together per site.

8. Incubation

The eggs began to hatch 20 days after laying. On May 25, the first small pit, 2 mm in diameter, appeared. There is movement at the bottom of the pit. A very small larva with U-shaped body is recognized and is taken from the sand and is put on a white paper background. The body length is less than 2 mm, pale yellow with black mouthparts and flattened, broad apex having 4 finger-like projections. When it is put back in the sand, it at once tries to dive downward with its head rather laboriously. It takes about 20 minutes to create a new small pit. After one week, there are many pits.

9. Life cycle

This is outlined here, compared with some field observations. Since finding larvae on June 9, 1971, only 2 empty shells were discovered on July 2. Pupae or adults were never seen again. It means that the emergence period of adults was over and they had oviposited in the soil. On July 25, there were many small pits found here, 4-5 mm in diameter. Obviously, small larvae were already incubated. The pits were not found in the field after the winter began. In early March, the pits appeared again. Reared larvae pupated in late March or mid and late April, 1972 and emerged in mid April and early May. Pupal period was about 20 days (13-25 days), adult stage very short (1-8 days), egg stage about 20 days. On late May the larvae of next generation appeared. Thus, they had one generation a year. But some of the larvae did not pupate and continued to prey with over 2 years to complete their life cycle.

10. Function of proleg

Finally a discussion on the proleg is made. There is a single projection with spines on abdominal sternite 1. Its structure is similar to the proleg on the ventral surface of the prothorax in chironomids and simuliids. It varies in shape and the number and arrangement of spines. It can be utilized to identify genera and species. For its function, there are three different opinions as follows. 1) Stop wheel [= a catch or ratchet for the spring-like movement of the neck]. Engel (1929) observed the larva of *Leptynoma sericea* Westwood from Africa. He believed that its function is 'stop wheel' for movement of the cervical part. When the head bends, the mouthparts grasp the proleg and then loosen it so that the head jumps back. This opinion is not acceptable from the structures of mouthparts and prolegs and from practical action. Wheeler (1930) denied this opinion. 2) Teeth-brush. Wheeler considered the proleg as a teeth-brush. The mouthparts are filled with sand and soil when larvae construct pits and catch prey, and the proleg can be used to clear mandibles and maxillae. 'It is not easy to see this action' as he stated. Once I saw that a larva actually brushes mouthparts after sucking for a long time. On this occasion, the head was brushed not against the proleg but against the soil. 3) Thumb. This is a widely accepted opinion. When the larvae attack, they use the anterior apex of the

body to roll the prey as if, when we grasp an object, the proleg acts as a 'thumb'.

I observed the prey habit of *V. ganquanensis*. It may prey on various insects with different size and hardness. It is reasonable to consider the proleg as a 'thumb.' I may consider the spines on the proleg as the 'nail.' Besides, when larvae dive into soil and move in soil, they always keep the apex of the head forward. The anterior apex of the larva is thin but the posterior apex is large. The anterior apex can easily move forward. The posterior apex wriggles forward with difficulty and the posterior apex fixing in the soil facilitates the movement of anterior apex. When the head and thorax curve downward, the proleg naturally projects forward just to make the anterior apex of the body stabilize. The transverse spine row is directed backward to strengthen this action. It has the same action as the spines on the posterior apex of the body, particularly 1 long spine row on the dorsal surface of the 7th abdominal segment to fix the posterior half of the body in the soil. Thus, the main function of the proleg is that of a leg. It must coordinate with the head and thorax in order to secure its prey and to move in the soil.

BIOLOGY OF SOME *VERMIOPHIS* SPECIES

(from C. Yang & Chen, 1993; newly translated by D. Yang)

C. Yang (1979) reported the biology of *Vermiophis ganquanensis*. Some additions are given here to the biology of *Vermiophis* species.

1. Living environment

The larvae always prefer a shaded place in which to live. At the time of field collection, more attention was paid to the places under the steep cliff, concave hole or near a building. Several photographs in fig. 5 show the living environment of *V. taihangensis*. Fig. 5A (Shangfungshaw, 500 m, Beijing): At the rock hole near the way to the mountain, we found the larvae of this species. At the same time we collected adults on the rockwall near it. Fig. 5B (Wutaishan, 1720 m, Shanxi): At the outer side of the wall of a large temple there are some pitfalls along the wall base. Fig. 5C & D (Yungang, Shanxi): There are many pitfalls in the grotto. [The photographs are not reproduced here].

2. Life history

Adults usually appear in May or June. The larvae can be collected throughout the year. In northern China, the larvae dive into the soil and do not build the pitfall during the winter. In several species, we collected the larvae and raised them to adults in the laboratory. It is easy to feed them with living insects. Generally they pupate and sometimes emerge as adults in late April. If larvae are in an early stage or do not take enough food, they would take one or two more years to become adults.

3. Abdomen of the male adult rubs the legs

The male genitalia of vermilionids have important characters for distinguishing species. But they have common characters: the tergum and sternum of the last segment are incised apically. During rearing we observed the cleaning action of males in *V. minshanensis* several times. The pair of hind tibiae rub the wing and abdomen, then rub with each other and pull the dust to the joint between tibia and tarsus. Then the abdominal apex is curved and rubs off the dust with its incision. This behaviour

is also observed in *V. wudangensis*. It seems that to rub (i. e. clean) legs is the one of the necessary functions for the incision (which has hairs) of male abdominal apex.

4. Oviposition

The oviposition action in *V. yanshanensis* is very peculiar. Fore and hind legs are outstretched to support the body, while the mid legs quickly stir (push) the soil to form the site. Then the apex of abdomen is curved and pushed down into the concavity to lay one egg. Finally she leaves off the place. On April 29, 1974 one emerged female oviposited for several times with the same action as shown in Fig. 404. Sometimes oviposition occurred on the side of the box and the mid legs also stirred (pushed) in the same way but the egg laid on the angle of the box could not be buried.

DISCUSSION

1. Sexual dimorphism

Wheeler (1930: 163) wrote, "The sexes are easily distinguished by the shape of the abdomen, that of the male being much narrower and more cylindrical than that of the female." Apart from the above matter no large sexual dimorphism is seen, since the eyes are widely separated in both sexes. An exceptional case is seen in *Vermilynx jasoni* and *V. vansoni* where the antenna differs in shape with sex (see Figs. 115-116, 284-285). There are two undescribed South African species of this genus in which no antennal sexual dimorphism is found (after personal communication of Stuckenberg).

For the many slight sexual differences, see original descriptions of the respective species.

2. Face

Face is flat in *Vermileo*, *Vermiophis* and *Vermitigris* but is produced forwards in *Lampromyia*, *Leptynoma*, *Vermilynx* and *Vermipardus*. Parafacial is thin or hardly visible. Clypeus ends near antennae; uppermost margin of clypeus is not well demarcated, and the boundary between clypeus and upper face is often obscure. Thus clypeus occupies almost whole surface of face. It is noted that there is a transverse furrow far below middle of clypeus in *Vermitigris sinensis* (♀) (this may be so in ♂ which is unknown). Face is narrower than one eye (in anterior view) and longer than distance from antenna to median ocellus. For the shape of face, see also diagnosis under each genus.

In the Tabanoidea (= Rhagionidae, Pelecorhynchidae, Athericidae and Tabanidae) and Xylophagoidea, the differentiation of clypeus, parafacials and upper face (= area between antennae and clypeus) (when present) is more distinct, and this is certainly plesiomorphic in relation to Vermilionidae.

It is noted that the protruded clypeus in Vermilionidae is seen only in *Lampromyia*, *Leptynoma*, *Vermilynx* and *Vermipardus* whose proboscis is sclerotized and conspicuously long, and is certainly apomorphic in relation to *Vermileo*, *Vermiophis* and *Vermitigris*.

3. Abnormal wing crossvein

It is often seen in various cells and we have found it in 7♂♂, 17♀♀ (of 2 genera and 7 species) from 71♂♂, 94♀♀ (of 5 genera and 11 species). The numbers of abnormal crossveins found are recorded below.

Vermileo comstocki (1 ♀ of 4 ♂♂, 3 ♀♀). 1 ♀: 4th posterior cell, 1 and 2 (in each wing).

Vermileo opacus (1 ♀ of 1 ♀). 1 ♀: 1st basal cell, 1 (on one wing).

Vermiophis ganquanensis (1 ♀ of 9 ♂♂, 19 ♀♀). 1 ♀: 1st posterior cell, 1 (on one wing).

Vermiophis minshanensis (4 ♀♀ of 8 ♂♂, 7 ♀♀). 1 ♀: 4th posterior cell, 1 (on one wing). 1 ♀: marginal cell, 1 (on one wing). 1 ♀: 2nd submarginal cell, 1 (on one wing); 1st posterior cell, 1 (on both wings); 4th posterior cell, 1 (on both wings). 1 ♀: 1st submarginal cell, 1 (on one wing); 1st posterior cell, 1 (on both wings); 4th posterior cell, 2 and 3 (in each wing).

Vermiophis taihangensis (4 ♂♂, 4 ♀♀ of 20 ♂♂, 32 ♀♀). 1 ♂: 1st posterior cell, 1 (on one wing). 2 ♂♂: 4th posterior cell, 1 (on one wing). 1 ♂: 4th posterior cell, 1 and 2 (in each wing). 1 ♀, 1st posterior cell, 1 (on one wing); 4th posterior cell, 1 and 2 (in each wing). 3 ♀♀: 4th posterior cell, 1 (on one wing).

Vermiophis taishanensis (1 ♂ of 1 ♂). 1 ♂: 4th posterior cell, 1 and 2 (in each wing).

Vermiophis yanshanensis (2 ♂♂, 6 ♀♀ of 15 ♂♂, 20 ♀♀). 2 ♂♂: 4th posterior cell, 1 (on one wing). 1 ♀: marginal cell, 1 (on one wing); 4th posterior cell, 1 (on both wings). 1 ♀: 1st basal cell, 1 (on one wing); 4 ♀♀: 4th posterior cell, 1 (on one wing).

4. Abnormally formed 1st posterior cell in *Vermiophis ganquanensis*

We have personally seen 5 genera (*Leptynoma*, *Vermileo*, *Vermipardus*, *Vermiophis* and *Vermitigris*) and 11 species, of which only *Vermiophis ganquanensis* often has an abnormal form of 1st posterior cell which is convergent apically, closed at wing margin or petiolate. Of 9 ♂♂, 19 ♀♀ examined, the following 1 ♂, 5 ♀♀ have such a cell: 1 ♂, convergent apically (on one wing); 1 ♀, narrowed somewhat apically (on both wings); 1 ♀, same as above (on one wing); 1 ♀, convergent apically (on one wing) and closed at wing margin (on another wing); 1 ♀, closed at wing margin (on both wings); 1 ♀, with long petiole (on one wing). See for example Fig. 127.

5. Wing venation in some *Lampromyia* and *Leptynoma*

Each of the veins R_{2+3} and R_4 is almost straight or gently curved and mouth of 2nd submarginal cell is not shorter than that of 1st submarginal cell in *Lampromyia* and *Leptynoma* (as in *Vermileo*, *Vermilynx*, *Vermiophis* and *Vermipardus*) but they are different in some species below.

Leptynoma (*Leptynoma*) *appendiculata* and *L. (L.) hessei* (see figs. 15-16 in Stuckenberg, 1996b): in vein R_{2+3} , apical portion bending upward and making almost a right angle with basal portion; apical half of vein R_4 bending upward.

Leptynoma (*Leptynoma*) *namaquaensis* (see fig. 16 in Stuckenberg, 1960) [= Fig. 57]: vein R_2 present independently; veins R_3 and R_4 each as in veins R_{2+3} and R_4 of *L. (L.) appendiculata* or *L. (L.) hessei*.

Leptynoma (*Perianthomyia*) *maculata* (see fig. 19 in Stuckenberg, 1996b): mouth of 2nd submarginal cell shorter than that of 1st submarginal cell.

Lampromyia pallida (see fig. 43 in Wheeler, 1930): vein R_{2+3} as in *L. (L.) appendiculata* and *L. (L.) hessei*; mouth of 2nd submarginal cell shorter than that of 1st submarginal cell, and vein R_4 not straight but bending upward to some degree.

Thus, veins R_{2+3} (or R_3) and R_4 differ markedly in venation in some species of *Lampromyia* and *Leptynoma*. I have seen directly 8 ♂♂, 2 ♀♀ of *L. (L.) namaquaensis* where the vein R_2 is always

present independently.

6. Status of *Leptynoma* (*Leptynoma*) *namaquaensis*

This species is peculiar by the presence of vein R_2 mentioned already (Fig. 57). Nevertheless, it belongs clearly to *Leptynoma* (*Leptynoma*). The presence of vein R_2 must be of no generic value, as Stuckenberg (1960: 228-230; 1996b: 252-253) discussed this character in detail.

7. Generic diagnosis in wing venation

Vein R_4 is almost vertical to vein R_5 in *Vermitigris* (Figs. 253, 257) but makes an acute angle with vein R_5 in other genera, i. e. *Lampromyia*, *Leptynoma*, *Vermileo*, *Vermilynx*, *Vermiophis* and *Vermipardus*, although it is nearly vertical in some species of *Leptynoma*.

8. Pulvilli and empodium

Stuckenberg (1960: 224) wrote, "The species of *Lampromyia* [s. lat.] are peculiar in having the pulvilli and empodium of the tarsi extremely reduced." He clearly shows three pad-like pulvilli and empodium (see his figs. 9-10) in *Vermipardus intermedius* and *Leptynoma hessei*.

In *Vermilynx vansoni*, "pulvilli minute, empodium represented only vestigially as the microscopic, setose apex of the ovate unguitractor plate (Stuckenberg, 1965a: fig. 2 [= Fig. 117])" (after Stuckenberg, 1995a: 31); "Character 5 [Pulvilli minute, empodium vestigial]: This is an autapomorphy for *Vermilynx*. The plesiomorphic condition, found in all other vermilionid genera, is for the pulvilli and empodium to be small, but clearly present and subequal" (after Stuckenberg, 1995b: 219).

On the other hand, Verrall (1909: 257) wrote, "Pulvilli in *Vermileo* and in [*Leptynoma*] *sericea* three, rather small but equal, and the claws outspread as usual; but in *Lampromyia pallida* there are only two minute pulvilli, which are under the long straight approximated claws (as in *Leptogaster*). The absence of empodium should be reconfirmed in *Lampromyia pallida*."

As shown in Fig. 117, the empodium is not bristle-like in *Vermilynx vansoni*. Yeates (1994: 83) wrote, "Woodley (1989) considered a bristleform empodium to be apomorphic for the Muscomorpha excluding the Nemestrinoidea. I have found the descriptive term 'bristleform' inadequate to describe many asiloid empodia. They are most often reduced in size and triangular in shape with an acute apex but bearing small setae (figs. 298-299). No asiloids were found with pulvilliform empodia in this study. Some asiloids have lost the empodium: Mydidae (Wilcox, 1981); some Apioceridae (Peterson, 1981); four genera of leptogastrine Asilidae (Martin, 1968), and *Hilarimorpha* (Webb, 1974)".

We have seen some specimens of *Systropus* spp. (Bombyliidae) which have a bristle-like empodium.

There is probably no fundamental difference between 'bristle-like' and 'triangular in shape with an acute apex.'

9. Male genitalia

We believe that most of the illustrations of male genitalia available have been reproduced in this paper.

The male genitalic characters of Vermileonidae are as follows: Aedeagus (= half-open umbrella

plate, trumpet tube, or sclerotic structure) bag- or tube-like. Aedeagus usually with lateral extension only in *Lampromyia*, *Leptynoma* and *Vermilynx*. Ejaculatory apodeme long and extruded anteriorly from aedeagus or reduced in size and situated inside aedeagus, and with or without lateral ejaculatory process. Posterior margin of fused gonocoxites is level with or higher than (= extending more posteriorly) base of gonostyli (it is not in Xylophagoidea [except Xylophagidae] and Tabanoidea). Sternum 9 (= hypandrium) absent. Gonostyle situated at dorsal posterolateral apex of gonocoxite. Dorsal bridge present and gonocoxal apodeme (if present) situated on dorsal bridge. Tergum 10 absent. Except for *cylindrica* group of *Lampromyia* and *Leptynoma* (*Perianthomyia*), tergum 9 (= epandrium) usually wider than long and with anterior and posterior margins concave. In *Vermileo ater* (from Crete) and *V. niloticus* (from the Sudan), tergum 9 has two projections on each posterolateral corner (Stuckenberg, 1965b: fig. 3 [= Fig. 99]).

Teskey (1981: 532) wrote, "The male genitalia of Vermileonidae most closely resemble the Therevidae." But this statement is incorrect and misleading.

The male genitalia of Vermileonidae are easily distinguished from those of other families whose characters are given below.

Xylophagoidea (except Xylophagidae) and Tabanoidea: See above.

Stratiomyidae: Tergum 10 well developed and produced posteriorly.

Pantophthalmidae: Dorsal bridge absent; gonocoxal apodeme conspicuously long.

Xylophagidae: Ejaculatory apodeme consisting of horizontal and vertical plates, of which the latter is flattened laterally.

Therevidae: Anterior part of aedeagus divided into dorsal and ventral apodeme. Base of gonostylus distant from dorsal posterolateral apex of gonocoxite. Each gonocoxite with three posterior appendages, that is, outer style (which is often not articulated with the gonocoxite), inner style (= gonostylus), and ventral lobe.

Rhagionempididae (= Apsilocephalidae): Ejaculatory apodeme mushroom-shaped, with a ring-like sclerite at base (= posterior end). Gonocoxite with a posterolateral crest. Large sternum 9 present. Tergum 9 with a large posterolateral process (= surstylus).

It is not necessarily certain whether aedeagus in Vermileonidae represents phallus (= aedeagal sheath) or endophallus (= aedeagus s. str.). It may possibly be homologous with half-opened umbrella plate inside phallus in Heterostomidae and some genera (*Atherimorpha*, *Neorhagio* and *Rhagina*) of Rhagionidae.

See Key 2 to genera based on male genitalia. Aedeagus, aedeagal lateral extension (seen in *Lampromyia*, *Leptynoma* and *Vermilynx*), fused gonocoxites, mid-posterior ventral concavity in fused gonocoxites, dorsal bridge, gonostylus, tergum 9, etc. vary in shape with species.

Lampromyia + *Leptynoma* + *Vermilynx* is put together in Key 2. The autapomorphic or synapomorphic characters of each genus or group are here cited according to Stuckenberg (1996a, b, 1998).

Lampromyia (*canariensis* group + *cylindrica* group; both from the Palaearctic region): T8 of ♂ deeply concave on posterior margin, much shortened over most of its width and withdrawn under T7, exposed and setose only laterally [Figs. 43-44, 48-50]. T8 is completely reflexed beneath T7 except for a small, triangular lateral section which is exposed on each side; the posterior edge of T7 is thus adjacent to the base border of the epandrium; T8 would be exposed only during mating, when the

infolded intersegmental membranes would be extended. Associated with this feature is a characteristic posture in which the hypopygium is flexed dorsad relative to the longitudinal axis of the abdomen, found in the palaearctic species except those in the Canary Islands. The very large size of the epandrium is made more conspicuous by this posture (figs. 24, 34). [This character state] occurs in the Palaearctic species of *Lampromyia* and is assumed for *nigripennis* and *lecerfi*; in the Southern African species, T8 is exposed and of normal, plesiomorphic form. The least modified form of T8 among the Palaearctic species occurs in *hemmingseni* [Fig. 33]. [1998: 73]

Lampromyia (*canariensis* group; 3 spp. from Canary Islands): *Mediodorsal projection present on upper margins of synsternite* [= fused gonocoxites]. This structural elaboration is apomorphically developed only in the species of the *canariensis* group [Figs. 18, 23, 28, cd]. It varies in form specifically as illustrated [as Figs. 18, 23, 25, 28-29, cd]. *Aedeagus almost straight* [Figs. 20, 26, 30]. A character of the Canary Islands species. In all other species of *Lampromyia*, and in other Afrotropical genera, the aedeagus is strongly curved upwards. [1998: 74-75]

Lampromyia (*cylindrica* group; 6 spp. from southern Europe and NW Africa): *Epandrium elongate, projecting well beyond apex of synsternite* [= fused gonocoxites] [Figs. 44, 48]. This condition contributes to the characteristic appearance of ♂♂ of the *cylindrica* group, which seemingly have a disproportionately large hypopygium. A relatively much shorter epandrium occurs in the *pilosula* and *canariensis* groups, and in other Afrotropical genera, except, but less markedly, in *Perianthomyia* [Figs. 70, 74]. *Aedeagus lacking lateral projections*. In *Vermipardus* the aedeagus is a curved tube, this evidently being the plesiomorphic form. In species of *Leptynoma*, *Vermilynx*, the *pilosula* group, and the *canariensis* group [Figs. 20, 26, 30, p], the aedeagus appears trifold because of two lateral extensions or arms; these do not include the lumen of the aedeagus. In species of the *cylindrica* group, only *funebis* and *pallida* have any structural development which may be comparable, in the form of dorsally directed flanges on each side of the aedeagus [Figs. 42, 54, p]. As the trifold condition is so widespread, the lack of these lateral arms must be a reversal. [1998: 74]

Lampromyia (*pilosula* group; 3 spp. from southern Africa): *Dorsal bridge extended posteriorly in a symmetrical, apically rounded lobe*. Found only in the Southern African species [Figs. 10, 15]. This is a more elaborate state; usually the dorsal bridge is transversely rectangular or arched. [1998: 73]

Leptynoma: Aedeagus trifold, strongly upcurved, lateral extensions dorsally directed, a basal keel present ventrally; gonostylus either bifid or with a ridge terminating in an angular point on dorsal margin. [1996b: 251]

Vermilynx: 1). *Epandrium*. In *V. vansoni* [Fig. 119] and in *V. jasoni* [Fig. 123], the epandrium is broader than long, deeply arched both transversely and longitudinally, and has a semicircular posterior emargination through which the proctiger (*pr*) protrudes; a unique feature is the extensive incurving of the rim (*vf*), especially posteriorly as a ventral secondary 'roof' around the emargination in which the proctiger lies. 2). *Proctiger*. In both species, the proctiger is elongate, tubular, and suspended at its base in the membrane enclosing the epandrium ventrally. 3). *Synsternite*. See *Sn* in [Figs. 118, 120-122]; the synsternite is short and deep, and its mediodorsal margins are produced into strongly projecting forms (*md*). 4). *Ventral aperture*. This is very small [Fig. 122, *va*], in contrast to the condition commonly seen in other genera in which it may occupy a much greater area in the ventral surface of the synsternite. 5). *Apical guides*. There are shaped as apically rounded, moderately elongate, posteriorly-directed flanges that incline ventrally, so that they approach one another closely

along their ventral margins and narrow the posterior opening to the ventral aperture [Figs. 122, 124, *ag*]. 6). *Gonostylus*. These are moderately elongate and incurved; they have a projection on the dorsal surface that appears in dorsal view as a flat, rounded lobe [Figs. 118, 120, 122, 124, *gs*]. [1996a: 198]

Further, the male genitalic characters are introduced according to Stuckenberg (1995b, 1996b) in the following taxa.

Vermipardus: Dorsobasal margins of fused gonocoxites joined by a sclerotised dorsal bridge on which aedeagus is attached by its posterodorsal edge; gonocoxites almost completely united ventrally, hypandrium absent or completely fused with them; mediolaterally the gonocoxites project posteriorly and partially enclose a ventral median aperture of specifically variable size; gonostylus inwardly articulating, moderately tapering, bifid in species of *brevirostris* group; epandrium broader than long, its basal margin concave; proctiger small, cerci scarcely protruding; aedeagus tubular, without lateral extensions, its base widened or flaring, its middle section directed posteriorly, apical section curved upwards, ventral keel rarely present on basal part, outer surface in most species with minute, flat, sharp, subtriangular, recumbent, sessile, distally-directed teeth (spiculae), sometimes these developed on margins of lateral flanges along aedeagal stem; ejaculatory apodeme a short, slightly flexed rod completely within basal part of aedeagus, its distal end attached to proximal end of membranous internal tube leading to aedeagal apex; [Figs. 195-249]. [1995b: 220-221]

Leptynoma (*Leptynoma*): Male genitalic features are as follows: hypopygium relatively inconspicuous, extending posteriorly in same axis as abdomen; epandrium short, about as long as broad, its margins progressively incurved posteriorly, not projecting beyond gonostylus; proctiger large and broad, occupying much of ventral surface of epandrium; apical guides in form of incurved flanges; gonostylus short, stout, incurved, asymmetrically bifid; lateral extensions of aedeagus arising almost on same plane as upcurved median intromittent tube, with minute denticles in most species; [Figs. 58-69]. [1996b: 252]

Leptynoma (*Perianthomyia*): Hypopygium relatively large, conspicuous, upwardly flexed relative to longitudinal axis of abdomen, this posture associated with a reduction of tergite 8 and increased intersegmental membrane between pregenital sternites; epandrium large, extending beyond gonostylus, with two dark, well-sclerotised, irregularly shaped bodies of unknown function on ventral surface, these visible externally as two dark spots; proctiger tubular, elongate, attached at base; gonostylus slender, almost straight, posteriorly directed, with conspicuous pointed angle on dorsal margin where width is abruptly reduced; apical guides slender, almost straight, converging, apically pointed; dorsal bridge strongly arcuate over posterior margin, produced on each side into a rounded flange which protrudes above dorsal edge of synsternite; aedeagus protruding dorsally above synsternite, lateral extensions situated at about midlength where deepest ventral part of curvature of aedeagal tube is attained; [Figs. 70-77] [1996b: 257].

10. Female genitalia

The diagnosis below is after Nagatomi & Iwata (1976: 39-40) on *Vermileo comstocki* and *Vermipardus intermedius*. See Figs. 294-295.

"Abdomen without telescopic ovipositor; intersegmental membrane between terga 7 and 8 short and hardly visible. Cercus is not elongate and *probably unsegmented but has two processes*. Sternum 10 undivided or *divided longitudinally according to genus or species*. Tergum 10 absent. Tergum 9

roughly rectangular or triangular according to genus or species. Sternum 8 elliptical in shape and wider than long. Tergum 8 wider than long and wider than sternum 8. [A reexamination is needed for the italic portion above]."

The cercus is two-segmented in *Vermiophis* spp. (see C. Yang, 1979: fig. 3; C. Yang & Chen, 1986: fig. 63; C. Yang & Chen 1993: figs. 1E, 2E, 4E) [Figs. 310-315], and *Lampromyia hemmingseni* and *L. lecerfi* (see Stuckenberg, 1998: figs. 14, 16) [Figs. 296, 298].

Stuckenberg (1995b: 221) on *Vermipardus*: "Female genitalia: ninth tergite deeply arched, posteriorly forming a cowl or hood over proctiger; proctiger not extensible; cerci two-segmented, basal segment deeper than long, its posterior corners rounded, a subsemicircular excision in hind margin in which the much smaller, short, squat second segment is placed; three subequal, spherical spermathecae; ninth sternite well sclerotised, transverse, posterior margin concave."

It should be noted that in *Vermiophis wudangensis* the sternum 9 is present as a pair of sclerites which are widely separated. The sternum 9 is not a genital fork but must be an external sclerite (if present).

Stuckenberg (1995b: 29) wrote, "my unpublished studies of *Lampromyia* Macquart show female cerci with a large first segment lacking a ventrolateral expansion."

Stuckenberg (1998: 73, 75) showed the four synapomorphic character states among the species groups of *Lampromyia*.

1). *T9 of ♀ extended posteriorly as a short hood with a broad, shallow excision in posterior margin* (fig. 15) [= Fig. 299] [apomorphic state]. The ♀ terminal tergite of all *cylindrica*-group species has this form.In the *canariensis* group T9 has other forms.

2). *Cerci of ♀ with apical segment short, largely hidden beneath basal segment* [apomorphic state]. In females of the *cylindrica* group, only the basal cercal segment is clearly exposed in dorsal view (fig. 16) [= Fig. 298]. In the *canariensis*-group species, the apical cercal segment is longer and clearly exposed (fig. 14) [= Fig. 296].

3). *S8 of ♀ with two closely adjacent, median, basal swellings* (figs. 14, 20) [= Figs. 296-297] [apomorphic state]. An exceptional apomorphy in *fortunata* and *hemmingseni* of the Canary Islands.

4). *Spermathecae small, gradually and irregularly swollen terminally, and darkly sclerotised* (fig. 10) [= Fig. 300] [apomorphic state]. This condition occurs in the Southern African species only; all Palaearctic species have subspherical, less thickly sclerotised, relatively larger spermathecae (figs. 11-13) [= Figs. 301-303], and the same general form occurs in the outgroups. Spermathecal form varies greatly in the Orthorrhaphous Brachycera, but the spherical condition occurs frequently and is probably plesiomorphic. The *canariensis* group and both subgroups of the *cylindrica* group may each be characterised by modifications of the spherical form: all species of the *canariensis* group have the same, least specialised spermatheca (fig. 11) [= Fig. 301]; in the *cylindrica* group, material was inadequate for all the species to be examined, but probably in the *cylindrica* subgroup the form in fig. 12 [= Fig. 302] is characteristic, in which a membranous sac arises from a relatively large aperture in the spermathecal wall and gives rise to the duct; in the *pallida* subgroup, the almost colorless spermatheca with long tapering extension and rugose inner surface as found in *pallida* (fig. 13) [= Fig. 303] probably occurs also in *iberica*.

For spermatheca in *Vermipardus*, see Stuckenberg (1997: 190-192) and Figs. 304-309.

It is necessary to study again more carefully the female genitalia of *Vermileo* and *Vermipardus*

based on more species. There is no illustration of female genitalia in *Leptynoma*, *Vermilynx* and *Vermitigris*. The female genitalia will be treated in several species by some of us in the future.

11. Some common characters among the genera

Lampromyia, *Leptynoma*, *Vermilynx*, *Vermiophis* and *Vermipardus*: (1) Antennal flagellomeres 2-5 gradually tapering apically and not arista-like in *Leptynoma*, *Vermilynx*, *Vermiophis* and *Vermipardus* (this is also so in some species of *Lampromyia*) and (2) last joint of flagellum much shorter than rest of flagellum in *Vermilynx*, *Vermiophis* and *Vermipardus* (it is always arista-like and roughly as long as or longer than rest of flagellum in *Lampromyia* and *Leptynoma*).

Vermileo, *Vermiophis* and *Vermitigris*: (1) Proboscis short and fleshy and much shorter than height of head; (2) ejaculatory apodeme large and extending far beyond anterior margin of aedeagus (= half-opened umbrella plate) in *Vermiophis* and *Vermileo* [but the male genitalia are not yet studied in *Vermitigris*].

12. Some distinguishing characters among the genera

Lampromyia, *Leptynoma*, *Vermilynx*, *Vermiophis* and *Vermipardus*: (1) Proboscis short and fleshy and much shorter than height of head in *Vermiophis* but long and sclerotized and much longer than height of head in *Lampromyia*, *Leptynoma*, *Vermilynx* and *Vermipardus*; (2) ejaculatory apodeme large and extending far beyond anterior margin of aedeagus (= half-opened umbrella plate) in *Vermiophis* but situated inside aedeagus (= trumpet tube) in *Lampromyia*, *Leptynoma*, *Vermilynx* and *Vermipardus*.

Vermileo, *Vermiophis* and *Vermitigris*: Antennal flagellomeres 2-5 gradually tapering apically and not arista-like and last joint of flagellum (= flagellomere 8 or sometimes 6) distinctly shorter than rest of flagellum in *Vermiophis* but flagellomeres 1+2 (which is as long to nearly twice as long as wide) with a long 3-segmented arista in *Vermileo* or flagellomere 1 (which is distinctly longer than wide) with a long 3-segmented arista in *Vermitigris*.

13. Generic status of 'Vermileo' spp. from North America and Central America

The male genitalia of *Vermileo comstocki* from North America are different markedly from those of *Vermileo* (4 spp.) from southern Europe, Crete and Sudan as shown in Key 2 and Figs. 87-91, 97-113. This suggests that *V. comstocki* may be separated generically from the Palearctic *Vermileo*. In the North American *V. comstocki* and *V. opacus*, frons is markedly wider towards vertex and occiput is swollen. In *V. vermileo* (from southern Europe), *V. dowi* (Cuba) and *V. willetti* (Mexico) the frons is parallel sided, and at least in the former two species the occiput is flat.

It is absolutely necessary to study all of the Central American species, especially *V. tibialis* (= the type species of *Pheneus* Walker, 1852) and *V. fascipennis* (= the type species of *Arthrostylum* Williston, 1895) for solution of their generic status.

14. Status of the recently erected genera of *Lampromyia* s. lat.

As the status of the four 'genera', various cases are conceivable as below.

(1). One genus: *Lampromyia* includes two subgenera, that is, *Lampromyia* s. str. (with *vansoni*, the type species of *Vermilynx* and *sericea*, the type species of *Leptynoma*) and *Vermipardus*, or includes four subgenera, that is, *Lampromyia*, *Leptynoma*, *Vermilynx* and *Vermipardus*.

(2). Two genera: *Lampromyia* and *Vermipardus*; *Lampromyia* consists of three subgenera, namely, *Lampromyia* s. str., *Leptynoma* and *Vermilynx*.

(3). Four genera: *Lampromyia*, *Leptynoma*, *Vermilynx* and *Vermipardus*.

At the present state of our knowledge, we follow Stuckenberg (1996b) and accept (3) above.

In *Lampromyia pallida*, the type species of *Lampromyia*, "the aedeagus is without accessory processes and resembles the type found in *Vermipardus*" (after Stuckenberg, 1960: 247) [Figs. 36-37]. However, this species differs markedly from the members of *Vermipardus* in other points including the male genitalia.

15. Phylogenetic relationships among the genera

In Fig. 416, the plesiomorphic (square mark), apomorphic (circle mark), and autapomorphic (in respective group) (triangular mark) character states are shown. The numbers 1-14 correspond to those in Fig. 416 and are explained below. The numbers 7-14 are entirely based on Stuckenberg (1960, 1995a,b, 1996b).

1. Proboscis short and fleshy and much shorter than the height of head (Figs. 78-79, 146, 174, 250, 258-259, 261, 268, 270) (square); proboscis long and sclerotized and much longer than the height of head (Figs. 1-3, 56, 114-116, 190-192, 263-264, 267) (circle).

2. Ejaculatory apodeme large and extending far beyond anterior margin of aedeagus (= half-opened umbrella plate or sclerotic structure in Stuckenberg, 1965b: figs. 1, 7-10) (square); ejaculatory apodeme situated inside aedeagus (= trumpet tube) (circle). [This character state is unknown in *Vermitigris*].

3. Antennal flagellum gradually tapering apically (square); abruptly widened base of flagellum (=

Fig. 416. A possible phylogenetic tree of the known genera of Vermileonidae. It is taken from Stuckenberg (1996b) concerning *Lampromyia* s. lat. Original.

Figs. 417-420. Possible phylogenetic trees of the known species of *Vermiphis*. Original.

apparently flagellomeres 1+2) roughly as long to nearly twice as long as wide, and with an arista (which is 3-segmented) (circle).

4. Palpal segment 2 widest around middle and then gradually tapering apically (square); palpal segment 2 swollen (in lateral view) and without an elongate portion gradually tapering apically (see fig. 405 in Nagatomi and Soroida, 1985) [= Fig. 260] (circle).

5. Vein R₄ making an acute angle with vein R₅ and mouth of 1st submarginal cell over 1/2 as long as mouth of 2nd submarginal cell (square); vein R₄ almost vertical to vein R₅ and mouth of 1st submarginal cell less than 1/2 as long as mouth of 2nd submarginal cell (circle).

6. Antennal flagellomeres 2-5 gradually tapering apically and last joint (flagellomere 8 or sometimes 6) shorter than rest of flagellum (square); flagellomere 1 (which is distinctly longer than wide) with a long arista (which is 3-segmented) (Figs. 80, 268) (circle).

7. "The labella are reduced, laterally compressed, and joined over about one-half of their length (1996b: fig. 8 [= Fig. 191 right]; 1960: fig. 7) [triangle]; the labella are in the form of elongate, dorsoventrally somewhat flattened tubes, separated over their entire length though normally held pressed together (1996b: fig. 9 [= Fig. 115]; 1960: fig. 8) [triangle]. These conditions are interpreted as different autapomorphies that distinguish *Vermipardus* from the other clades collectively, and demonstrate their monophyly (1996b: 245)."

8. "6-7 pseudotracheae in a dendritic cluster in each labellum [square]; the main pseudotrachea in each labellum has only either two or three branches, and forms a relatively elongate entity corresponding to the modified form of the labellum [circle] (1996b: 245)."

9. "Aedeagus a simple, tubular organ without lateral accessory processes (1960: 227) [see aedeagus in Figs. 195-249] [square]; aedeagus nearly always with a pair of lateral accessory processes (1960: 227) [Figs. 4-6, 12-13, 18, 21, 37-38, 42, 59-60, 62-63, 65-66, 68-69, 118-120] [circle]." In *Lampromyia* (cylindrica group), aedeagus has no lateral projection as a result of reversal.

10. Pseudotrachea with two branches in *Lampromyia* (1996b: Fig. 274) (triangle); pseudotrachea with three branches in *Vermilynx* and *Leptynoma* (1996b: Fig. 275) (triangle). "While it appears to be a parsimonious argument that the two-branched condition could be derived from the three-branched and is thus more apomorphic, no evidence exists for such a progression, so these two conditions are regarded as unlinked autapomorphies indicating two monophyletic lineages (1996b: 245)."

11. "Hind tibia is clearly narrower apically than the hind femur [square]; hind tibia is considerably swollen apically and is enlarged as the hind femur....though it is not so marked in the female [of *Vermilynx*] [circle] (1996b: 245-246)."

12. "Proboscis is shorter (about 1.3-1.6 times mesonotal length), directed rigidly downwards, and is apparently incapable of forward or posterior flexure [square]; proboscis is extremely elongate and slender [Figs. 2-3]; at rest, the proboscis is folded beneath the body, directed obliquely backwards between the legs. In addition, the flies are able to flex the proboscis forwards when feeding at flowers [circle] (1996b: 246)."

13. "Wing form is unremarkable and generalised [square]; wing has an apomorphic form (figs. 14-19) [see notes under the genus *Leptynoma*] [circle] (1996b: 247)."

14. "No marked apomorphies in the radial field [square]; R₂₊₃ shows a variety of strongly apomorphic conditions, with each group having its own specialisations....directed in the key [circle] (1996b: 247)."

16. Possible phylogenetic tree in *Vermiphis* species

Vermiphis tibetensis differs from other species of the genus by having last joint of antennal flagellum shorter than preceding two joints together. However, it is undetermined whether this antennal character is apomorphic or plesiomorphic. Synapomorphic and autapomorphic characters can be seen only in the structure of male genitalia at present. There are definite two species groups, that is, A (= *ganquanensis* + *minshanensis* + *wudangensis*) and B (= *taishanensis* + *yanshanensis*). Furthermore, there are two peculiar species, that is, C (= *taihangensis*) and D (= *tibetensis*). *V. wudangensis* is unusual among the A group.

For A, B, C, D, and *V. wudangensis* above, synapomorphic or autapomorphic character state is given below separately.

Two combinations of species groups are easily supposed and they are (C+D) and (B+D), of which one must be rejected if the other stands. A reversal of the character state inevitably occurred in one of the two cases. It is unknown which case is the correct one.

Four possible phylogenetic trees are conceivable (see Figs. 417-420). However, no synapomorphic character state has been detected for (B+C+D), (A+B) or (A+C). It is suggested that (B+C+D) is the stronger possibility than (A+B) or (A+C), because the reversal of character in question seems to be easier in (B+C+D).

1. Synapomorphic character for A (= *ganquanensis* + *minshanensis* + *wudangensis*): Dorsomedial posterior projection of gonocoxite triangular and pointed apically.

2. Autapomorphic character for *wudangensis*: Dorsomedial posterior projection of gonocoxite conspicuous and demarcated from rest of gonocoxite.

3. Synapomorphic character for B (= *taishanensis* + *yanshanensis*): Aedeagus with two dorsal projections (in lateral view), of which anterior one represents a transverse band in dorsal view.

4. Possible synapomorphic character for C (= *taihangensis*) + D (= *tibetensis*): Tergum 9 with a mid posterior large ventral plane directed anteriorly.

When (B+C+D) is formed, a reversal occurred and the synapomorphic character state for (B+D) disappeared at C.

5. Possible synapomorphic character for B (= *taishanensis* + *yanshanensis*) + D (= *tibetensis*): Aedeagus with two dorsal projections (in lateral view), of which anterior one represents a transverse band in dorsal view. When (C+B+D) is formed, a reversal occurred and the synapomorphic character state for (C+D) disappeared at B.

6. Autapomorphic character for *taihangensis* (= C): Posterior curved part of aedeagus (in lateral view) with apical process directed anteriorly, which is rectangular (somewhat trapezoid) in dorsal view; dorsomedial posterior corner of gonocoxite with a process directed ventrally.

7. Autapomorphic character for *tibetensis* (= D): Posterolateral ventral part of gonocoxite long and narrow; within aedeagus behind ejaculatory apodeme there are paired longitudinal *strong* sclerites.

17. Evaluation of four character states

1. Antennal flagellum with a long arista or with last joint roughly as long as or longer than rest of flagellum: This character state is certainly apomorphic. It would occur four times, i. e. at *Vermileo*, *Vermitigris*, *Lampromyia* and *Leptynoma*. It is certain that *Vermipardus* + *Vermilynx* + *Leptynoma* + *Lampromyia* is not paraphyletic but monophyletic, because they are compactly built with one another by having two definite synapomorphic characters (see nos. 1-2 in Fig. 416).

2. Hind coxa with an elongate ventral (= anterior) process: Of the genera of Vermileonidae, only *Vermileo* (at least in North American *V. comstocki* and *V. opacus*) has this character state. It is judged at first sight as apomorphic. However, the presence of knob-like process on hind coxa is widely distributed in a number of families of orthorrhaphous Brachycera. It is possibly plesiomorphic in Vermileonidae too. If so, it would be lost two times, i. e. at *Vermipardus* + *Vermilynx* + *Lampromyia* + *Leptynoma* and at *Vermiophis* + *Vermitigris*. If not, it would be developed secondarily in *Vermileo*.

3. Palpal segment 2 widest around middle and then tapering apically: (1) This character is seen in *Vermiophis* and *Vermitigris*. There are two other states, that is, (2) swollen in lateral view and without

an elongate portion gradually tapering apically in *Vermileo* (at least in *V. comstocki* and *V. opacus*) and (3) clavate in *Lampromyia*, *Leptynoma*, *Vermilynx* and *Vermipardus*. The character state (1) is apparently plesiomorphic relative to (2) but is undetermined to be apomorphic or plesiomorphic relative to (3). Nagatomi (1981: 400) wrote: The palpus of the lower Brachycera is broadly classified here into two shapes. *A*: tapering distally at least in apical portion and pointed at apex, and curved downward or more or less so; *B*: log-like, knob-like, leaf-like, spatulate. The *A*-shape is seen in *Rhagio* (as well as *Neorhagio* and *Rhagina*), Athericidae, and female Tabanidae. Sometimes male Tabanidae may have an *A* palpus.

4. Presence of vein R_2 in *Leptynoma* (*Leptynoma*) *namaquaensis*: Stuckenberg (1996b: 252-253) wrote: This angled form was evidently the precursor of the remarkable condition in *namaquaensis* (figs. 17, 18) [or Fig. 57], in which vein R_2 and R_3 both seem to be present; distal extension of a stump vein and its forward curvature to the costa could result in a complete-spurious vein distal to R_{2+3} . This seemingly primitive venation with five radial veins, not otherwise recorded in Brachycera, was discussed previously (Stuckenberg, 1960). Its possibly primitive nature was mooted, but the condition is now seen to be secondary and highly apomorphic; a four-branched radial section is unknown in fossil Brachycera, even in the ancient Alinkidae from the Upper Triassic (Krzemiński, 1992) and Rhagionidae from the Lower and Middle Jurassic (Kovalev, 1982). Also, as the cladogram for this subgenus reveals, *namaquaensis* is one of the youngest and most derived of the species.

18. Larval head skeleton

Wheeler (1930: 199, 201) wrote: Comparison of *Vermileo* and *Lampromyia* with the few other Rhagionids that have been studied in detail (*Atherix*, *Rhagio*, *Ptiolina*) reveals numerous resemblances, but also striking differences, as will be found by the curious reader who cares to consult the figures of Brauer, Becker and de Meijere,.....

Teskey (1981: 532) wrote: The larval mouthparts of the Vermileonidae bear remarkable similarities to some of the Empidoidea (compare fig. 4 with figs. 48.41 and 48.43), especially in the configuration of the mandibular-maxillary sclerites and the membranization of the lateral covering of the head that bears the antennae and maxillary palpi. The retracted cranial portions of the head capsule have retained a more primitive, extensively sclerotized condition.

The following are quoted from Sinclair (1992) who made a study on mandible of *Vermileo comstocki* and *V. opacus*.

The mandible of this family is of similar structure to that of the ground plan of the Brachycera. The mandible is composed of two components, consisting of a distal hook with a serrated posterior margin and an inverted 'U'-shaped basal sclerite, lacking the mandibular brush. Contrary to Teskey (1981b), the mandible of Vermileonidae is not similar to that of the Empidoidea (see Eremoneura ground plan).

[In *Tabanomorpha* (Rhagionidae, Pelecorhynchidae, Athericidae and Tabanidae)] mandible with brush on posterior margin of external arm of basal sclerite (apomorphic). [In Pelecorhynchidae + Athericidae + Tabanidae] mandibular brush associated with an articulated rod (apomorphic). [In Rhagionidae] mandibular brush associated with a simple fold of cuticle (plesiomorphic). [In Empidoidea] mandible of final instar larvae comprised of four components (apomorphic). [As ground plan of Brachycera including Rhagionidae and Vermileonidae] mandible of final instar larvae

composed of two components (plesiomorphic).

In the plesiomorphic state, the maxilla is composed of well-defined sclerites, articulated posteriorly with the basal mandibular sclerite (figs. 8, 10). In the *Eremoneura*, the maxilla is an elongate, primarily membranous lobe, delimited at its base by the antenna (fig. 19; Teskey, 1981a, figs. 3.12-13). This character is proposed as a possible synapomorphy of the *Eremoneura*. Elongate maxillae with similarly placed antennae in Vermileonidae are probably the result of convergence ([see] Teskey, 1981b, fig. 39.4).

Character 20. Labium composed of a large rectangular plate (plesiomorphic) / labial hypopharyngeal sclerite 'V'-shaped (apomorphic). The plesiomorphic condition occurs in the Tabanomorpha and Asiloidea, where the labium is composed of a large rectangular submental plate. The apomorphic state, a 'V'-shaped labial hypopharyngeal sclerite, is present in the Empidoidea (Vaillant, 1948, fig. 16) and retained in primitive Cyclorrhapha (e. g. *Melanderomyia*, *Megaselia*). This character was first proposed as a possible synapomorphy of the *Eremoneura* by Hennig (1952) and adopted by Griffiths (1972).

Tsacas (1962: 231) on *Vermileo vermileo*: Le labium, très réduit chez *Vermileo*, est muni de palpes très courts et de soies bien développées. L'hypopharynx, par sa forme en Y, rappelle la pièce en V des Empidiformia. Le salivarium, par la sclérotisation de sa moitié basale et l'absence d'une pompe salivaire individualisée, montre des particularités évidentes. L'ensemble de des trois organes, labium, hypopharynx, salivarium, fait penser plutôt aux Empidiformia, aussi étonnant que cela puisse paraître.

The maxillary palpus (Figs. 351-356) identified by Engel (1929), Wheeler (1930) and Tsacas (1962) was interpreted as antenna by Teskey (1981) (Fig. 358). According to Tsacas (1962), antenna is short and inconspicuous in *Vermileo vermileo* (see Figs. 359-361) but long and conspicuous in *Rhagio scolopaceus* and *Chrysopilus auratus* (see Figs. 364-365, 371-372). The antenna of *Vermileo comstocki* was evidently misidentified by Teskey (1981), as pointed out already by Ludwig *et al.* (1996).

Of the two synapomorphic character states (shown in italics) for *Eremoneura* proposed by Sinclair, one (as to labial hypopharyngeal sclerite) may also be common to Vermileonidae but the other (as to maxilla and antenna) would be unsuitable for Vermileonidae, when the antenna of Vermileonidae was misidentified by Teskey. Nevertheless, there may be some similarity in the structure and position of maxilla and true antenna between Vermileonidae and *Eremoneura*.

It should be noted that the Vermileonidae is neither *Eremoneura* nor Empidoidea in assignment but has been judged as a sister group of Rhagionempididae (= Apsilocephalidae) + *Eremoneura* by Nagatomi *et al.* (1991) and Nagatomi (1992, 1996).

19. Presence of poison canal within larval mandible

Woodley (1989: 1380) wrote: Tabanidae (Ch. 31), Athericidae (Webb, 1977), and Pelecorhynchidae (Teskey, 1970, for *Glutops*), based on what we know of their larvae, all possess a canal in the center of each mandible, used for injecting poison into their prey. Known larvae of Rhagionidae (Ch. 33) have no such poison canal. This apomorphic feature indicates a monophyletic origin of the three families that possess the poison canal. Stuckenberg (1973: 669) gives a brief summary of this character.

Ludwig *et al.* (1996: 9) on *Vermileo vermileo*: "The mandibles are channeled on the inner side and allow the injection of saliva and to sucking out of the digested prey."

It seems that the poison canal in Vermileonidae occurs independently from that in Pelecorhynchidae, Athericidae and Tabanidae.

20. Larval and pupal spiracles

There are many illustrations for spiracles of Diptera in several publications, e. g. McAlpine *et al.* (1981, 1987), Ferrar (1987), and Smith (1989) where the numbers are smaller in Nematocera and orthorrhaphous Brachycera than in Cyclorrhapha. Keilin (1944), Whitten (1960), Teskey (1981a), etc. discussed or synthesized the respiratory systems and spiracles of Diptera. Their taxonomic importance is distinct but the phylogenetic significance becomes unclear, if their form changes considerably for adaptation. Regrettably neither personal study nor review of the literature has been made by us and no comment is possible on spiracles for taxonomic importance and phylogenetic significance within Vermileonidae or orthorrhaphous Brachycera.

21. Antennal flagellum, etc., phylogenetically important

Antennal flagellum is tapering apically, long and with segments often reduced in number and often becoming a terminal segmented arista (Figs. 276-293, etc.). But in *Vermilynx (jasoni)* and *vansoni* and some *Vermipardus* the shape of antennal flagellum is similar to or difficult to separate from that of Xylomyidae, Coenomyiidae, Rhagionidae, Pelecorhynchidae, Tabanidae, etc. In *Vermilynx (jasoni)* and *vansoni* the male antennal flagellum is shorter than the female. Probably the male is apomorphic relating to the female in this respect.

Nagatomi (1996: 131) wrote: "Vermileonidae has the following apomorphic characters: female cercus possibly 1-segmented at least in some species of *Vermileo* and *Lampromyia*; antennal flagellum often 3-4 segmented; in *Lampromyia pallida* Macquart, empodium absent (after Verrall, 1909); in *Vermileo comstocki* Wheeler and *Lampromyia (Vermipardus) brevirostris* Bezzi, the lateral part of male tergum 9 protrudes posteriorly. Each of these characters may indicate a transitional sign towards the Apsilocephalidae + *Eremoneura*."

However, the above characters occurred within the Vermileonidae and are never synapomorphic attributes of Vermileonidae + Rhagionempididae (= Apsilocephalidae) + *Eremoneura*.

Let us suppose that there is one stock including Vermileonidae, and a shoot from this stock is the Rhagionempididae (= Apsilocephalidae) + *Eremoneura* which has the following common characters: typical antennal flagellum widely distributed in Vermileonidae, Rhagionempididae, Empidoidea and primitive Cyclorrhapha; larval head skeleton mentioned in a separate section; 1-segmented female cercus; empodium bristle-like, triangular at apical portion, or absent; male tergum 9 with lateral part protruded posteriorly; etc.

There must be another stock different from the one including Vermileonidae, and from this stock the Nemestrinoidea + Asiloidea or Asiloidea (if Nemestrinoidea is a sister group of Tabanoidea) is derived.

22. Synapomorphic and autapomorphic characters

The Vermileonidae is most closely related phenetically (not phylogenetically) to the Tabanoidea by having the following characters (which are plesiomorphic relating to Nemestrinoidea and Asiloidea): tibial spurs (1 : 2 : 2 or rarely 0 : 2 : 2) always present; three pad-like pulvilli and empodium present (with a few exceptions); antennal flagellum with five or more segments in some genera; female cercus

2-segmented (sometimes possibly 1-segmented).

Its synapomorphic characters (relating to Tabanoidea) are as follows: differentiation of clypeus, parafacials and upper face is indistinct and clypeus occupies almost whole surface of face; hind leg markedly long; abdomen slender and clavate. Its autapomorphic characters are given below. Adult: scutellum flattened dorsally and plank-like. Larva: see P. 76 in this paper.

23. Some notes to Nagatomi (1996) on phylogeny of the orthorrhaphous Brachycera

Pp. 97, 104, 129, for '*Vermileomorpha*', read '*Vermileonomorpha*'.

P. 121, line 13 (from bottom), for 'Tergum 10 is absent in Stratiomyoidea', read 'Tergum 10 is absent in Xylomyidae, Pantophthalmidae, but present (well developed and produced posteriorly) in Stratiomyidae.'

There are two facts which may exert against the scheme Vermileonidae + Rhagionempididae (= Apsilocephalidae) + Empidoidea + Cyclorrhapha.

(1) The apical portion of empodium becomes triangular in shape or bristle-like, or is sometimes absent. This may be a synapomorphic character state for Asiloidea + Rhagionempididae (= Apsilocephalidae) + Empidoidea + Cyclorrhapha. However, one of us (A. N.) thinks that it occurred three times independently at *Vermilynx vansoni*, Rhagionempididae + Empidoidea + Cyclorrhapha, and Asiloidea.

(2) The male genitalia are different conspicuously among three groups, that is, Vermileonidae, Rhagionempididae, and Empidoidea + Cyclorrhapha, and the phallus of Rhagionempididae resembles that of Asilidae. However, they are also different markedly in several parts (besides male genitalia). The resemblance of phallus between Rhagionempididae and Asilidae seems to be a result of parallelism. It should be noted that the Vermileonidae is one of the lower forms but the Empidoidea is the highest one among the orthorrhaphous Brachycera.

More comparison is needed for larval head structures among Vermileonidae, Rhagionempididae and Empidoidea. It is hoped that the appearance of new data and the development of molecular biology in the future can clarify the position of questionable taxa of the orthorrhaphous Brachycera including Vermileonidae and Rhagionempididae.

24. Sister group of Vermileonidae

The Vermileonidae had been treated as a subfamily of Rhagionidae. It becomes recognized as an independent family since the work of Nagatomi (1977), Teskey (1981b) and Woodley (1989). The sister group of Vermileonidae seems to be the Rhagionempididae (= Apsilocephalidae) + Empidoidea + Cyclorrhapha, as discussed by Nagatomi *et al.* (1991) and Nagatomi (1992, 1996). More detailed explanations for larval mouthparts, antennal flagellum, etc. are given separately in this paper.

ACKNOWLEDGEMENTS This paper is dedicated to Dr. Brian R. Stuckenberg (Pietermaritzburg, South Africa) for the advancement in knowledge of the Vermileonidae through his many fine contributions.

Our heart-felt thanks must be expressed to the previous authors whose passages and illustrations are reproduced here. Without them, the value of this paper would be markedly reduced.

Grateful acknowledgement is made to Mr. K. G. V. Smith (London, Editor in Chief,

Entomologist's Monthly Magazine) for his critical reading of the manuscript.

We are also much indebted to Mr. Fasheng Li (China Agricultural University, Beijing), Mr. H. Chen (Wuhan, China), Professor Io Chou (Northwest Agricultural University, Shaanxi) for their help in many ways. This study is supported partly by the National Natural Science Foundation of China.

Postscript

There are many undescribed Vermileonidae at Dr. Stuckenberg's disposal. They are introduced below, based on the information from his letters to one of us (A. N.).

(On July 21, 1997): The vermilionid fauna of South Africa is proving to be much larger than previously suspected, and there will certainly be even more new species discovered when searches are made in the arid interior of this country, as well as in mountain ranges in the north and north-west.

(On March 2, 1998): New discoveries of other interesting Vermileonidae continue to be made! I have just finished a manuscript in which I am describing a wonderful new species of *Leptynoma* s. str. found in Namibia. It is the sister-species of *Leptynoma* (*L.*) *namaquaensis*, and has elaborated the unusual radial venation of that species by further modifications. In this paper I am presenting a biographical analysis of *Leptynoma* s. str. (now 5 species),

The first known vermilionid [probably *Lampromyia* sp.] in the Arabian peninsula was recently found (in Oman)!

(On April 7, 1998): I also have the strange new species from Namaqualand on the western side of South Africa, which is probably related to *Leptynoma* but has some extraordinary features and may require a new genus. One of its peculiarities was that the flies emerged in mid-winter, which no other vermilionid is known to do in South Africa.

(On October 13, 1998): *Vermipardus* continues to grow; This genus must certainly have at least 20 species, but possibly even 25 species, as there are still extensive parts of South Africa where no collecting of vermilionids has yet been done.

I also have new species of *Vermilynx* and *Perianthomyia* to describe,

It is also now clear that the palaearctic *Vermileo* badly need revision. I now have a new species from Malta, and it seems that the total number of species might be at least twice what are already described.

REFERENCES

- Becker, T. 1921. Neue Dipteren Meiner Sammlung. *N. Beitr. Syst. Insektenk., Berlin* 2: 41-48.
 Bequaert, J. 1922. The predaceous enemies of Ants. *Bull. Amer. Mus. Nat. Hist.* 45: 271-331.
 Bezzi, M. 1926. South African Rhagionidae (Diptera) in the South African Museum. *Ann. S. Afr. Mus.* 23: 297-324.
 Biró, L. 1897. Biologische Mitteilungen aus Neu Guinea. I. Die Ameisen-Löwen in Neu Guinea. *Berl. Ent. Zeitschr.* 42: 129-138.
 Brauer, F. 1883. *Vermileo Degeeri* Macq. *Wien. ent. Ztg.* 2: 114.
 Brauns, A. 1954. *Puppen terricoler Dipterenlarven*. 156 pp. Musterschmidt. Göttingen.
 Brunetti, E. 1927. Notes on Malayan Diptera, with descriptions of new species. *J. fed. Malay. St. Mus.* 13: 281-309.

- Buchner, P. 1940. Über den Wurmlöwen (*Vermileo vermileo*). *Natur. u. Volk*, Frankfurt a. M. **70**: 116-131.
- Chassagnard, M. T. & Tsacas, L. 1974. Morphologie de la tête et des pièces buccales imaginaires de *Chrysopilus auratus* F. et de *Vermileo vermileo* de G. (Diptera: Rhagionidae). *International J. Insect Morph. Embryol.* **3** (1): 13-32.
- De Geer, G. 1752. Rön om Mask-Lejonet. *Vetensk. Akad. Handl.* 1752: 180-192, 261-265.
- Deleon, D. 1938. Notes on the distribution of *Vermileo* in the United States and Mexico with a description of a new species (Diptera: Rhagionidae). *Entomol. News* **49**: 167-171.
- Edwards, F. W. 1932. A note on the genus *Vermilegrus* Wheeler (Diptera, Rhagionidae). *Stylops* **1**: 217-220.
- . 1935. "Worm-lions" from Portugal and the Sudan (Diptera, Rhagionidae), with a description of a new species. *Proc. R. ent. Soc. Lond.* **10**: 54-57.
- Engel, E. O. 1929. Notes on two larvae of South African Diptera belonging to the families Leptidae and Asilidae. *Trans. R. soc. S. Afr.* **18**: 147-162.
- Evenhuis, N. E. 1994. *Catalogue of the fossil flies of the world (Insecta: Diptera)*. 600 pp. Backhuys Publishers, Leiden.
- Falcoz, L. 1927. Observations biologiques sur divers insectes des environs de Vienne en Dauphiné. *Bull. Soc. Ent. France*, pp. 226-232.
- Fernández, J. M. 1952. La *Lampromyia canariensis* Macq. *Graellsia* **10**: 47-55. (In Spanish)
- Ferrar, P. 1987. A guide to the breeding habits and immature stages of Diptera Cyclorrhapha. *Entomograph* **8** (1, text): 1-478; (2, figs): 479-907.
- Frederiksen, J. & Hemmingsen, A. M. 1972. Ethological and autecological studies on Canarian Vermileoninae (Diptera Brachycera, Rhagionidae= Leptidae). *Vidensk. Meddr dansk naturh. Foren.* (1972) **135**: 37-59.
- Frey, R. 1937. Die Dipterenfauna der Kanarischen Inseln und ihre Probleme. *Soc. Scient. Fenn. Comm. Biol.* **6** (1): 1-237.
- Greene, C. T. 1926. Descriptions of larvae and pupae of two-winged flies belonging to the family Leptidae. *Proc. U. S. Nat. Mus.* **70**: 1-20.
- Griffiths, G. C. D. 1972. The phylogenetic classification of Diptera Cyclorrhapha, with special reference to the structure of the male postabdomen. *Series Ent.* **8**: 1-340.
- Hafez, M. & El-Moursy, A. A. 1956a. Studies on desert insects in Egypt. I. Field and laboratory investigations on the worm-lion, *Vermileo vermileo* L. [= possibly *Vermileo niloticus* Edwards] (Diptera: Rhagionidae). *Bull. Soc. Entom. Egypte* **15**: 279-299.
- & ———. 1956b. Studies on desert insects in Egypt. II. On the general biology of *Vermileo vermileo* L. [= possibly *Vermileo niloticus* Edwards] (Diptera: Rhagionidae). *Bull. Soc. Entom. Egypte* **15**: 333-348.
- & ———. 1964. Comparative notes on the habits of the pit-building worm-lions and ant-lions. *Bull. Soc. Entom. Egypte* **47**: 125-132.
- Hemmingsen, A. M. 1963. The ant-lion-like trap of the larva of *Lampromyia canariensis* Macquart [= correctly *Lampromyia fortunata* Stuckenberg] (Diptera, Leptidae= Rhagionidae, Vermileoninae). *Vidensk. Meddr dansk naturh. Foren.* **125**: 237-267.
- . 1968. A review of instinctive behaviour in the worm-lions *Vermileo vermileo* L. and

- Lampromyia pallida* Macquart (Diptera Brachycera, Rhagionidae, Vermileoninae). *Vidensk. Meddr dansk naturh. Foren.* **131**: 289-302.
- . 1977. Ormeløvestudier. *Ent. Meddr.* **45**: 167-188.
- . & Nielsen, B. R. 1971. Species differences in ovipository instincts within the Vermileoninae (Diptera Brachycera, Rhagionidae= Leptidae). *Vidensk. Meddr dansk naturh. Foren.* **134**: 149-203.
- Hennig, W. 1952. *Die Larvenformen der Dipteren*. Vol. **3**. 628 pp. Akademie-Verlag, Berlin.
- . 1967. Die sogenannten "niederen Brachycera" im Baltischen Bernstein (Diptera: Fam. Xylophagidae, Xylomyidae, Rhagionidae, Tabanidae). *Stuttgarter Beitr. Naturkunde* **174**: 1-51.
- James, M. T. 1965. Family Rhagionidae. pp. 342-348. In: Stone, A., Sabrosky C. W., Wirth, W. W., Foote, R. H., and Coulson, J. R. (eds.), *A catalog of the Diptera of America north of Mexico*. 1696 pp. Washington, D. C.
- . 1968. 29. Family Rhagionidae (Leptidae). pp. 1-12. In: Dept. Zool. Secretaria da Agricultura (ed.), *A catalogue of the Diptera of the Americas south of the United States*. São Paulo.
- Keilin, D. 1944. Respiratory systems and respiratory adaptations in larvae and pupae of Diptera. *Parasitology* **36**: 1-66.
- Kertész, K. 1908. *Catalogus dipterorum hucusque descriptorum*. Vol. **3**. 367 pp. Leipzig, Budapest.
- Le Fauchaux, M. 1961. Contribution à l'étude du cycle biologique de *Vermileo Degeeri* Macquart (Diptera Rhagionidae): Ponte et éclosion des jeunes larves. *Bull. Soc. Scientif. Bretagne* **36**: 133-141.
- Leonard, M. D. 1930. A revision of the Dipterous family Rhagionidae (Leptidae) in the United States and Canada. *Mem. Am. Ent. Soc.* **7**: 1-181.
- Lindner, E. 1925. 20. Rhagionidae. 49 pp. In: Lindner, E. (ed.), *Die Fliegen der paläarktischen Region*. Stuttgart.
- Ludwig, P., Smola, U. & Melzer, R. 1996. Die Mundwerkzeuge des Wurmlöwen *Vermileo vermileo* L. und ihre Funktion (Diptera, Vermileonidae). *NachrBl. bayer. Ent.* **45** (1/2): 9-14.
- Majer, J. 1988. Family Vermileonidae. pp. 34-35. In: Soós, A. & Papp, L. (eds.), *Catalogue of Palaearctic Diptera*. Vol. **5**. 446 pp. Akadémiai Kiadó, Budapest.
- Mansell, M. W. 1985. [Chapter] 19. Neuroptera. In: Scholtz, C. H. & Holm, E. (eds.), *Insect of Southern Africa*. p. 186. Durban: Butterworths.
- Marchal, L. P. 1897. Notes d'entomologie biologique sur une excursion en Algérie et en Tunisie. *Mém. Soc. Zool. de Fr.* **10**: 5-25.
- Martin, C. H. 1968. The new family Leptogastridae (the grass flies) compared with the Asilidae (Robber flies) (Diptera). *J. Kansas Entomol. Soc.* **41**: 70-100.
- Matsura, T. 1989. 'The antlions of sandhills.' 215 pp. Shisaku-sha, Tokyo. (In Japanese)
- . 1997. Myrmeleontidae. p. 14. In: Ishii, M., Ohtani, T. & Johki, Y. (eds.), *The encyclopedia of animals in Japan*, Vol. **9**, Insects II. 181 pp. Heibonsha, Tokyo. (In Japanese)
- McAlpine, J. F., Peterson, B. V., Shewell, G. E., Teskey, H. J., Vockeroth, J. R. & Wood, D. M. 1981. *Manual of Nearctic Diptera*. Vol. **1**. Research Branch Agriculture Canada. Monograph No. 27. 674 pp.
- , ———, ———, ———, ——— & ———. 1987. *Manual of Nearctic Diptera*. Vol. **2**. Research Branch Agriculture Canada. Monograph No. 28. pp. 675-1332.

- Morisita, M. 1952. Habitat preference and evaluation of environment of an animal. Experimental studies on the population density of an antlion, *Glenuroides japonicus* M'L. [= correctly *Hagenomyia micans*]. I. *Physiology and Ecology* 5 (1): 1-16. (In Japanese with English summary)
- Mostovski, M. B. 1997. To the knowledge of fossil Dipterans of superfamily Archisargoidea (Diptera, Brachycera). *Paleontol. J.* No. 1, pp. 72-77. (In Russian with English summary)
- Nagatomi, A. 1975. Family Rhagionidae (Leptidae). pp. 82-90. In: Delfinado, M. D. & Hardy, D. E. (eds.), *A catalog of the Diptera of the Oriental Region*. Vol. 2. 459 pp. Univ. Press of Hawaii, Honolulu.
- . 1977. Classification of lower Brachycera (Diptera). *J. nat. Hist.* 11: 321-335.
- . 1981. Some characters of the lower Brachycera (Diptera) and their plesiomorphy and apomorphy. *Kontyu*, Tokyo 49: 397-407.
- . 1982. Geographical distribution of the lower Brachycera (Diptera). *Pacific Insects* 24: 139-150.
- . 1984. Male genitalia of the lower Brachycera (Diptera). *Beitr. Ent.*, Berlin 34: 99-157.
- . 1991. History of some families of Diptera, chiefly those of the lower Brachycera (Insecta: Diptera). *Bull. Biogeogr. Soc. Japan* 46: 21-37.
- . 1992. Notes on the phylogeny of various taxa of the orthorrhaphous Brachycera (Insecta: Diptera). *Zool. Sci.* 9: 843-857.
- . 1996. An essay on phylogeny of the orthorrhaphous Brachycera (Diptera). *Entomol. Mon. Mag.* 132: 95-148.
- . 1997. Vermileonidae. pp. 447-458. In: Papp, L. & Darvas, B. (eds.), *Contribution to a Manual of Palaearctic Diptera*. Vol. 2. Nematocera and lower Brachycera. 592 pp. Science Herald, Budapest.
- . & Iwata, K. 1976. Female terminalia of lower Brachycera (Diptera), I. *Beitr. Ent.*, Berlin 26: 5-47.
- . & Soroida, K. 1985. The structure of the mouthparts of the orthorrhaphous Brachycera (Diptera) with special reference to blood-sucking. *Beitr. Ent.*, Berlin 35: 263-368.
- . & Yang, D. 1998. A review of extinct Mesozoic genera and families of Brachycera (Insecta, Diptera, Orthorrhapha). *Entomol. Mon. Mag.* 134: 95-192.
- , Saigusa, T., Nagatomi, H. & Lyneborg, L. 1991. The systematic position of the Apsilocephalidae, Rhagionempididae, Protrepididae, Hilarimorphidae, Vermileonidae and some genera of Bombyliidae (Insecta, Diptera). *Zool. Sci.* 8: 593-607.
- Oldroyd, H. 1947. A new *Vermitigris* (Diptera: Rhagionidae) from Borneo. *Proc. R. ent. Soc. Lond.* (Ser. B) 16: 21-22.
- Osten Sacken, C. R. 1883. Synonymica concerning exotic dipterology. No. II. *Berl. ent. Zeit.* 27: 295-298.
- Pechuman, L. L. 1938. A synopsis of the new world species of *Vermileo* (Diptera-Rhagionidae). *Bull. Brooklyn Entomol. Soc.* 33: 84-89.
- Peterson, B. V. 1981. Apioceridae. pp. 541-548. In: McAlpine, J. F., Peterson, B. V., Shewell, G. E., Teskey, H. J., Vockeroth, J. R. & Wood, D. M. (eds.), *Manual of Nearctic Diptera*. Vol. 1. Research Branch Agriculture Canada, Monograph No. 27. 674 pp.
- de Réaumur, R. A. F. 1753. Histoire du ver lion mouche. *Mém. Acad. Sc., Paris* 1753: 402-419. 1 pl.

- Ed. 1762: 604-631.
- Rohdendorf, B. R. 1938. [Dipterous insects of the Mesozoic Karatau. I. Brachycera and Nematocera]. *Trudy Paleontol. Inst. Akad. Nauk SSSR* 7 (3): 29-67. (In Russian)
- de Romand, B. E. 1833. Note sur la larve du *Leptis vermileo*. *Ann. Soc. Ent. France* 2 Bull.: 498-499.
- Séguy, E. 1926. *Faune de France*. 13. Diptères (Brachycères). 308 pp. Paul Lechevalier. Paris.
- . 1928. Diptères nouveaux de l'Afrique mineure. *Bull. Soc. ent. France* (1928): 45-46.
- . 1930. Contributions a l'étude des diptères du Maroc. *Mém. Soc. Sci. nat. phys. du Maroc*. 24: 1-206.
- Shelford, R. S. 1901. Notes on some Bornean insects. *Rep. Brit. Assoc. Section D. Glasgow*, pp. 689-691.
- von Siebold, C. T. E. 1861. Ueber die Larve von *Leptis vermileo*. *Amtl. Ber. 35. Versam. Deutsch. Naturf. u. Aerzte in Königsberg* (1860), pp. 105-107.
- Sinclair, B. J. 1992. A phylogenetic interpretation of the Brachycera (Diptera) based on the larval mandible and associated mouthpart structures. *Syst. Entomol.* 17: 233-252.
- , Cumming, J. M. & Wood, D. M. 1994. Homology and phylogenetic implications of male genitalia in Diptera-Lower Brachycera. *Entomologica scand.* 24: 407-432.
- Smith, K. G. V. 1989. An introduction to the immature stages of British flies. *Handbk Ident. Br. Insects* 10 (14): 1-280.
- Struck, M. 1992. Pollination ecology in the arid winter rainfall region of southern Africa: a case study. *Mitteilungen des Institutes für Allgemeine Botanik* 24: 61-90.
- Stuckenberg, B. R. 1955. New and little-known South African Rhagionidae (Diptera). *J. Ent. Soc. S. Africa* 18: 255-265.
- . 1960. Diptera (Brachycera): Rhagionidae. *S. Afr. anim. Life* 7: 216-308.
- . 1961. Records and descriptions of Blepharoceridae, Erinidae and Rhagionidae from South Africa (Diptera). *Ann. Natal Mus.* 15: 109-124.
- . 1965a. A new species of *Lampromyia* Macquart (Diptera: Rhagionidae) from the Richtersveld. *J. Ent. Soc. S. Africa* 28: 107-109.
- . 1965b. Notes on the Palaearctic species of *Vermileo*, with the description of a new species from Crete (Diptera: Rhagionidae). *Ann. Mag. nat. Hist.* (13) 8: 495-500.
- . 1971. A taxonomic study on the species of *Lampromyia* Macq. in the Canary Islands (Diptera: Rhagionidae). *Vidensk. Meddr dansk naturh. Foren.* (1971) 134: 77-88.
- . 1973. The Athericidae, a new family in the lower Brachycera (Diptera). *Ann. Natal Mus.* 21: 649-673.
- . 1980. 22. Family Rhagionidae. pp. 309-311. In: Crosskey, R. W. (ed.), *Catalogue of the Diptera of the Afrotropical Region*. 1437 pp. British Mus. (Nat. Hist.), London.
- . 1995a. *Vermilyn*, a new genus for the wormlion fly *Lampromyia vanson* Stuckenberg of the Richtersveld, Southern Africa (Diptera: Vermileonidae). *African Entomology* 3: 29-34.
- . 1995b. A taxonomic revision of *Vermipardus* Stuckenberg, 1960, with descriptions of new species and notes on the biology and biogeography of the genus (Diptera: Vermileonidae). *Ann. Natal Mus.* 36: 215-253.
- . 1996a. A new species of the wormlion genus *Vermilyn* Stuckenberg (Diptera: Vermileonidae) from Namaqualand, South Africa. *African Entomology* 4: 197-202.

- . 1996b. A revised generic classification of the wormlion flies of Southern Africa previously placed in *Lampromyia* Macquart, with reinstatement of *Leptynoma* Westwood 1876, and descriptions of a new subgenus and two new species (Diptera, Vermileonidae). *Ann. Natal Mus.* **37**: 239-266.
- . 1997. Three new species of *Vermipardus* Stuckenberg, and a revised cladogram for the genus with novel biogeographical implications (Diptera: Vermileonidae). *Ann. Natal Mus.* **38**: 173-196.
- . 1998. A revision of the Palaearctic species of *Lampromyia* Macquart (Diptera, Vermileonidae), with the description of a new Iberian species and a cladogram for the genus. *Bonn. zool. Beitr.* **48**: 67-96.
- Szilády, Z. 1934. Die palaearktischen Rhagioniden. *Ann. Mus. Nat. Hung.* **28**: 229-270.
- Teskey, H. J. 1970. The immature stages and phyletic position of *Glutops rossi* (Diptera: Pelecorhynchidae). *Can. Ent.* **102** (9): 1130-1135.
- . 1981a. Morphology and terminology — larvae [Chapter] 3. pp. 65-88. In: McAlpine, J. F., Peterson, B. V., Shewell, G. E., Teskey, H. J., Vockeroth, J. R. & Wood, D. M. (eds.), *Manual of Nearctic Diptera*. Vol. 1. Research Branch Agriculture Canada, Monograph No. 27, 674 pp.
- . 1981b. Vermileonidae. pp. 529-532. In: McAlpine, J. F., Peterson, B. V., Shewell, G. E., Teskey, H. J., Vockeroth, J. R. & Wood, D. M. (eds.), *Manual of Nearctic Diptera*. Vol. 1. Research Branch Agriculture Canada, Monograph No. 27, 674 pp.
- Tsacas, L. 1962. Recherches sur la structure et le fonctionnement de la tête et des pièces buccales larvaires des Rhagionidae (Diptères). *Mém. Mus. Nat. d'Hist. Naturelle. Série A, Zool.* **27**: 147-235.
- Vaillant, F. 1948. Les premiers stades de *Liancalus virens* Scop. (Dolichopodidae). *Bull. Soc. Zool. France* **73**: 118-126.
- Verrall, G. H. 1909. *British flies*. Vol. V. Stratiomyidae and succeeding families of the Diptera Brachycera of Great Britain. 780 pp. Gurney & Jackson. London.
- Webb, D. W. 1974. A revision of the genus *Hilarimorpha* (Diptera: Hilarimorphidae). *J. Kansas Entomol. Soc.* **47**: 172-222.
- . 1977. The Nearctic Athericidae (Insecta: Diptera). *J. Kans. Ent. Soc.* **50** (4): 473-495.
- Wheeler, W. M. 1930. *Demons of the dust*. 378 pp. Kegan Paul, Trench, Trubner & Co., London.
- . 1931. A Cuban *Vermileo*. *Psyche*, Camb. **38**: 165-169.
- . 1934. A specimen of the Jamaican *Vermileo*. *Psyche*, Camb. **41**: 236-237.
- Whitten, J. M. 1960. The tracheal system as a systematic character in larval Diptera. *Syst. Zool.* **8**: 130-139.
- Wilcox, J. 1981. Mydidae. pp. 533-540. In: McAlpine, J. F., Peterson, B. V., Shewell, G. E., Teskey, H. J., Vockeroth, J. R. & Wood, D. M. (eds.), *Manual of Nearctic Diptera*. Vol. 1. Research Branch Agriculture Canada, Monograph No. 27, 674 pp.
- Woodley, N. E. 1989. Phylogeny and classification of the "Orthorrhaphous" Brachycera. pp. 1371-1395. In: McAlpine, J. F. & Wood, D. M. (eds.), *Manual of Nearctic Diptera*. Vol. 3. Research Branch Agriculture Canada. Monograph No. 32. pp. 1333-1581.
- Yang, C. 1979a. A new genus and species of wormlion from China (Diptera: Rhagionidae). *Entomotaxon.* **1**: 83-89. (In Chinese with English summary)
- . 1979b. Biological observations on *Vermiophis ganquanensis* Yang. *Entomotaxon.* **1**: 131-140. (In Chinese)

- . 1988. A primary note on Chinese *Vermitigris* and a new species of Guangxi (Diptera: Vermileonidae). *Entomotaxon.* **10**: 178-182. (In Chinese with English summary)
- . & Chen, H. 1986. A new species of the genus *Vermiophis* from the Wudang Mountains, China (Diptera: Vermileonidae). *J. Huazhong Agric. Univ.* **5** (4): 321-325. (In Chinese with English summary)
- . & ———. 1987. Diptera: Vermileonidae. *Agricultural insects, spiders, plant diseases and weeds of Xizang* **2**: 157-160. (In Chinese with English summary)
- . & ———. 1993. Notes on Vermileonidae and four new species of *Vermiophis* (Diptera: Brachycera) from China. *Entomotaxon.* **15**: 127-136. (In Chinese with English summary)
- Yeates, D. K. 1994. The cladistics and classification of the Bombyliidae (Diptera: Asiloidea). *Bull. Am. Mus. Nat. Hist.* No. 219. 191 pp.
- Zimsen, E. 1964. *The type material of I. C. Fabricius*. 656 pp. Munksgaard, Copenhagen.

Received July 10, 1998

Accepted November 15, 1998

永富昭, 楊集昆, 楊定 中国と世界のアナアブ科 (双翅目)

アナアブ科は長くシギアブ科の一亜科とされたが、現在は独立の科となる。永富ら (1991) と永富 (1992, 1996) は本科とオドリバエ主科+ハエ亜目を姉妹関係にあるとする。オーストラリアと南米を除いた世界に不連続に分布する。1979年以降中国で2属8種が発見命名された。幼虫は砂中にすり鉢状の巣を作り、落ちた節足動物を捕食して、その生活は脈翅目のウスバカゲロウ科の一部 (アリジゴク) に似る。

Vermiophis は中国特産で、最も原始的な属である。*Vermitigris* は中国南部とチベット以外ではボルネオ、スマトラ、マレー半島、印度南部に分布する。

この総説は中国産全種の再記載と共に世界の7属55種を目録とする。地理的分布、化石種、習性、生活史、幼生期の形態について既往の論文中の多数の章句と付図を引用紹介する。若干の形質について分類学的、系統学的意義を論議する。

しかし分類、生態、生理のすべてにわたって得られた知識は少なく、今後の研究に待つところ大である。

INDEX OF TAXA

- Apogon* 16
dufouri 23
Arthrostylum 16, 136
Lampromyia 1-4, 10, 12, 25, 51-53, 61, 64-65, 67, 70-71, 78, 84, 86, 89-90, 92-93, 95-96, 98, 103, 114, 116, 118, 126, 129-133, 135-137, 139-141, 143, 145, 149-150
canariensis 4-5, 7, 10-11, 65, 67, 70-71, 76, 78-79, 83-84, 88, 90, 92-93, 95-98, 106, 112, 117-118, 132-133, 135, 146
cylindrica 4-5, 8-10, 11, 25, 65, 70, 118, 132-133, 135, 139
flavida 4-6, 10, 64, 70-71
fortunata 4, 7, 10-11, 65, 70-71, 76-78, 83, 92-98, 100, 105-109, 111-113, 115-117, 122, 135, 146
funnebris 5, 8, 11, 65, 70, 93, 95-96, 98, 133
hemmingseni 4, 7, 11, 65, 67, 70-71, 93, 95-98, 101, 118, 133, 135
iberica 5, 8, 11, 65, 70, 118, 135
lecerfi 5, 11, 65, 67, 70, 118, 133, 135
miki 12
nigripennis 5, 11-12, 70, 118, 133
pallida 4-5, 9-10, 12, 65, 67, 70, 76, 84, 86, 91-93, 95-98, 100, 105-106, 108-109, 111, 113, 118, 120-123, 130-131, 133, 135, 137, 143, 147
pilosula 4, 6, 10, 12, 64, 70-71, 133
rebecca 4, 6, 12, 64, 70-71
sp. from Spain 80, 84
Leptynoma 1-4, 10, 12-14, 25, 52, 64, 71, 78, 86, 90, 129-134, 136-137, 139-141, 145, 150
appendiculata 2, 12, 14, 64, 71, 130
hessei 12-15, 64, 72, 117, 130-131
maculata 12, 15-16, 64, 71, 130
namaquaensis 1-2, 12-15, 64, 72, 130-131, 141, 145
phantasma 12, 15-16, 71
sericea 2, 12-14, 16, 71, 76-77, 79-88, 90, 102, 105, 111-113, 115-117, 121, 127, 131, 136
Perianthomyia 3, 12-13, 15-16, 64, 71, 132-134, 145
Pheneus 16, 136
Protovermileo 1, 69, 71
electricus 69-70
Psammorycter 16
Vermileo 2-4, 16-18, 20-21, 23-24, 71-72, 78, 84, 88-92, 96, 99, 101, 103, 114, 116-117, 126, 129-131, 136-137, 140-143, 145-146, 148-150
ater 20-21, 22, 72, 118, 132
balearicus 24
comstocki 1-4, 16-19, 21, 62, 66, 72, 76-80, 84-86, 88, 90-91, 93-94, 98, 100-102, 104-109, 111-112, 116, 118-119, 121, 124, 130, 134, 136, 140-143
degeeri 16-17, 23, 145, 147
dowi 2, 17-18, 20-21, 22, 72, 102, 136
fascipennis 16, 18, 22, 24, 72, 136
nigriventris 20-22, 24, 72, 93, 95-96, 98
niloticus 21-22, 23-24, 72, 132, 146
opacus 1-3, 16-18, 23, 72, 118, 130, 136, 140-141
sp. from Egypt = possibly *V. niloticus* 76-78, 84, 87-88, 91-99, 101, 105, 109-112, 114-123, 146
tibialis 16, 18, 20-23, 72, 136
vermileo 2, 16-18, 20-23, 24, 63, 72, 76-81, 84-87, 91-93, 95-98, 100, 102, 104-106, 112-116, 118, 120-122, 136, 142, 146-147, 149
willetti 18, 24, 72, 136
Vermilyn 1-4, 12, 14, 25, 27, 52, 64, 72, 78, 90, 129-133, 136-137, 139, 140-141, 143, 145, 149
jasoni 4, 25, 27, 64, 72, 93-94, 129, 133, 143
vansoni 25-27, 72, 94, 129, 131, 133, 136, 143-144, 149
Vermiophis 1-2, 4, 26, 29-31, 68-69, 72-74, 90, 118, 128-131, 135-141, 151
ganquanensis 26, 29-32, 33, 35, 46, 68, 73, 76-78-79, 87, 89, 91, 124, 126, 128, 130, 138-140, 150
minshanensis 29-32, 35, 37, 46, 68, 73, 128, 130, 138-140
taihangensis 28-32, 37-38, 40-41, 49, 51, 68, 73, 128, 130, 138-140
taishanensis 28-32, 38, 40-42, 49, 74, 130, 138-140
tibetensis 28-32, 42-45, 69, 73, 118, 138-140
wudangensis 29-32, 46-48, 68, 73, 118, 129, 135, 138-140
yanshanensis 28-32, 38, 40-41, 49-51, 68, 74, 96, 129-130, 138-140
Vermipardus 1-4, 12, 14, 25, 51-52, 54-58, 64, 68, 75, 78, 90, 93, 99-100, 102-103, 117, 119, 129-132, 133-137, 139-141, 143, 145, 149-150
barraclooughi 51-52, 54, 58
basuto 51-52, 55, 119
brevirostris 51-52, 55, 75, 100, 118-119, 134, 143
brincki 51-52, 55, 58, 75, 119
intermedius 1-2, 19, 51, 53, 56, 62, 64, 66, 68, 75, 117-119, 131, 134
londti 51, 53, 56, 119
munroi 51, 53, 57, 119
promontorii 51, 53, 57, 64, 68, 76, 81, 86, 103-104, 118-119
sathon 51, 53, 57, 118-119
sylphe 51, 53-54, 56, 99-100, 102-103, 119
tresignatus 51, 53-54, 58
univittatus 51, 53, 56, 68, 75, 102-103, 118-119
whiteheadi 51, 54, 58
Vermitigris 2, 4, 54, 60-61, 69, 71, 74-75, 78, 90, 129-131, 136-137, 140-141, 146, 151

<i>fairchildi</i>	54, 60-61, 69, 75-76, 78-79, 84, 86-87, 89-91, 102, 114	<i>sinensis</i>	2, 59-60, 63, 65, 69, 74-76, 78, 84, 87, 90-91, 129
<i>infasciatus</i>	60, 75, 118	sp. from Sarawak	60, 115
<i>orientalis</i>	60-61, 62, 65, 75-76, 78-79, 84, 89-91	sp. from Tibet	67, 69, 74-75

GENERAL INDEX

adult phenology	117	Vermileonidae	80
antenna	64	larval spiracles	84
Palaeartic <i>Lampromyia</i>	65	larval stage, duration	120
breeding site	99	leaping	109
cannibalism	105	life cycle	117
copulation	94	longevity	118
diagnosis, family	2	male genitalia	19
diapause	121	<i>Lampromyia</i>	6, 7, 8, 9
dispersal in field	98	<i>Leptynoma</i>	14, 15
distribution	69	<i>Vermileo</i>	20, 21, 24
egg	91	<i>Vermilynx</i>	27
incubation period	119	<i>Vermiophis</i>	33, 37, 40, 42, 45, 48, 50
laying	96	<i>Vermipardus</i>	55, 56, 57, 58
number	98	metamorphosis	114
emergence	116	moulting	114
feeding habit	92	moults, number	121
female terminalia	66	mouthparts	62, 63
of <i>Vermiophis</i>	68	oviposition	95
first instar larva	109	site in the field	95
fossil	69	pit, shape	112
genral biology	92	postures and movements	105
hatching	114	prepupation and prepupal period	115
key, genera based on external characters	3	prey	103
genera based on male genitalia	4	capture	106
New World species of <i>Vermileo</i>	18	pseudopod on abdominal segment 1	79
Palaeartic species of <i>Vermileo</i> based on male genitalia	20	pseudotracheae	64
<i>Vermiophis</i> species based on aedeagus and ejaculatory apodeme	31	pupa	86
<i>Vermiophis</i> species based on antenna and coloration of body	29	Vermileonidae	87
<i>Vermiophis</i> species based on coloration of abdomen and legs	29	pupal spiracles	90
<i>Vermiophis</i> species based on fused gonocoxites and gonostylus	30	pupal stage, duration	122
<i>Vermiophis</i> species based on male genitalia	30	pupation	115
<i>Vermiophis</i> species based on tergum 9	31	sex ratio	92
<i>Vermitigris</i> species based on external character	60	sleeping posture	98
labellum	64	spermatheca	
larva	75, 99	<i>Lampromyia</i>	67
spiracles	85	<i>Vermipardus</i>	68
Vermileonidae	77	swarming	94
larval head		trap	111
skeleton	84	watch and wait attitude	106
parts of <i>Chrysopilus auratus</i>	82	wing venation in <i>Leptynoma</i>	13
parts of <i>Rhagio scolopaceus</i>	83		

日本熱帯生態学会 The Japan Society of Tropical Ecology

日本熱帯生態学会は熱帯地域の生態学的研究の進歩と研究成果の交流と普及に努めます。熱帯地域の生態学的研究といっても、熱帯生態学という確立した学問分野があるわけではありません。それぞれの研究者がそれぞれの研究方法によって熱帯地域の理解を深めていけばよいのですが、熱帯地域の自然に対する開発のすすみかたを考えると、熱帯地域の自然と人間生活をどのようにして持続的なものとして維持し発展させるかという視点が重要になってくることはさけられません。生物過程が主体となった自然生態系に注目するだけでなく、自然を組織している社会的なしくみにも目をつぶってはならないでしょう。このような意味でこれからの熱帯研究はより広い視野をもとめられるにちがいありません。幅の広ささまざまな研究の成果の相互交流が重要です。さらに研究成果は学界が独占すべきものではありません。広く社会に還元し、熱帯地域に生じている社会と自然をめぐる諸問題が正しく理解され、解決されるよう努力する必要があります。

TROPICS に投稿される方へのお願い

日本熱帯生態学会で編集をしているTROPICS は、国際的な学術雑誌として刊行されていますが、日本語による論文・総説も掲載します。現在、雑誌の編集・印刷は編集委員会事務局でマック DTP システムを使って編集・組版をして、印刷原版を打ち出し、それを印刷所でオフセット印刷しております。そのような事情がありますので投稿される方は以下の点に留意して原稿を作成・送付してください。

- 1 投稿原稿は手書きでもワープロを使用した打ち出しでも受付ますが、NEC-98 や DOS-V 系のワープロソフトを使った場合は必ずテキスト文 (ASCII-file) に変換したファイルもつけてお送りください。またマッキントッシュを使って作成したファイルの場合もかならずテキスト文に変換したファイルをつけておいてください。ファイルは 3.5 インチフロッピーディスク (2DD, 2HD どちらでもかまいません)、又は MO ディスクに入れて、原稿と同時に送ってください。使用ソフトは特に限定はいたしません、必ずテキスト文 (ASCII-file) に変換をしたファイルを加えてお送りくださるようお願いいたします。また使用機種とソフトを記録しておいてください。
- 2 字体の指定は編集委員会で行ないます。イタリック、上付きや下付きの字体指定は必ず打ち出し原稿にしておいてください。またタブ指定は表作成には必要ですが、本文中では使わないでください。指定を解除する余計な手間がかかります。
- 3 英文原稿の場合は日本語の、日本語原稿の場合は英文の摘要 (abstract) を忘れずにつけてください。外国からの投稿の場合は日本語摘要を編集委員会が準備することがありますが、そのような特別な場合を除いては摘要がないと完全な投稿原稿とは認められません。
- 4 図や写真は、充分完成されたものを作成してください。ただし図中の文字、数字、記号などは鉛筆書きの指定でもかまいません。地図などは編集委員会でリライトをすることがしばしば起こっていますが、つまらない誤りが発生することがあります。また、コンピュータで作図された図は、そのファイルを送られるときは必ず Mac の PICT 形式のファイルに変換しておいてください。
- 5 図や表は出来るかぎり頁幅で収まるようにしてください。表の組み上げが編集事務では多大の時間と神経を使うところです。特に大きな表や特殊な表は、そのままオフセット印刷原版に使えるように仕上げてください。
- 6 投稿原稿には、著者名、所属と住所、表題、見出しの表題 (短縮した)、キーワード、本文・図・表などの原稿枚数、原稿作成に使用したコンピュータの機種名と使用ソフト名、連絡先、投稿日などを送付状として付けてください。なお、くわしいことは日本熱帯生態学会投稿要項を参照してください。

投稿原稿は各編集委員または編集委員会事務局 (〒890-0065 鹿児島市郡元一丁目 21-35 鹿児島大学理学部地球環境科学科 (内) 堀田満 Tel. 0992-85-8166; Fax 0992-85-8172, e-mail: hotta@sci.kagoshima-u.ac.jp) あてにご送付ください。現在、TROPICS は年 4 回の発行がなされています。招待論文や十分に完成された論文がフロッピー入稿された場合であれば、最も速く処理がされると初校の打ち出しまでに必要な原稿整理の日数は数日、印刷に必要な日数は約 1 週間です。審査や校正のやりとりで手間取ることはありますが、迅速に研究成果を発表することが出来るシステムになっています。またページ制限は約 20 頁としてありますが、数十頁を越えるような大部な報文は Monographs として印刷報告することも可能です。

日本熱帯生態学会の事務局住所は 558-0022 大阪市住吉区杉本 3-3-138 大阪市立大学理学部 (気付け) 日本熱帯生態学会事務局 (山倉拓夫) です。入会・退会、会費納入等の学会事務は事務局で行なっています。

Monograph Series No. 1

1999年 5 月

熱帯研究

日本熱帯生態学会

永富昭, 楊集昆, 楊定 中国と世界のアナアブ科 (双翅目) 1-154

編集・発行

日本熱帯生態学会編集委員会 堀田満 (編集委員長)

〒890-0065 鹿児島市郡元1丁目21-35

鹿児島大学理学部地球環境科学科 (気付)

Tel. 099-285-8166 Fax 099-285-8172

E-mail: hotta@sci.kagoshima-u.ac.jp

印刷

斯文堂株式会社 鹿児島市南栄3丁目1 Tel. 099-268-8211

印刷・発行日 1999年 5 月25日印刷 1999年 5 月30日発行