STUDY ON THE MALE HORMONE (VI)
STUDY ON THE MALE HORMONE FROM BOAR
TESTES (III)
A NEW CRYSTAL MALE HORMONE*

A. Ogata and S. Hirano.

*From the Department of Pharmaceutical Chemistry, Medical Faculty,
Imperial University of Tokyo.*

(Received October 20, 1934)

We find that Butenandt (1) and Frattini & Maino (2) are the only workers who have written on the crystal male hormone up to the present moment, but Butenandt used the urine of human male as his material, so it is an open question whether or not the crystal obtained by him is identical with the hormone extracted from testis. Frattini & Maino got a crystal substance from horse testes but it appears to us that, for a pure hormone, it is too weak in its activity.

We began our study on the male hormone from testes in October, 1930, by using boar testes as our material, and, we have already reported upon a new method of extraction (3) and a method of purification (4). From January, 1934, we spent more time and labor in trying to separate crystal-fractions and oil-fractions from the testicular extract but so far we find the hormone is present only in the oil-fractions and the crystal-fractions were always inactive. Regardless of the fact that the estrus-producing hormone has already been taken out from the urine of pregnant women in crystalline form, and that a great progress has been made in the chemical studies of it, no one has yet succeeded in extracting it from the liquor folliculi in crystal state. Accordingly, we often thought that we, too, may not be able to get the crystal male hormone from testes, but we have finally

* We wish to express our sincere gratitude to the Imperial Academy for the support it has given to our study.

succeeded in obtaining an active crystal substance by purifying the oil-fraction after inactive crystals have been removed from it. The male hormone from testes is not an oily substance, can be confirmed by the fact that the activity of this crystal substance is far more potent than the mother liquor (oil-fraction) from which the active crystals are isolated.

TABLE 1.

Testes, being freed from epididymides and tunics, finely hashed.

	Extracted with ether after being stirred up with the same amount of 5 % NaOH	
Ether extract		Ether-insoluble substance (inactive)
Dissolved in acetone and cooled to -10°C		
Acetone extract		Cholesterol
Extracted with 70 % alcohol		
70 % alcoholic solution		Insoluble oily substance
Washed with hexane		
70 % alcohol extract (S 10)		Hexane-soluble substance (S 11)
Ether		
Ether-soluble substance (S 12)		Ether-insoluble crystals (S 13)
Crystals separate out; washed with acetic ether		
Acetic ether-soluble substance		Crystals (K 10) (Inactive)
Dissolved in a very small volume of absolute alcohol; hexane added		
Alcohol-hexane-insoluble oil		Alcohol-hexane-soluble crystals (K 11) (Inactive)
Extracted with 50 per cent alcohol		
50 % alcoholic solution		50 % alcohol-insoluble oil (K 7)
Allowed to evaporate naturally		
Crystal hormone (Fp. 128° - 130°)	Mother liquor (K 7 and K 10)	Saponified with 10 per cent alcoholic potassium hydroxide
	Unsaponifiable substance	Saponifiable substance
	Crystal hormone (Fp. 129 - 130°).	

Every fraction shown on the left hand side of the table, is active. Every fraction on the right hand side shows either inactive or slightly active.

TABLE 2.

Pre- paration	Fusing point	Daily dose of injection, mg. (injected once daily for two days)	Average percentage increase* in dimensions of part B of comb.					
			After completion of injections					
			First day	Second day	Third day	Fourth day	Fifth day	Sixth day
S 12	Heavy oily substance	5,337	20	22	18			
		8,539	26	30	24			
		12,809	45	55	67			
		14,944	42	56	50			
		18,149	47	65	68			
		21,348	49	73	83			
K 10	Fp 210°-217°	4,750	13	18	—	16		
K 11	Fp 93°-95°	2,000	4	4				
K 7	Heavy oily substance	6,809	28	—	35	29	—	33
K 6	Fp 130°-140°	5,500	15	—	84	109	—	124
K 8	Fp 138°-140°	3,500	34	77	99	—	—	127
K 9	Fp 128°-140°	0,850	1	13	17	25	36	24
K 12	Fp 128°-130°	0,867	—	14	14	23	—	—

*For dimensions of part B of comb, refer to Ogata, A. and Ito, Y.: J. Pharmaceut. Soc. (Japan) **53**, 39 (1933).

Only about 10 mg. of crude crystal hormone can be obtained from 80 kg. of testes. By repeatedly recrystallizing them from dilute alcohol, they are reduced to 2 mg. This crystal hormone has a melting point of 129°-130°, and is insoluble in dilute acid and alkali but soluble in ethanol, methanol, acetic ether, ethyl ether, benzene, chloroform, and acetone. Its activity is not destroyed by being boiled with alcoholic potassium hydroxide even for a long time, and the hormone does not contain any nitrogen. The melting point of the crystal hormone which Butenandt obtained from the male urine is 178°, it is therefore obvious that his crystal hormone is different

from that of ours. Figs. 2 and 3 are microscopic photographs of our crystals.

Table 1 shows the procedure of preparing the crystal hormone and table 2 the result of our comb test of each fraction, which is indicated in table 1. We have already reported up to S 10-fraction in our previous reports (3, 4), to which refer.

Explanation of table 2.

K6, K8, and K9 are the crude crystals of hormone which we obtained from different stocks and only purified by washing them with dilute alcohol. Their melting points vary more or less according to the stocks K6, K8, and K9. K12 is the pure crystals which were obtained by recrystallizing the crude crystals. In the case of K6 or K8, it shows the rate of increase* more than 120 % even on the sixth day after the completion of injections, and moreover, its reaction is still increasing. This may be considered as the result of too large a dose has been injected, but even when a far less dose is injected as in the case of K9 or K12, it shows the highest reaction on the fourth and fifth day after the completion of injections. The extracts we obtained so far, for example, S12 given in table 2, when large doses are injected, they naturally exhibit more strong reaction on the third day than on the second day after they are injected, but if proper doses are injected, they show the highest reaction on the second day after injections and the reaction decreases on the third day. If it exhibits the rate of increase of more than 20 % on the second day, it is generally observed that it has already exhibited the rate of increase close on to 20 % on the first day after injections. But in the case of K9, while it exhibited the rate of increase of 25% on the fourth day after injections, showed only 1% on the first day. In the case of K12, we did not measure the result of the first day after injections,

*The rate of increase in our comb test is the result of the measurement of the dimensions of part B of comb. The rate of increase in the dimensions of part B of comb does not differ substantially from the rate of increase in total dimensions of the silhouette of comb by planimetric measurement. For dimensions of part B of comb, refer to the foot-note of table 2.

as we judged from the result of K9 that it would not be large, but after comparing a subsequent rate of increase with that of K9, we thought that it will be safe to consider that it is almost identical with the result of the first day of K9. That is to say, when our crystal hormone is injected, it will take a little more time in exhibiting its reaction than when impurer extracts are injected, but the reaction is far more potent and lasts longer.

In testing the increase in weight of seminal vesicles of castrated rats with our purest crystal male hormone (Fp. 128-130°), we find that by injecting 330 γ of it to each rat daily for three days, *viz.* 990 γ , it exhibited an average rate of increase of 135 % when compared with the average weight of seminal vesicles of uninjected castrated rats. Fig. 1 is a photograph showing the result of this seminal vesicle test.

Experiment

6.12 litres of 5 per cent aqueous sodium hydroxide are added to the boar testes (6.12 Kg.), after being freed from epididymides and tunics finely hashed and stirred up for several hours. Then the mixture is put into a continuous extractor and extracted with ethyl ether. The ether is distilled from the hormone ether solution and the residue is dried, 60 g. of yellow creamy ether extract are obtained. 1100 cc. of acetone (Generally, 2~4cc. of acetone are employed for *ag.* of ether extract.) are added to 300 g. of the ether extract and shaken at room temperature and placed in a refrigerator of -10°C. for 12 to 24 hours, the cholesterol separates out. This is filtered by suction and the residue is again treated with equal volume of acetone. Both filtrates are mixed, acetone is completely evaporated and the residue cooled off again to -10°C., then cholesterol separates out, and this is centrifuged, 180 g. of red transparent adhesive oil (acetone extract) are obtained.

250g. of the acetone extract (Testes 42 Kg.) are extracted twice with 500 cc. of 70 per cent alcohol each time for two hours on boiling water bath.

The 70 per cent alcohol solutions of both times are mixed and washed with hexane until the hexane becomes colorless. The 70 per cent alcohol solution is next distilled under diminished pressure, ca. 1 g. of 70 per cent alcohol extract is obtained.

About 10 cc. of ether are added to ca. 1 g. of 70 per cent alcohol extract and shaken now and then and set aside for some time, the ether insoluble crystals (S 13 given in table 1.) are filtered. Occasionally, S 13 does not become crystalline powder, adhering to the walls of flask, so only the ether solution is transferred to another flask and ether is distilled; the residue remaining in the bottom of the flask is dried *in vacuo* and allowed to stand for a few days at room temperature, crystals separate out gradually from what was at first a red transparent adhesive oily substance and the mass becomes intransparent. Then the oily substance is dissolved quickly in a small volume of acetic ether and filtered by suction. (The crystals are inactive, K10) The acetic ether is distilled from the filtrate, a dark red transparent oily substance is obtained. This is dissolved in the least possible volume of absolute alcohol and hexane of 50 times by volume of absolute alcohol aforementioned, is poured in as it is stirred up, an oily substance separates out. After allowing it to stand overnight, the alcohol-hexane layer which has become transparent is separated from the oily substance that has settled in the bottom of the flask. This oily substance is boiled with ca. 10 cc. of 50 per cent alcohol, the alcoholic layer is separated from insoluble oily substance (K7) after cooled. The alcoholic layer is allowed to evaporate naturally for a few days, the hormone separates out in crystals for the first time. They are filtered by suction and when they are washed with a small volume of 50 per cent alcohol, 5mg. (Testes 42 Kg.) of fine crystalline powder are obtained. When this crystalline powder is recrystallized several times from 50 per cent alcohol, 1 mg. of crystals which has a melting point of 128°-130° is obtained.

Further, after saponifying the 50 per cent alcohol soluble substance

aforementioned with 10 per cent alcoholic potassium hydroxide for five hours, unsaponifiable substance is extracted with ether. This ether soluble substance is saponified again with 10 per cent alcoholic potassium hydroxide for five hours and unsaponifiable substance is extracted with ether. The ether is distilled from the ether solution and the unsaponifiable substance obtained at this stage, is dissolved again in a small volume of ether and the ether solution is allowed to evaporate naturally. When nearly all the ether is evaporated, crystals separate out from oily substance. This oily substance is dissolved in a small volume of ether and filtered by suction, the residual crystals on the filter are repeatedly recrystallized from dilute alcohol. Thus purified crystals have melting point of 129° – 130° , which differs no way from that of unsaponified crystal hormone noted above and physiological actions of both crystals are also identical.

Seminal vesicle test of our crystal male hormone.

The rats used in this test have been castrated at the same time when they were thirty days old, and seven months elapsed after castration. We injected to each of two castrated rats 330 γ of the crystals, dissolved in dilute alcohol, once daily for three consecutive days. The following is the result of measurements of weight of seminal vesicles taken on the fourth day :

Weight of seminal vesicles. mg.		The rate of increase in weight of seminal vesicles
Uninjected castrated rats	Injected castrated rats	
5.9 } 7.6 } 6.8	17.5 } 14.5 } 16	135 %

Discussion

The crystal male hormone which we obtained is more potent in its activity than any of the testicular extracts that has hitherto been reported.

Recently, Ruzicka et al (5) synthesized a substance identical with crystal

male hormone, so-called the "Androsteron", which Butenandt obtained from human male urine, and reported its physiological activities namely, when 200 γ of "Androsteron", dissolved in oil, are injected to each capon once daily for two days, the rate of increase in comb-growth is 20 %. But when 470 γ of our crystal male hormone, dissolved in dilute alcohol, are injected to each capon once daily for two days, they ought to show the rate of increase of 20 %, so we believe there is no substantial difference in the reactions between the "Androsteron" and our crystal hormone, when the difference in solvents of injection is taken into consideration. This fact has already been experienced by many workers when the estrus-producing hormone was tested. According to the experiment of Ruzicka et al the activity of the "Androsteron" shows the highest reaction on the second day after injections and the reaction is regressing on the third day, while, on the contrary, our crystal male hormone gradually becomes active from the second day after injections and exhibits the highest reaction on the fifth day, in other words, the reaction of our crystal male hormone lasts much longer (refer to K9 in table 2). Ruzicka et al, in their seminal vesicle test of "Androsteron", report that they injected 30 γ to each rat once daily for fourteen days, viz 420 γ in fourteen days, seminal vesicles of injected castrated rats which exhibited the maximum reaction weighed 30–40mg., while the average weight of seminal vesicles of uninjected castrated rats was 10 mg. But when 330 γ of our crystal male hormone were injected once daily for three days, viz. 990 γ in three days, the average rate of increase in weight of seminal vesicles was 135 %. We have already experienced in the seminal vesicle test that, the longer the period of injection is, the lesser the dose is enough to produce a larger rate of increase in weight of seminal vesicles than in the shorter period of injection.

For example, we show in table 3 the result of testing S10-Fraction.

The rats, each of which has been injected 9.945mg. in three days, exhibited only 248 % increase in the weight of seminal vesicles, while each of

which has been injected 6.489mg. in seven days, indicated an increase of

TABLE 3.

Dose mg.	Number of days of injection	Total amount of injected substance, mg.	Weight of seminal vesicles, mg.		Rate of increase in weight of seminal vesicles. %
			uninjected castrated rats	injected castrated rats	
0.927	7	6.489	5.2 5.2 5.0 } 5.13	32.5 25.7 35.2 26.0 } 29.85	482
3.315	3	9.945	4.7 4.0 } 4.4	17.5 15.5 13.0 } 15.3	248
1.658		4.974		9.8 9.0 12.5 } 10.4	136
3.315	2	6.630	3.8 4.9 5.1 } 4.6	9.7 8.4 8.2 } 8.8	91

(Every rat we used in this experiment is castrated thirty days after being bred and 27-37 days elapsed after castration and the uninjected castrated rats and the injected castrated rats used in each test are those which have been castrated at the same time.)

482 %. Again, the rats, each of which has been injected 6.630mg. in two days indicated an increase of only 91 %, while each of which has been injected 4.974mg. in three days, showed an increase of 136 %. Namely, the longer the period of injection, the less total amounts of substance are used than in the shorter period and moreover larger rates of increase are obtained than in the shorter period. In the example given in table 3 the differences in the periods of injection are one or four days, but it is very evident that the period of injection has exerted such a great influence on the reaction of the hormone, whilst the period of injection of "Androsteron" is eleven days longer than ours, so the activity of our crystal male hormone is probably more potent than the "Androsteron".

If the male hormone from urine shows the same physiological action as the male hormone from testes, then it makes us consider these two points, namely, whether the hormone from urine is identical with the hormone from testes, or whether a part of the molecule of the testicular hormone is decom-

posed and still it possesses the active atomic group or groups. But the hormone circulates all over the body with the blood and while developing its special physiological activity, it is subjected to such chemical change as oxydation or reduction, usually excreted with the urine, so we conclude that the crystal of Butenandt is probably an active product of decomposition of the male hormone from testes. Accordingly, we think it is quite natural that the crystal obtained by Butenandt does not correspond with our crystal. We believe this can be proved by the fact that notwithstanding the estrus-producing hormone has already been obtained a long ago from the urine of pregnant women in a crystalline form, but no crystal possessing the same chemical composition as the crystal obtained from the urine has yet been extracted from the ovary which is considered to be the organ producing the female hormone, although the active oily substance has been extracted from it.

Gallagher & Koch (6) note that the hormone extracted from the testes and the hormone obtained from human male urine possess different chemical properties. When the former is boiled with 5 per cent alcoholic alkali, its action against the comb is destroyed but the latter possesses resisting power against boiling alkali. But we have already noted under the section of experiment that the crystal male hormone which we obtained by boiling the 50 per cent alcohol-soluble substance with 10 per cent alcoholic potassium hydroxide for ten hours, has not only the same melting point and crystal form, but also the same activity against comb and seminal vesicles as the crystalline hormone extracted without taking this treatment, so it appears to us that Gallagher & Koch may have probably made mistake in their experiment.

Although we believe that the crystal obtained by us is pure, yet we will hereafter continue our work and find whether our crystal male hormone is wholly pure or mixture of crystal hormone and inactive principle by preparing various chemical derivatives.

Summary.

1. We have succeeded in extracting the crystal male hormone from boar testes and described the method of preparation. The physiological activities of the crystal hormone are more active than any of the testicular preparation that have been reported heretofore.
2. The activities of the crystal male hormone extracted from boar testes against comb and seminal vesicles are not destroyed by alkaline hydrolysis.
3. We have proved by fairly well-grounded experiments that the crystal male hormone from testes and the hormone from urine are not chemically identical.

Fig. 1

A photograph showing the reaction of our crystal male hormone exhibited in the seminal vesicles of castrated rats.

Left figure: Seminal vesicles of two rats; 0.330 mg. are injected once a day for 3 days.

Right figure: Seminal vesicles of two uninjected rats.

Fig. 2

Fig. 3

Fig. 2 and 3 are photographs showing the crystals obtained from absolute alcohol solution by natural evaporation.

Fig. 2 Photographed by a penetrated light, ca. 400 $\times$

Fig. 3 Photographed by a reflected light, ca. 200 $\times$.

References.

- (1) Butenandt, A.: Zt. angew. Chem. **44**. 905 (1931).
- (2) Frattini, B. & Maino, M.M.: B. ges. Physiol. **60**. 308 (1930).
- (3) Ogata, A. and Hirano, S.: J. Pharmaceut. Soc. (Japan) **52**. 187 (1932).
- (4) Ogata, A. and Hirano, S.: J. Pharmaceut. Soc. (Japan) **53**. 153 (1933).
- (5) Ruzicka, L. et al.: Helv. **17**. 1395-1406 (1934).
- (6) Gallagher, T. F. and Koch, F. C.: J. Biol. Chem. **100**. xlviii (1933);
Endocri. **18**. 107. (1934).

Y. Kondo: Über den diagnostischen Wert des Blatt-Aschenbildes mit besonderer Rücksicht auf die Systematik der Rutaceae. (Résumé.)

Studien über die Erkennung der Drogen auf Grund des Aschenbildes. (V. Mitteilung.)

[Aus dem Institut für Warenkunde der Handelshochschule Nagoya.]

(Eingegangen am 7. November 1934.)

Um den diagnostischen Wert der Merkmale festzustellen, die vom Verfasser seit 1929 bei seinen Untersuchungen¹⁾ zur Erkennung der Drogenblätter und Krautdrogen auf Grund des Aschenbildes benutzt wurden, und um daneben die Ergebnisse der vorangegangenen Untersuchungen zusammenfassend darzustellen, wurde eine Reihe Untersuchungen ausgeführt. Da die Diagnostik der Drogen nichts anderes ist als die Erkennung der Pflanzen-

¹⁾ Ohara, K. u. Kondo, Y., Studien über die Erkennung der Drogen auf Grund des Aschenbildes. (I. Mitteil.) Aschenbilder der Drogenblätter in Pharmacopoea Japonica. IV. Diese Zeitschrift. 573, 164 (Vol. 49, Nov., 1929).

Ohara, K. u. Kondo, Y., dasselbe (II. Mitteil.) Über die Aschenbilder wichtiger Drogenblätter. Arch. d. Pharm., Jahrg. 1931, S. 292.

Kondo, Y., dasselbe (III. Mitteil.) Aschenbilder wichtiger, in den japanischen und österreichischen Pharmakopöen fehlender Drogenblätter. Diese Zeitschrift. 597, 124 (Vol. 51, Nov., 1931).

Kondo, Y., dasselbe (IV. Mitteil.) Blatt-Aschenbilder wichtiger Krautdrogen. Diese Zeitschrift. 615, 71 (Vol. 53, Mai, 1933).

Kondo, Y., Nachtrag zu Aschenbilder wichtiger, in den japanischen, deutschen und österreichischen Pharmakopöen fehlender Drogenblätter. Diese Zeitschrift.

男性ホルモンの研究 (第六報)

睾丸ホルモンの研究 (其三)

一新結晶男性ホルモン

緒 方 章

平 野 四 郎

(東京帝國大學醫學部藥學科)

(昭和 9 年 10 月 20 日受理)

現在までに結晶性男性ホルモンの就て記述しているのは A. Butenandt (Ztsch. angew. Chem. 44. 905. 1931) と B. Frattini u. M. M. Maino (B. ges. Physiol. 60. 308. 1930) のみである。然し Butenandt は男尿を原料としているから、彼の得た結晶が睾丸から産出されるホルモンと同一物であるか何うかは未決の問題である。又 Frattini u. Maino は馬睾丸を原料として結晶性物質を得たが純粹のホルモンとしては効力が餘り弱い様に思はれる。

余等は 1930 年 10 月より豚睾丸を原料として睾丸ホルモンの研究を始め一新抽出方法(本誌 52 1060. 1932) 及精製方法(本誌 53. 869. 1933) を報告した。其後余等は本年 1 月より更に多大の勞力を費して睾丸エキスを結晶性部分と油狀部分に分離することに勉めたが今度はホルモンは常に油狀部分にのみ存在して結晶部分は常に無効であつた。卵巢濾胞ホルモンは妊婦尿から既に早くから結晶として取出され其化學的研究も大に進歩しているに拘はらず卵巢濾胞からは未だ何人も結晶狀態に抽出していないから余等は睾丸ホルモンも結晶體として睾丸から得られないものかも知れぬと思つたこともある。然るに今回無効結晶を除去した油狀部分を更に精製して遂に効力の強い結晶を得ることに成功した。而して睾丸ホルモンが油狀物質でないことの證據は此結晶の効力が此者を去つた母液(油狀部分) よりも遙に強いことによつて確認出来る。

粗製ホルモン結晶は睾丸 80kg から僅に約 10mg を得るに過ぎず、之を數回水性アルコールから再結晶すると 2mg に減ずる。此結晶は Fp. 129-130° で稀酸アルカリに不溶, Methanol, Äthanol, Chloroform, Benzol, Essigester, Äther, Aceton に可溶である。10%酒精性カリと長時間煮沸しても効力は破壊されず窒

素を含有しない。Butenandt が男尿より得た結晶は $Fp. 178^{\circ}$ であるから余等の結晶と異物であることは明である。歐文の Fig 1 及 Fig 2 は余等の結晶の顯微鏡寫眞である。

第 1 表
睾丸白膜を除去磨潰したる睾丸

5% NaOH の同量と攪拌後エーテル抽出	
Ätherextrakt	残 渣
Aceton に溶し $-10^{\circ}C$ に冷却	
Acetonextrakt	類脂體
70% Alkohol 加熱抽出	
70% Alkohol 溶液	不溶油狀物
Hexan で洗滌	
70% Alkohol extrakt (S10)	Hexan 可溶物 (S11)
Aether	
Aether 可溶物 (S12)	Aether 不溶結晶 (S13)
數日後析出した結晶を Essigester で洗滌	
Essigester 可溶物	結 晶 (K10)
極少量の純 Alkohol に溶し 50 倍量の Hexan を注加	
不溶油狀物	Alkohol-Hexan 可溶結晶 (K11)
50% Alkohol 溶液	
自然蒸發(數日放置)	10% Alkohol 性 KOH で鹼化
ホルモン結晶 母液 (Fp 128-130°) (K7, K10)	冷後不溶の油 (K7)
	不鹼化物
	酸 Fraktion
	ホルモン結晶 (Fp 129-130°)

表中左側の各 Fraktion は有効、右側の各 Fraktion は無効或は微弱の效力を示すに過ぎぬ。

第 2 表

製品	熔 融 點	1 回注射量 mg (1 日 1 回 2 日間注射)	平均雞冠 B 面積増加率 %					
			注射終了 第 1 日	注射終了 第 2 日	注射終了 第 3 日	注射終了 第 4 日	注射終了 第 5 日	注射終了 第 6 日
S 12	油狀粘稠物	5.337	20	22	18			
		8.539	26	30	24			
		12.809	45	55	67			
		14.944	42	56	50			
		18.146	47	65	68			
		21.348	49	73	83			

K 10	210-217°	4.750	13	18		16		
K 11	93-95°	2.000	4	4				
K 7	油状粘稠物	6.809	28		35	29		33
K 6	130-140°	5.500	15		84	109		124
K 8	138-140°	3.500	34	77	99			127
K 9	128-140°	0.850	1	13	17	25	36	24
K 12	128-130°	0.867		14	14	23		

第 1 表はホルモン結晶の抽出方法を表示したもの、第 2 表は第 1 表に示す各 Fraktion の去勢雞雞冠試験の結果を示す。但し S13 までは既に前報に記載したから夫等を参照され度い。

第 2 表 の 説 明

K6, K8 及 K9 は異なる原料から得たホルモンの粗製結晶で原料の異なるにつれて幾分 Fp を異にするが夫等を合して數回再結晶を繰返して得たものが K12 である。K6 或は K8 の場合には注射終了後第 6 日に於ても 120% 以上の増加率*を示し而も効力が尙ほ上り坂である。此事は注射量が多過ぎた爲とも考へられるが夫よりも遙に少量を注射した K9 或は K12 の場合でも尙注射終了後第 4 日、第 5 日目に最高反應を示している。之までに余等が得たエキス例へば第 2 表に於ける S12 では多量を注射した場合には注射終了後第 2 日より第 3 日の方が効力が強く顯はれるのは勿論のことであるが注射量が適量であれば注射終了後第 2 日に最高反應を示し第 3 日には既に効力が下り坂となる。而して第 2 日に 20% 以上の効力を示す時には注射終了後第 1 日に於て既に 20% に近い増加率を與へるのが普通である。然るに K9 の時には注射終了後第 4 日に 25% を示すに拘はらず第 1 日には僅に 1% である。K12 の場合には K9 の結果から推察して第 1 日の結果は大きい増加率を示さぬだらうと思ひ測定しなかつたが其後の増加率を K9 と比較することによつて第 1 日の結果は K9 の第 1 日と略々同様と考へて間違はない。即ち余等のホルモン結晶を注射した場合には不純エキスを注射した時よりも効力の發顯は幾分遅れるが効力は遙に強く又長時間持続する。

余等の最も精製した結晶(Fp.128-130°)に就て 去勢ラツテの 精囊重量増加試験

* 余等の雞冠試験に於ける増加率は總て雞冠 B 面積の増加率である。雞冠 B 面積の増加率は雞冠陰影像全面積の Planimeter による測定結果の増加率と大差がなく只正確度に於て前者が優れている。

を行ふに, 1日 330r 宛 3 日間即ち 990r の注射で試験開始後第 4 日目に對照動物の平均精囊重量に比し 135% の平均重量増加率を示した. 歐文 Fig3 は此精囊試験の結果を寫眞で示す.

余等の得た男性ホルモン結晶は從來發表された睾丸ホルモンの何れよりも効力が強いが結晶尿ホルモンの効力との比較は未だ此者の効力に就て信頼し得る報告がなかつた爲め行ひ得なかつたが, 最近 L. Ruzicka et al (Helv. 17. 1395-1406. 1934) は Butenandt が男尿より得た男性ホルモン結晶 Androsteron と同一物を Dihydrocholesterin を酸化して人工的に製し比較的詳細に其生理的効力をも發表してゐる. 即ち Androsteron は 200r づつ 1 日 1 回 2 日間注射して雞冠面積増加率 20% を示すに對し余等の結晶は 470r づつ 1 日 1 回 2 日間注射すると増加率 20% を示す筈である. 故に Androsteron の場合には注射液は油溶液であり, 余等の結晶の場合は水性アルコール溶液であるから, 兩者の溶媒の相違を考慮に入れるならば兩者の効力には大した相違がないと信ずる. 此事實は既に女性ホルモンの試験に際して多數の研究者が経験してゐる處である. Androsteron の効力は注射終了後第 2 日に最高反應を示し第 3 日目には既に効力が減少しつつあるに反し余等の結晶は注射終了後第 2 日より次第に効力を發顯し第 5 日に最高反應を示してゐる. 則ち余等の結晶ホルモンの効力の方が遙かに長續きする (第 2 表 K9 参照). 尙 Ruzicka et al は Androsteron の精囊試験に於て 1 日 1 回 30r づつ 14 日間即ち 420r 注射して最大反應を示した精囊の重量は 30-40mg であつて, 對照去勢ラットの平均精囊重量は 10mg であつたと云つてゐる. 然るに余等は余等の結晶について, 1 日 1 回 330r づつ 3 日間即ち 990r の注射で 135% の精囊重量増加率を示した. 精囊試験に於ては注射期間が長い程注射量は少量でも却つて大きい重量増加率を示すものであることは既に余等の経験する所である. 今余等の製品 S10 (第 1 表参照) について注射期間と効力の關係を表示すると第 3 表の如くなる. 即ち全量 9.945mg を 3 日間に注射したものは精囊重量増加率僅に 248% なるに反し全量 6.489mg を 7 日間に注射したものは増加率 482% を示した. 又全量 6.630mg を 2 日間に注射したものは 91% なるに反し 4.974mg を 3 日間に注射したものは 136% を示した. 余等の今茲に示した例は注射期間の相違が僅かに 1 日或は 4 日の場合であるが尙ほ且つ注射期間が長い程注射全量は少量でも却つて大きい増加率を示すことが明瞭に認められる. 却説,

此例より判断する時 Ruzicka et al の注射期間は 14 日であり余等のは 3 日であつて 11 日の相違があるから余等の男性ホルモン結晶の去勢ラツテ精嚢に對する効力は Androsteron よりも寧ろ強力であると考へるのが妥當であらう。

第 3 表

1 回注射量 (mg)	注射日数 (1日1回注射)	注射全量 (mg)	精嚢重量 (mg)		平均精嚢重量 増加率 %
			對照動物	注射動物	
0.927	7	6.489	5.2 5.2 5.2 5.0	5.13 25.7 35.2 26.0 32.5	482
3.315	3	9.945	4.7 4.0	4.4 17.5 15.5 13.0	248
1.658	3	4.974	4.7 4.0	4.4 9.8 9.0 12.5	136
3.315	2	6.630	3.8 4.9 5.1	4.6 9.7 8.4 8.2	91

尿製品が睾丸ホルモンと同じ生理作用を顯す時は、此製品は睾丸ホルモン共者であるか或は睾丸ホルモン分子が一部分分解し、然も尙ほ未だ有効原子團を保有してゐるものであるか此の二つのことが考へられる。然しホルモンは血液と共に身體を循環して特殊の生理作用を發顯する間に多くは酸化或は還元の如き化學的變化を受けて尿中に排泄される場合が多いから余等は Butenandt の得た結晶従つて Ruzicka et al の製した結晶 Androsteron も恐らくは睾丸ホルモンの有効分解産物であらうと理解する。従つて Androsteron と余等の得た結晶とが化學的に一致しないのは寧ろ當然の結果ではなからうかと考へる。此事實は妊娠尿からは卵巢濾胞ホルモンと同じ生理作用を顯すホルモンを既に早く結晶として得られたに拘はらず其產出器官と考へられてゐる卵巢よりは未だ之と同一化學組成を有する結晶の抽出されてゐないことによつて實證されると思ふ。

Gallagher u. Koch (J. biol. Chem. 100. XLVIII. 1933; Endocr. 18. 107. 1934) は睾丸より得べきホルモンと男尿より得べきホルモンとの化學的性質の相異として前者は 5% 酒精性アルカリと煮沸することによつて鶏冠に對する效力を消失するが後者はアルカリによつて効力が破壊されないと云つてゐる。然るに余等は實驗之部に示した如く 50% Alkohol に可溶の Fraktion を 10% 酒精性 KOH

と 10 時間煮沸鹼化したものから加水分解をせずに單に抽出した結晶と結晶形, 熔融點を同じうするのみならず雞冠及精囊に對する效力も何等變化のないホルモン結晶を得たから Gallagher u. Koch の説は間違つてゐると思ふ。

余等は余等の得た結晶がホルモンなりと信ずるものであるが今後引續き研究を行ひ諸種の化學的誘導體を製して此結晶が單一性のものなるか又は混合結晶であるかを追究せんとするものである。

次に余等の前報に發表した精製法に改良を加へた點に就て述べる。辜丸ホルモンの研究其二(前掲)の 70% C_2H_5OH エキス(S10)は Aceton エキスを先づ Methanol で抽出し其可溶成分を 70% C_2H_5OH で煮沸し C_2H_5OH 層を Hexan で完全に洗滌して製したが其後の實驗により Aceton エキスを直接 70% C_2H_5OH で抽出しても效力が前報のものに劣らぬのみならず此 S10-Fraktion を Äther で所理すれば前報通り Äther 不溶の無效結晶(S13)及 Äther 可溶の有効 Fraktion(S12)を得る。故に爾後 CH_3OH 所理を省くこととした。尚ほ 70% C_2H_5OH 層を洗滌するのは必ずしも Hexan に限らず石油エーテル(Kp.40-70°)で充分であることを知つた。又 Aceton エキスを 70% C_2H_5OH で何回抽出すれば完全にホルモンを抽出し得べきやを検する爲 Aceton エキス 200-300g に付き毎回 70% C_2H_5OH 500cc 宛を用ひて 4 回抽出を試みたが最初の 2 回で大部分抽出され次の 2 回で尙幾分の有效物質が抽出される。然し後 2 回には不純物も多く溶出し來り従つて製品の效力は半減することを知つた。故に後の 2 回の抽出はホルモンを回収するには良いが純粹抽出の目的には最初の 2 回だけに止める方が得策である。若しそれ以上の抽出を行つても第 5 回目の分には極めて少量のホルモンが溶けて來るに過ぎぬ。

今回余等はホルモン結晶以外に K10 及 K11 なる結晶を得た。此等の結晶は前報に記した結晶 S13 と共にホルモンに比して收得量が稍良好であるから、精製の上分析結果を發表し得るのは近き將來であらうと信ずる。

實 驗 之 部

結 晶 ホ ル モ ン の 製 法

S12-Fraktion ca. 1g (辜丸 42kg)を數日間室溫に放置すると最初赤色透明粘稠の油狀物質であつたものが次第に結晶を析出して不透明になる。此時に少量の醋酸エステルで油狀物質のみを迅速に溶して吸引濾過する(結晶 **K10** は幾分ホル

モンを附着し従つて生理試験結果は幾分有効であるが此附着したホルモンは此結晶を純 C_2H_5OH より再結晶すると母液より回収することが出来る)。次に濾液より醋酸エステルを去ると暗赤色透明の油狀物が残留するから之を出来るだけ少量の純 C_2H_5OH に溶し 50 倍量の Hexan を攪拌しつつ注加すると不溶油狀物が析出する。1 夜放置後透明になつた Alkohol-hexan 層を器底に沈んだ油狀物質から分離する (Alkohol-hexan 層を濃縮すると無効結晶 **K11** を得る)。此油狀物質を 50% C_2H_5OH ca 10cc で煮沸し冷後 Alkohol 層を不溶の油狀物 (**K7**) と分離し Alkohol 層を數日放置して自然に蒸發させるとホルモンは始めて結晶となつて析出する。之を吸引濾過し少量の 50% C_2H_5OH で洗滌するとさらさらの粉末 (**K6, K8** 及 **K9**) になる。收量 5mg 之を數回稀 Alkohol から再結晶すると Fp. 128-130° を示す結晶 **K12** (收量 1mg) を得る。此際母液よりは前記 **K7** 及 **K10** を得る。

尙ほ前記 50% C_2H_5OH 溶液からホルモンを自然蒸發によつて析出させる代りに此 50% C_2H_5OH 可溶物 (睾丸 120kg) を 10% Alkohol 性 KOH 30cc で 5 時間鹼化後不鹼化物を Äther 抽出しそれを更に同様に鹼化し、再び不鹼化物を Äther で抽出し Äther 溶液から Äther を溜去した後殘渣を少量の Äther に溶し自然に Äther を蒸發させると殆んど全部の Äther が蒸發し去つた時に油狀物中に結晶を生じる。これを少量の Äther を加へて吸引濾過し稀 Alkohol より再結晶すると Fp. 129-130° を示し **K12** と生理的效力も變らないものが得られる。粗結晶の收量約 12mg であつたから **K6, K8** 及 **K9** の時と收量に大差なくアルカリ煮沸によつてホルモンが分解されないことがわかる。

前記の無効結晶 **K10** は數回純 C_2H_5OH より再結晶を反覆すると Fp. 216-223° の無色長板狀の結晶となり **K11** は純 C_2H_5OH より再結晶すると Fp. 93-95° の無色針狀結晶となる。

K6, K8, 及 **K9** はホルモンの粗製結晶で原料を異にするにつれ幾分 Fp を異にする即ち Fp は夫々 130-140°, 138-140° 及 128-140° で之等を數回再結晶したものは Fp=128-130° になる。而して此母液より上記 **K10** に相當する Fp 216-223° の結晶を回収する。

動物試験

ホルモン結晶 **K6, K8, K9, K12** 及油狀 Fraktion **K7** 及無効結晶と見做すべき

K10 及 K11 の雞冠試験を次に示す. 注射は何れも 1 日 1 回 2 日間, 注射液は稀 C_2H_5OH 溶液.

製 品	雞番號	1回注射量 mg	雞 冠 B 面 積 増 加 qmm 及		
			第 1 日	第 2 日	第 3 日
K6 (Fp130-140°)	Nr. 51	5.500	363-317=46 (15)	—	582-317=265(84)
K8 (Fp138-140°)	Nr. 484	3.500	489-364=125(34)	646-364=282(77)	726-364=362(99)
K9 (Fp128-140°)	Nr. 51	0.850	290-287=3 (1)	324-287=37 (13)	337-287= 50(17)
K12 (Fp128-130°)	Nr. 51	0.867	—	292-257=35 (14)	294-257= 37(14)
K7	Nr. 56	6.809	558-441=117(27)	—	620-441=179(41)
	Nr. 133		368-286=82 (29)	—	369-286= 83(29)
K10 (Fp210-217°)	Nr. 33	4.750	189-167=22 (13)	197-167=30 (18)	—
K11 (Fp93-95°)	Nr. 51	2.000	281-271=10 (4)	281-271=10 (4)	

製 品	雞番號	1回注射量 mg	増 加 率(%) 注射終了後		
			第 4 日	第 5 日	第 6 日
K6 (Fp130-140°)	Nr. 51	5.500	664-317=347(109)	—	709-317=392(124)
K8 (Fp138-140°)	Nr. 484	3.500	—	—	825-364=461(127)
K9 (Fp128-140°)	Nr. 51	0.850	360-287=73 (25)	389-287=102 (36)	355-287=68 (24)
K12 (Fp128-130°)	Nr. 51	0.867	316-257=59 (23)	—	—
K7	Nr. 56	6.804	604-441=163 (37)	664-441=223 (51)	646-441=205 (46)
	Nr. 133		346-286=60 (21)	327-286=41 (14)	312-286=26 (9)
K10 (Fp210-217°)	Nr. 33	4.750	193-167=26 (16)		
K11 (Fp93-95°)	Nr. 51	2.000			

K10 の精囊試験 生後 30 日に去勢し去勢後 4 ヶ月半経過した雄ラツテに 1 日 1 回 2.666mg (0.5cc 稀 Alkohol 溶液)宛 3 日間繼續して皮下注射し第 4 日目に精囊重量を検すると注射動物 2 匹の精囊は夫々 8.0mg 及 9.2mg で對照動物 2 匹の精囊は 6.7mg 及 7.0mg で極めて少量のホルモンが附着する程度に過ぎぬことがわかる.

ホルモン結晶 K12 の精囊試験 生後 30 日に去勢し去勢後 7 ヶ月経過した雄ラツテに 1 日 1 回 0.330mg (0.1cc 稀 Alkohol 溶液)宛 3 日間注射し第 4 日目に

精囊重量を検する.

注射動物精囊重量(2匹)	$\left. \begin{array}{l} 17.5 \\ 14.5 \end{array} \right\} 16\text{mg}$	∴ 増加率 135%
対照動物精囊重量(2匹)	$\left. \begin{array}{l} 5.9 \\ 7.6 \end{array} \right\} 6.8\text{mg}$	

70% C₂H₅OH エキス(S.10)製造の變法

Aceton エキス 300g に 70% C₂H₅OH 500cc を加へ還流冷却器下に 2 時間煮沸し 1.5 時間水冷後アルコール層(幾分濁濁)を分液漏斗中に傾瀉移入し残留する油層には更に 70% C₂H₅OH 500cc を加へて先と同様に煮沸後新しい分液漏斗中にコルベン内容を全部移入し 1 夜放置後完全に分離析出した油層を分離しアルコール層(Cholesterin の結晶析出する場合あり)は前回分離のアルコール層に合し Hexan(石油エーテル Kp.40-70° で代用する)が無色になるまで完全に Hexan で洗滌し最後にアルコール層を減壓蒸溜乾燥する. 收得量 1g 内外.

本品は S.10(本誌 53, 883, 1933 参照)相當物で Äther で所理すると S.12 及 S.13 相當物(本誌 53, 885 参照)を得.

S.10 の雞冠試験 本品 370.7mg を水性 C₂H₅OH 10cc 溶液とし 1 日 1 回 2 日間注射したとき注射終了翌日の雞冠 B 面積増加率は次の如くである.

雞番號	1回注射量 (mg)	注射前B面積 (qmm)	注射後B面積 (qmm)	B面積増加率 (%)	平均増加率 (%)
Nr.122	11.121	114	280	146※	58
Nr.443	„	134	199	49}	
Nr.56	„	264	441	67}	
Nr.443	5.561	201	231	15}	19
Nr.122	„	341	308	28}	
Nr.137	„	261	294	13}	

※此數を平均増加率計算に除外した理由は B 面積増加率最大差異 170% 以上を示したからである(本誌 54, 355 参照). 以下之に準じる.

又 2 回 70% C₂H₅OH で抽出した Aceton エキス 300g に就て更に 70% C₂H₅OH 500cc 宛で 2 回熱時抽出し Hexan で洗滌したものは 0.2004g あり. 其雞冠試験は

1回注射量 (mg)	雞番號	注射前B面積 (qmm)	注射後B面積 (qmm)	B面積増加率 (%)	平均
10.020	Nr.15	331	374	13	14
	Nr.51	280	317	13	
	Nr.133	281	324	15	
12.826	Nr.485	350	467	33	26
	Nr.122	256	301	18	

14.028	Nr.162	230	281	22}	25
	Nr.33	248	315	27}	
	Nr.177	234	249	6	

以上にて Aceton エキスの 70% C_2H_5OH 抽出は合計 4 回で充分なるを知る。5 回或は夫以上は無効物質の混入が多くなる。本報告に於ける S.10 或は S.12 は 70% C_2H_5OH 抽出 2 回分を意味する。

本研究には帝國學士院に學術研究費を補助され、又帝國社臓器薬研究所は多量の罂丸エキスを供給せられ又介補平澤金太郎君は原料製造に終始熱心に従事された。茲に記して厚く謝意を表す。

葉部灰像の鑑識學的價值竝にへんろうだ科

植物葉部灰像の系統學的研究

灰像による生薬鑑識の研究〔第五報〕

近 藤 良 男

(昭和 9 年 11 月 7 日受理)

(名古屋高等商業學校商品學教室)

Y. Kondo: Über den diagnostischen Wert des Blatt-Aschenbildes mit besonderer Rücksicht auf Systematik der Rutaceae.

Studien über die Erkennung der Drogen auf Grund des Aschenbildes.

(V. Mitteilung.)

目 次

1. 緒 論 2. 鑑識學的特徴と系統學的特徴 3. 葉部灰像に於ける特徴 (1) 蓆酸石灰結晶 (2) 結晶の分布型 4. 葉部灰像特徴の系統學的意義 5. へんろうだ科植物の葉部灰像 (1) 實驗材料 (2) 蓆酸石灰結晶 (3) 分布型	(4) へんろうだ科植物葉部灰像特徴一覽表 6. 葉部灰像に及ぼす内外の生理的影響 (1) 産地の差異による影響 (2) 葉の成, 若による影響 7. 考 察 (1) 系統學的灰像研究の既往の業績 (2) 結果の考察及結論 8. 摘 要