Jpn. J. Med. Sci. Biol., 49, 129 - 149, 1996.

**Review: SCHISTOSOMA MANSONI: THE ULTRASTRUCTURE OF LAR-
VAL MORPHOGENESIS IN BIOMPHALARIA GLABRATA AND OF ASSO-
CIATED HOST-PARASITE INTERACTIONS**

S. Chia-Tung PAN*

*Department of Tropical Public Health, Harvard School of Public Health, 665 Hun-
tington Avenue, Boston, MA 02115, USA*

(Received July 15, 1996. Accepted September 13, 1996)

CONTENTS

1.	Introduction.....	130
2.	Materials and Methods.....	131
3.	Results.....	131
	Postmiracidium.....	131
	Mother sporocyst.....	136
	Microvilli of mother sporocyst.....	139
	Germ cells.....	140
	Excretory system.....	142
	Musculature.....	142
	Interstitial cells.....	142
	Nervous system.....	142
	Miscellaneous.....	142
4.	Comments.....	144
5.	Summary.....	146
6.	References.....	147

*Present address: 91-219 Waopio Place, Kapolei, Hawaii 96707, USA

INTRODUCTION

It is estimated that schistosomiasis affects over 200 million people primarily in developing countries and is spreading rapidly as irrigation schemes are expanded for agricultural needs. Yet our knowledge on the basic biology of the schistosome is still far from adequate (1). A rational approach to the effective control of human schistosomiasis may depend on such knowledge.

The miracidium is the first infective stage in the complex life-cycle of the parasite, and the future of the entire life-cycle depends on the capacity of the larval stage to locate and enter a suitable snail host. Once the miracidium enters a snail host, it must go through a radical morphological and physiological reorganization to develop and multiply asexually in the snail. In turn the snail host reacts to the presence of the foreign object. Studies on the host-parasite interactions with the light microscope have not revealed details of the early morphogenesis of the miracidium after invasion of a suitable or resistant snail (2-7). It has been shown that some miracidia may degenerate in 24 hr after penetration and that on the average only 3-6 mature mother sporocysts develop per snail to produce daughter sporocysts, even the snails may have been invaded by as many as 86 miracidia (5).

We completed in 1980 a monographic study of the fine structure of the miracidium of *Schistosoma mansoni* (8). This provides the basis for the present study of the morphogenesis of the miracidium in susceptible snails and on the fine structure of the host-parasite interactions. We observed that certain structures of the miracidium are not only responsible for the rapid formation of the new tegument of the mother sporocyst, but also may prevent close contact of snail amoebocytes with the postmiracidium stage (8). Such defense mechanisms by the parasite appear to be essential for survival during the first few days of parasitic life. On the other hand, the defense mechanisms of the parasite may not be functional in resistant snails as evidenced by several in vitro studies (9-11). While the mechanisms affecting compatibility between parasite and snail host have begun to attract researchers, detailed knowledge on host-parasite relations, both in compatible and incompatible hosts, is still inadequate, especially at the subcellular level. Such studies may eventually lead to a fundamental understanding of the biology of host (snail)-parasite (schistosome) interactions.

In this review, we have described the fine structure of morphogenesis of the miracidium in the susceptible snail host *B. glabrata* and the host-parasite interactions between them.

MATERIALS AND METHODS

The strains of *B. glabrata* and of *S. mansoni* and their maintenance have been described (5). Snails (5-7 mm in diameter) were mass-exposed to numerous, freshly hatched miracidia for one hour in small finger bowls (10 × 4.5 cm). Exposed snails were removed from maintenance aquaria at intervals for processing of their tissues for light and electron microscopy as before (8). It should be noted that among several buffers tried for dilution of glutaraldehyde and osmium tetroxide, Chernin's handling solution (12) appeared to be the most suitable for snail and parasite tissues. Extensive light microscopy of thick sections (0.25-0.5 μ) stained with toluidine blue O was essential to locate the minute parasites in snail tissues (tentacles, head-foot, mantle collar, etc.), prior to electron microscopy.

RESULTS

The most important miracidial structures involved in the miracidium-mother sporocyst transformation during the first 24 hr of parasitic life are the syncytial epidermal ridges with accompanying cytons, 21 ciliated epidermal plates that are attached to the ridges by septate desmosomes and the terebratorium. The terebratorium appears to be a modified epidermal ridge containing at least 12 peripheral sensory organelles and openings of three secretory ducts of glandular cells (8).

The ages of the organisms are expressed from the end of exposure of snails to miracidia.

Organisms within 5 hr after exposure: Postmiracidium

The miracidium of *S. mansoni* penetrates its snail host without discarding the ciliary epidermal plates, although the larva loses its swimming capacity immediately after penetration (2). Figure one illustrates the anterior portion of a sagittal section of a miracidium immediately after penetration, showing no recognizable changes in miracidial fine structure [see (8) for comparison]. During the first 3 hr after infection, the ciliated epidermal plates detach from the miracidium and curl up in the vicinity of the parasite (Fig. 2). As soon as the epidermal plates are discarded or loosened, epidermal ridges start to expand into the areas vacated by the plates, forming a new tegument by 5 hr postinfection (Fig. 3). Before the formation of a complete tegument, the parasite has neither

Explanations of Figs. 1 to 17.

Magnifications are indicated by a bar ($2\ \mu\text{m}$) besides each electron micrograph except for that in Fig. 10 which is equivalent to $1\ \mu\text{m}$. Letters demarcating structures are independent for each figure. All electron micrographs are from organisms in the tissues of *B. glabrata* at various ages which are indicated in the legends.

Fig. 1. Sagittal section of the anterior portion of a postmiracidium immediately after exposure (0 hr). At this stage of infection, little morphological changes are discernible. a, terebratorium; b, ciliated epidermal plate of the first tier; c, apical gland duct; d, apical gland; e, nerve; f, extra-CNS neuron; g, outer circular muscle; h, longitudinal muscle; i, processes of interstitial cells and j, host snail tissue.

Fig. 2. Postmiracidium (1 hr). Practically all epidermal plates are detached at this stage and the organism is devoid of protective surface covering except for the epidermal ridges. a, epidermal ridge; b, detached epidermal plate; c, muscle; d, cilia; e, ridge cyton; f, dilated RER cisternae; g, interstitial cyton; h, cilia within host histiocyte; i, host histiocyte and j, snail muscle.

Fig. 3. A young healthy mother sporocyst (5 hr) covered entirely by a newly formed tegument. a, tegument; b, lateral gland; c, circular muscle; d, peri-flame cell space holding many leptotriches; e, nerve; f, coiled septate desmosome covered over by tegument; g, histiocyte of host snail containing endocytosed cilia or ciliary plate of miracidium; h, host amoebocyte containing cilia and i, discarded ciliary plate in the snail tissues.

Fig. 4. Degenerating postmiracidium without tegument and surrounded by amoebocytes, histiocytes and fibroblasts (24 hr). Note that many cell nuclei show degenerative changes (clumping of heterochromatins) and cytoplasm appear granular. a, CNS containing neurons and neuropiles; b, degenerating nuclei; c, normal-appearing germ cell with prominent nucleolus in the nucleus; d, fibroblasts/histiocytes; e, amoebocytes containing many pseudopodia, some of which are closely attached to the "naked" parasite surface and f, fibroblast.

the characteristics of a miracidium (motility of cilia, swimming capacity) nor those of a mother sporocyst (a complete surface covering, i.e., tegument). This

Fig. 5. Expanding epidermal ridge and connecting cytoplasmic bridge to its cyton holding numerous membrane-bound vesicles (5 hr). a, ridge; b, connecting bridge; c, membrane-bound vesicles with electron dense contents; d, circular muscle of parasite; e, multiciliated sarcular sensory organ; f, lateral papilla; g, contents of vesicles discharged in the snail tissues and h, pseudopodia of amoebocyte.

Fig. 6. Tegument formation of young mother sporocyst (5 hr). a, tegument containing many vesicles; b, membrane-bound vesicle in the process of rupturing its membrane to discharge electron-dense contents; c, vesicle fusing its membrane to tegument membrane; d, electron dense vesicle contents discharged in the snail tissues and e, amoebocyte pseudopodia.

Fig. 7. Participation of membrane-bound vesicle in the formation of tegument in a young mother sporocyst (5 hr). Note a wide gap between pseudopodia and tegument. a, membrane-bound vesicle just ruptured its membrane; b, electron-dense contents of vesicle discharged in the host tissues; c, vesicle fusing to the tegument membrane; d, parasite muscle; e, basal lamina and f, host amoebocyte pseudopodia.

Fig. 8. Expansion of epidermal ridge at the early stage of tegument formation (1 hr). Note that the cytoplasm of histiocyte is closely attached to the expanding ridge surface without interference of vesicle contents (arrow). a, epidermal ridge containing numerous vesicles; b, vesicle; c, newly expanded ridge cytoplasm without vesicle; d, septate desmosome; e, loosened epidermal plate; f, epidermal ridge expanding beneath the plate; g, ridge cyton containing many vesicles; h, circular muscle; i, basal lamina; j, snail histiocyte containing endocytosed cilia and k, collagenous fibers invested on the surface of histiocyte.

young parasite is termed a postmiracidium. The postmiracidium is perhaps the most vulnerable stage of the intramolluscan parasite as all or parts are devoid of a protective surface covering. Thus, the internal tissues such as basal lamina and muscles are directly exposed to the host defense activities (Figs. 4 and 8) and may be subjected to the attack by the host amoebocytes (Fig. 8). Expansion of the epidermal ridges starts shortly after the epidermal plates are loosened or lifted off from the basal lamina but are still attached to the ridges by septate desmosomes. Sometimes the expansion of the epidermal ridges begins between the plates and the basal lamina (Fig. 8). The detached epidermal plates and cilia are quickly internalized by nearby host cells (Figs. 2, 3, 8 and 17). Some of these cells appear to be wandering amoebocytes (Figs. 14 and 17), while others have numerous collagenous fibers invested on their surface and appear as fibroblasts (Fig. 8). Both types of phagocytes are morphologically similar to normal free amoebocytes (Fig. 9). The latter cells, however, are fixed in the connective tissues and may be regarded as fixed phagocytes or snail histiocytes. This observation collaborates our earlier finding (5) that the connective tissues of *B. glabrata*, especially the loose connective tissues can mobilize hypertrophic amoebocytes under proper stimuli, suggesting that many cells in the connective tissues may be histiocytes. In those postmiracidia which fail to develop a new tegument the surface may be completely surrounded by amoebocytes and fibroblasts forming concentric layers (Fig. 4) and the organisms show extensive degeneration (Fig. 4).

Mother Sporocyst

By about 5 hr after penetration, the formation of a new tegument is usually complete and the parasite is enveloped by a thin syncytial tegument whose cytons are located beneath the muscle layers and are connected to the tegument by narrow cytoplasmic bridges (Fig. 5). The remarkably rapid formation of a new tegument by the postmiracidium appears to be accomplished with the help of membrane-bound vesicles present in the ridge cytons but rarely seen in the ridges during the miracidial stages [see (8), Figs. 17 to 20]. Soon after the epidermal plates are detached and the ridges begin to expand to form a new tegument, membrane-bound vesicles in the ridge cytons greatly increase in number and may also be seen in the ridges and their connecting cytoplasmic bridges, indicating movement of vesicles to ridges from their cytons (Figs. 5 and 7). The activities of ridge cytons are enhanced as indicated by the increased numbers of mitochondria and dilatation of rough endoplasmic reticulum (RER) cisternae (Fig. 2). As the vesicles move into ridges, many of them attach to the outer membrane of the

Fig. 9. Free amoebocyte near a young mother sporocyte (5 hr). a, nucleus; b, pseudopodia; c, mitochondrion and d, membrane-bound vesicle resembling that of parasite ridge cyton.

Fig. 10. Enlarged portion of Fig. 9. a, nucleus; b, lysosome-like vesicle with electron dense contents; c, Golgi's; d, mitochondrion and e, lamellar body.

ridges, and then the vesicles rupture to empty the electron-dense contents into the snail tissue (Figs. 6 and 7), while the vesicle membranes become a part of the membrane of the newly formed tegument (Figs. 6 and 7). None of the vesicles were seen free in the snail tissues. Thus the surface of the newly formed tegument become irregular (Figs. 5 to 7). In areas where the contents of the

Fig. 11. Retracted terebratorium being covered over by expanding epidermal ridge of young mother sporocyst (5 hr). a, profile of interlaced cytoplasmic expansion; b, cross section of apical gland duct lined with microtubules; c, circular muscle fibers forming a sphincter muscle; d, anterior-most epidermal ridge; e, membrane-bound vesicle; f, nerve containing neurosecretory vesicles and g, circular muscle.

vesicles are seen in the snail tissues, the pseudopodia of the amoebocytes are spread over the parasite tegument but are always separated from the tegument by a gap of 200 to 300 nm (Figs. 6 and 7). Therefore, it is postulated that the membrane-bound vesicles of ridge cytons are devices for storing membrane to be used in the rapid formation of a new tegument of the mother sporocyst. The contents of the vesicles may protect the early stages of the mother sporocyst from attack by host phagocytes by preventing their close attachment to the surface of the newly formed tegument. The abundance of RER, ribosomes, Golgi's and complex carbohydrates in the ridge cytons suggests that these cells are active in the production of exportable proteins. These activities are especially prominent shortly after the miracidium enters the snail host. The protective effect of the vesicle contents may also be indirectly visualized in that discarded epidermal plates and cil-

ia are quickly phagocytized by the host amoebocytes (Figs. 2, 3, 8, 14 and 17). However, the amoebocytes are unable to attach to the parasite tegument and are separated by a gap containing vesicle contents (Figs. 6 and 7). The vesicles resemble closely those present in amoebocytes (Figs. 9 and 10) and may be lysosome-like structures.

The septate desmosomes lining the peripheries of the epidermal plates and ridges usually detach from the plates but are intricately folded in a small area and are subsequently covered over rapidly by the expanding tegument (Fig. 3). The complex terebratorium with highly interlaced cytoplasmic folds on the surface (8) appears to keep its morphological integrity for over 5 hr postinfection (Fig. 11). In some organisms at 5 hr the terebratorium retracts into the body of the postmiracidium and is covered over by the expanding new tegument which flows from the anterior-most epidermal ridge (Fig. 11). By 72 hr after infection, the membrane-bound vesicles which were so abundant in the epidermal ridges during their expansion to form new tegument (Fig. 5) have practically disappeared from the tegument and ridge cytons (Fig. 13). The tegument cytons now contain numerous ribosomes, polysomes and large mitochondria, but few other organelles or complex carbohydrate particles. The septate desmosomes and sensory cilia which became embedded or enclosed within the new tegument begin to lose their identities by 72 hr after infection.

Microvilli of Mother Sporocyst

Occasional microvilli were recognized on the new tegument of young mother sporocysts fixed at 5 hr postinfection (Fig. 12). Extensive development of microvilli was first observed among organisms fixed at 24 hr after infection, and by 72 hr the development seemed almost complete (Fig. 13) and appeared little different from the microvilli of organisms fixed in 24 days postinfection. By this time mother sporocyst is completely surrounded by a space of ca. 3 μ m in which numerous collagenous fibers are present. The microvilli are frequently branched. The fine structure of microvilli of mother sporocyst resemble that of the microvilli of higher animals (13). The unit membrane of the microvilli is covered with a layer of dense fine fibrous material, ca. 10-15 nm thick and the intervillous spaces are filled with dense fine fibrous material. The cross section of each villus is more or less round in shape (Figs. 13 and 15).

Fig. 12. Young mother sporocyst (5 hr) showing early stages of microvilli formation. a, expanding tegument with many membrane-bound vesicles; b, early microvilli; c, septate desmosomes covered over by tegument; d, muscle and e, nerve to lateral papilla with clear neurosecretory vesicles.

Fig. 13. Young mother sporocyst (3 days) with well-formed microvilli. a, tegument devoid of vesicles but containing mitochondria and alpha glycogen particles; b, well-developed microvilli; c, still prominent circular muscle and d, collagenous fibers of snail tissues.

Germ Cells

Germ cells in the miracidium contain a relatively large nucleus enclosing a prominent nucleolus but that is poor in heterochromatins (8). The scarce granular cytoplasm is crowded with numerous ribosomes, many small mitochondria,

Fig. 14. Young mother sporocyst (5 hr) with an amoebocyte loosely attached on the irregular surface of tegument. Note that the amoebocyte is separated by a gap containing discharged electron dense contents of vesicles. a, tegument with many vesicles (b) and mitochondria (c); d, circular muscle; e, dislodged epidermal plate and cilia; f, amoebocyte with endocytosed cilia and g, pseudopodium.

Fig. 15. Mature mother sporocyst (24 hr) with complete complement of microvilli. a, tegument devoid of membrane-bound vesicles; b, microvilli; c, inconspicuous muscle and d, snail histiocyte.

rare Golgi's, a few RER and beta glycogen particles. These features are retained by the organism for the first 24 hr postinfection. Thereafter, the ribosomes are replaced by numerous polysomes and the scarce ribosomes are dispersed in small patches. By 48 hr, RER, Golgi's and mitochondria become more abundant.

Excretory System

The system consists of flame cells, excretory tubules, excretory vesicles and excretory pores in a miracidium (8). The excretory system appeared to maintain its morphological integrity during the observation period, i.e., up to 24 days postinfection (Fig. 16).

Musculature

Muscles maintain their prominence until about 3 days after infection (Fig. 13). However, the musculature is thinner and less prominent in organisms studied after 24 days (Fig. 15). It may be assumed that mother sporocysts retain some muscle fibers throughout their lives [5 to 7 weeks, (5)], to support the integrity of the tegument and the entire organism.

Interstitial Cells

These cells consist of irregularly shaped cytons located in the core areas behind the neural ring of the miracidium and have many long cytoplasmic processes extending throughout the miracidium (8). The major contents of their cytoplasm are alfa particles of complex carbohydrates and lipid droplets, suggesting their main functions are space-filling and energy storage (8). Changes of interstitial cell morphology were not observed in organisms younger than 5 hr. Thereafter alfa particles are mostly replaced by beta particles and they are more dispersed in the cytoplasm and stain less densely by lead citrate. By 24 hr after penetration, vacuoles of various sizes appear and the cytoplasm appears to lose its structural integrity, i.e., the contents are lost, the cytoplasm appears empty and the nucleus shows condensation of heterochromatins (Fig. 16). The changes suggest that space is being provided for the increasing numbers of dividing germ cells after the stored energy is exhausted.

Nervous System

Neurons within and without the neural ring (8) begin to show degenerative changes in organisms fixed in 24 hr after infection. The nuclei of these neurons exhibit clumping of heterochromatins and the neuropiles show less contents (Fig. 16).

Miscellaneous

Within 2 hr after the miracidium enters the snail host, many ciliated epidermal plates detach and are seen free in the snail tissues (Fig. 2). A few of the

Fig. 16. Yong mother sporocyst (24 hr) with normal-appearing flame cell, degenerating interstitial cell and neuron. a, flame cell; b, leptotriches in the peri-flame cell space; c, interstitial cyton with pyknotic nucleus; d, neuronal nucleus showing clumping of heterochromatins and e, neuropiles.

Fig. 17. Young amoebocyte near a mother sporocyst (5 hr) with endocytosed parasite epidermal plate/cilia. This amoebocyte is perhaps just transformed from a histiocyte containing no lysosome-like membrane-bound vesicles. a, nucleus; b, endocytosed epidermal plate/cilia; c, pseudopodia and d, histiocyte.

plates or cilia may be seen within host cells nearby. Endocytosis of discarded epidermal plates by host phagocytes reaches a peak by 5 hr postinfection but by 24 hr hardly any host cells containing discarded plates were observed indicating rapid digestion of the foreign bodies by snail phagocytes. The snail collagenous fibers have a thickness of 25 nm and numerous striations with periodicities of 50 nm, and are essentially similar in appearance to those of higher animals (13).

Comments

Although the fine structure of larval morphogenesis and of the host-parasite interactions between schistosomes and their hosts have been studied sporadically in the past (7,9-11, 14-22) systematic studies of the fine structure have not appeared in the literature. We have attempted to clarify the mechanism of the rapid formation of a new tegument during the miracidium-mother sporocyst transformation in vivo after the ciliated epidermal plates of the miracidium are discarded. We agree with most investigators (15,17-19) that the new tegument of the mother sporocyst is formed by the expansion of epidermal ridges. However, we have observed for the first time two important mechanisms of the formation of tegument and the survival of the parasite in the snail: [1] the participation of membrane-bound vesicles formed in the ridge cytons as ready-made membrane for the new tegument and [2] the contents of the vesicles which are discharged into snail tissues to prevent host amoebocytes from close apposition to the parasite surface. Failing to do so would make the "naked" parasite (postmiracidium) vulnerable to the host-defense mechanisms (Fig. 4). It was determined that only about 5 hr are required by the postmiracidium to completely form a new tegument in susceptible *B. glabrata*. Such rapid formation of a new tegument is an essential activity on the part of the parasite for survival in the snail host. Otherwise the parasite may quickly be destroyed by amoebocytes, as is seen in Fig. 4 where a degenerating organism without a tegument is being entirely enclosed by concentric fibroblasts [Type I tissue reaction of Pan (5)]. Southgate (23) and Wilson et al. (24) also reported that *Fasciola hepatica* postmiracidium forms a new tegument by utilizing preformed membranes but did not describe the mechanism.

During the miracidial stage, the membrane-bound vesicles are primarily located in the ridge cytons and in relatively small numbers, but in the postmiracidial stages shortly after infection, the numbers of these vesicles increase extensively and many are seen located in the ridges and their connecting bridges to the cytons, indicating their importance in relation to new tegument formation and to the survival of the postmiracidium. The vesicles are morphologically similar to the membrane-bound vesicles present in the snail amoebocytes near the postmiracidium (Figs. 9 and 10) which in turn resemble the lysosomes of vertebrate macrophages. It has been observed that hemolymph and/or amoebocytes of resistant *B. glabrata* may impair the survival of mother sporocysts of *S. mansoni* in vitro (9,10). These authors indicated also that amoebocytes from susceptible snails may be able to destroy mother sporocysts in

vitro in the presence of hemolymph from resistant snails. Our observations may explain why amoebocytes of susceptible *B. glabrata* are not capable of destroying the healthy postmiracidium/mother sporocyst. The contents of membrane-bound vesicles of the ridges discharged into the snail tissues may be responsible for the expression of the tegumental surface antigen of primary sporocysts in snail tissues seen by Dunn et al (21). The action of hemolymph from resistant snails may possibly neutralize the vesicle contents thus enabling amoebocytes from susceptible snails to closely appose to the mother sporocysts in vitro and destroy the parasite (9,10). Additional evidence that amoebocytes from susceptible snails are capable of destroying some mother sporocysts and postmiracidia was earlier reported (5) and the recent studies indicate that many miracidia die shortly after penetration and are quickly (within 24 hr) surrounded by amoebocytes and absorbed. A snail may be invaded by as many as 86 miracidia but the majority die shortly after penetration. Although the causes of death are not clear, it is likely that some postmiracidia do not mount a defense mechanism by releasing contents of membrane-bound vesicles to ward off the attack by amoebocytes. It would be interesting to pursue this aspect both in susceptible and resistant snails in the future. It is interesting to note that Yoshino et al. (22) showed that postmiracidia may grow to produce daughter sporocysts in the presence of snail embryonic cells (BGE), although these cells may internalize discarded ciliary plates and send out filopodia to interdigitate with microvilli of parasites. Perhaps these cells have not the capacity of "professional" phagocytes (amoebocytes) and are unable to destroy the parasites in vitro.

In my earlier publications (4,5,25), I indicated that some fibroblasts may transform into phagocytic free amoebocytes after proper stimulations by foreign bodies, such as schistosome larvae, microsporidia, etc. We now have more evidences to show that some fibroblasts in snail connective tissues behave like fixed macrophages of vertebrates as seen in Figs. 3 and 8. These cells are invested with numerous collagenous fibers on their surfaces, endocytose foreign bodies (cilia, epidermal plates, etc.) and apply themselves to the surface of the parasite tegument.

The excretory system appeared morphologically intact for the length of observation (24 days) and perhaps is maintained over the life of mother sporocyst without modification as similar organ systems are present throughout the life cycle stages of schistosomes.

It is interesting to note that most of ribosomes of the germ cells of the miracidium are replaced by polysomes in 24 hr after infection indicating that the germ cells are preparing for division which are seen 48 hr after penetration.

Although a small number of microvilli appear on the tegument in 5 hr, the number increased steadily thereafter and by 3 days postinfection they had become numerous reflecting the heightened absorptive activities of mother sporocysts during their rapid development. This is reflected in the disappearance of alpha particles of complex carbohydrates and later of beta particles in the newly formed tegument. Obviously the loss or exhaustion of these energy sources is compensated by the increased absorptive activities of numerous microvilli. Previous investigators (26,27) described two types of hemocytes (amoebocytes) in *B. glabrata* and *Bulinus* spp. We also observed two kinds of phagocytes in *B. glabrata*: free wandering amoebocytes containing many membrane-bound vesicles and large numbers of active Golgi's and abundant pseudopodia; and fixed amoebocytes invested on the surface with numerous collagenous fibers, many Golgi's but scarce pseudopodia and membrane-bound vesicles. The former type is probably equivalent to the granulocytes and the latter type to the hyalinocytes of Harris (26). In accordance with my earlier observations (5), I believe that the fixed amoebocytes (snail histiocytes) of *B. glabrata* may under proper stimulations transform into free wandering amoebocytes, becoming hypertrophic and producing numerous lysosome-like vesicles.

Finally, the roles of apical and lateral glands are difficult to interpret. The dense darkly stained droplets are still present in the cytons and ducts in large numbers in young mother sporocysts (5 hr). The droplets are discharged into snail tissues as amorphous granular material. Perhaps the material may help maintaining a space filled with snail hemolymph around mother sporocysts as we have observed previously (5).

SUMMARY: An electron microscopic study has been carried out to describe the transformation of the miracidium of *S. mansoni* into the mother sporocyst in the susceptible *B. glabrata* and the associated host-parasite interactions. The miracidium enters the snail host without morphological alterations. Within 3 hr after entering, all the ciliary epidemral plates of the miracidium are discarded. A new tegument is quickly formed by 5 hr postinfection by the expansion of epidermal ridges. The rapid formation of the new tegument reflects the participation of membrane-bound vesicles in the ridge cytons. The membranes of these vesicles become the new tegument membranes with the discharge of their

electron-dense contents into the snail tissues. The vesicular contents discharged into the tissues apparently prevent snail amoebocytes (phagocytes) from attachment to the parasite tegument and thus prevent their interference with the further development of the postmiracidium. If a postmiracidium fails to mobilize membrane-bound vesicles in the formation of tegument, the parasite becomes surrounded by closely attached concentric layers of fibroblasts formed by amoebocytes and histiocytes within 24 hr. The membrane-bound vesicles are present in small numbers in the ridge cytons of the miracidium and become numerous in the postmiracidium stage and with many migrate to the ridges through connecting bridges within 24 hr. By 3 days postinfection when extensive microvilli have formed on the tegument the vesicles have disappeared and are replaced by mitochondria, ribosomes and complex carbohydrate particles. Many fibroblasts in the snail connective tissues have phagocytic capacities and are regarded as snail tissue histiocytes or fixed amoebocytes that eventually may become hypertrophic and detached.

ACKNOWLEDGMENTS

I wish to thank Dr. Thomas H. Weller of this department for reading the manuscript and to Dr. Alfred A. Pan for preparing the manuscript.

REFERENCES

1. Weller, T. H. (1975): The increasing impact of schistosomiasis in a changing global ecology. p.21-29. In M. Marois (ed.), Development of chemotherapeutic agents for parasitic diseases, North Holland, Amsterdam.
2. Maldonado, J. F. and Acosta-Matienzo, J. (1947): The development of *Schistosoma mansoni* in the snail intermediate host, *Australorbis glabratus*. P. R. J. Publ. Hlth Trop. Med., 22, 331-373.
3. Ottolina, C. (1957): El miracidio del *Schistosoma mansoni*. Anatomia-citologia-fisiologia. Rev. Sani. Asist. Soc., 22, 1-435.
4. Pan, C. (1963): Generalized and focal tissue responses in the snail, *Australorbis glabratus*, infected with *Schistosoma mansoni*. Ann. New York Acad. Sci., 113, 475-485.
5. Pan, C. (1965): Studies on the host-parasite relationship between *Schistosoma mansoni* and the snail *Australorbis glabratus*. Am. J. Trop. Med. Hyg., 14, 931-976.
6. Schutte, C. H. J. (1974): Studies on the South African strain of *Schistosoma mansoni*. Part I: Morphology of the miracidium. S. Afr. J. Sci., 70, 299-302.

7. Meuleman, E. A. (1972): Host-parasite interrelationships between fresh water pulmonate *Biomphalaria pfeifferi* and the trematode *Schistosoma mansoni*. *Neth. J. Zool.*, 22, 355-427.
8. Pan, S. C. (1980): The fine structure of the miracidium of *Schistosoma mansoni*. *J. Inverteb. Pathol.*, 36, 307-372.
9. Bayne, C. J., Buckley, P. M. and Dewan, P. C. (1980): *Schistosoma mansoni*: Cytotoxicity of hemocytes from susceptible snail hosts for sporocysts in plasma from resistant *Biomphalaria glabrata*. *Exp. Parasitol.*, 50, 409-416.
10. Bayne, C. J., Buckley, P. M. and Dewan, P. C. (1980): Macrophage-like hemocytes of resistant *Biomphalaria glabrata* are cytotoxic for sporocysts of *Schistosoma mansoni* in vitro. *J. Parasitol.*, 66, 413-419.
11. Loker, E. S., Bayne, C. J., Buckley, P. M. and Kruse, K. T. (1982): Ultrastructure of encapsulation of *Schistosoma mansoni* mother sporocysts by hemocytes of juveniles of the 10-R2 strain of *Biomphalaria glabrata*. *J. Parasitol.*, 68, 84-94.
12. Chernin, E. (1957): A method of securing bacteriologically sterile snail (*Australorbis glabratus*). *Proc. Soc. Exp. Biol. Med.*, 96, 204-210.
13. Fawcett, D. W. (1981): *The Cell*. 2nd ed., Saunders, Philadelphia, PA.
14. Smith, J. E. and Chernin, E. (1974): Ultrastructure of young mother and daughter sporocysts of *Schistosoma mansoni*. *J. Parasitol.*, 60, 85-89.
15. Basch, P. F. and DiConza, J. J. (1974): The miracidium-sporocyst transition in *Schistosoma mansoni*: Surface changes in vitro with ultrastructural correlation. *J. Parasitol.*, 60, 935-941.
16. Koie, M. and Frandsen, F. (1976): Stereoscan observations of the miracidium and early sporocyst of *Schistosoma mansoni*. *Z. Parasitenk.*, 50, 335-344.
17. Meuleman, E. A., Lyaruu, D. M., Khan, M. A., Holtzmann, P. J. and Sminia, T. (1978): Ultrastructural changes in the wall of *Schistosoma mansoni* during the transformation of the miracidium into mother sporocyst in the snail host *Biomphalaria pfeifferi*. *Z. Parasitenk.*, 56, 227-242.
18. Samuelson, J. C., Quinn, J. J. and Caulfield, J. P. (1984): Hatching, Chemokinesis and transformation of miracidia of *Schistosoma mansoni*. *J. Parasitol.*, 70, 321-331.
19. Samuelson, J. C. and Caulfield, J. P. (1985): Role of pleated septate junctions in the epithelium of miracidia of *Schistosoma mansoni* during transformation to sporocysts in vitro. *Tissue Cell*, 17, 667-682.
20. Chiang, C. P. and Caulfield, J. P. (1988): *Schistosoma mansoni*: Ultrastructural demonstration of a miracidial glycocalyx that cross-reacts with antibodies raised against the cercarial glycocalyx. *Exp. Parasitol.*, 67, 63-72.
21. Dunn, T. S. and Yoshino, T. P. (1988): *Schistosoma mansoni*: Origin and expression of a tegumental surface antigen on the miracidium and primary sporocyst. *Exp. Parasitol.*, 67, 167-181.

22. Yoshino, T. P. and Laursen, J. R. (1995): Production of *Schistosoma mansoni* daughter sporocysts from mother sporocysts maintained in synxenic culture with *Biomphalaria glabrata* embryonic (BGE) cells. J. Parasitol., 81, 714-722.
23. Southgate, V. R. (1970): Observations on the epidermis of the miracidium and on the formation of the tegument of the sporocyst of *Fasciola hepatica*. Parasitology, 61, 177-190.
24. Wilson, R. A., Pullin, R. and Denison, J. (1971): An investigation of the mechanism of infection by digenetic trematodes: the penetration of the miracidium of *Fasciola hepatica* into snail host *Lymnaea truncatula*. Parasitology, 63, 491-506.
25. Pan, C. (1958): The general histology and topographic microanatomy of *Australorbis glabratus*. Bull. Museum Comp. Zool., Harvard Coll., 119, 237-299.
26. Harris, K. R. (1975): The fine structure of encapsulation in *Biomphalaria glabrata*. Ann. New York Acad. Sci., 266, 446-464.
27. Krupa, P. L., Lewis, L. M. and Vecchio, P. D. (1977): *Schistosoma haematobium* in *Bulinus guernei*: Electron microscopy of hemocyst interactions. J. Invertebr. Pathol., 30, 35-45.